Medium-sized cyclophanes. Part 58.¹ Synthesis and conformational studies of [2.n] metacyclophan-1-enes and [n.1] metacyclophanes

Takehiko Yamato,*" Koji Fujita" and Hirohisa Tsuzuki^b

- ^a Department of Applied Chemistry, Faculty of Science and Engineering, Saga University, Honjo-machi 1, Saga-shi, Saga 840-8502, Japan. E-mail: yamatot@cc.saga-u.ac.jp; Fax: +81 0952 28 8591
- ^b Tohwa Institute for Science, Tohwa University, 1-1-1 Chikushigaoka, Minami-ku, Fukuoka-shi, Fukuoka 815-8510, Japan

Received (in Cambridge, UK) 13th December 2000, Accepted 27th June 2001 First published as an Advance Article on the web 13th August 2001

A series of *syn*- and *anti*-[2.*n*]metacyclophan-1-enes and [2.*n*]metacyclophane-1,2-diols are prepared in good yields by a McMurry cyclization of 1,*n*-bis(5-acetyl-2-methoxyphenyl)alkanes. Interestingly, in the same coupling reaction in the absence of pyridine the pinacol rearrangement of [2.*n*]metacyclophane-1,2-diols to afford [*n*.1]metacyclophanes is observed, attributable to the TiCl₄ or acids generated from the McMurry reagent occurring during the cyclization reaction. In fact, protic acid- or Lewis-acid induced pinacol rearrangements of [2.*n*]metacyclophane ratio of the products is strongly governed by the number of the methylene bridges. The proportion of the rearrangement product increases with increasing length of the bridge. Conformational studies of [*n*.1]metacyclophanes as well as of [2.*n*]metacyclophane-1-enes in both solution and solid state are also described.

Introduction

Although [*n*.1]MCPs (MCP = metacyclophane) have been prepared by various workers, the synthetic routes used in previous methods were too long for practical purposes. Vögtle² reported the first synthesis of both [4.1]- and [5.1]MCP³ by use of the sulfone pyrolysis method. Later, Lin *et al.* succeeded in preparing the lower [3.1]homologue by a photochemical method.⁴ However, it was quite difficult to obtain sufficient amounts for further studies by this route.

In cyclophane chemistry the reductive coupling of carbonyl compounds by low-valent titanium, the McMurry reaction,⁵ has been used before by Mitchell *et al.*⁶ to synthesize cyclophanes with glycol units as bridges, by Tanner and Wennerström,⁷ and recently by Hopf⁸ and Grützmacher *et al.*⁹ for a cyclization of suitable dialdehydes to yield unsaturated cyclophanes. On the other hand, it is well known that acid-catalyzed pinacol rearrangement of 1,2-glycols affords the corresponding ketones;¹⁰ *e.g.*, 2,3-diaryl-2,3-dihydroxybutanes afford 3,3-diarylbutan-2-ones or 1,2-diaryl-2-methylpropan-1-ones *via* aryl or methyl shift, respectively. Thus, there is substantial interest in the systematic investigation of the pinacol rearrangement of [*n*.2]cyclophanes with glycol units at the ethylene bridges to afford either [*n*.1]cyclophanes by ring contraction or [*n*.2]cyclophane ketones.

In this paper, we describe a new preparative route for a series of anti-[2.n]MCP-1-enes and [2.n]MCP-1,2-diols using the low-valent-titanium-induced McMurry reaction. The latter compounds were further converted to [n.1]MCPs by pinacol rearrangement. Conformational studies of [n.1]MCPs as well as of [2.n]MCP-1-enes in both solution and solid state are also described.

Results and discussion

1,*n*-Bis(5-*tert*-butyl-2-methoxyphenyl)alkanes **1** have been prepared according to our previous papers.¹¹ The AlCl₃-MeNO₂catalyzed acetylation of compounds **1** with acetic anhydride or acetyl chloride at 20 °C led to an *ipso*-acylation reaction,¹² affording the desired 1,*n*-bis(5-acetyl-2-methoxyphenyl)alkanes **2** in good yield (Scheme 1).¹³



Scheme 1 Reagents and conditions: i, MeCOCl, $AlCl_3$ -MeNO₂, CH_2Cl_2 room temp. for 1 h.

1,2-Bis(5-acetyl-2-methoxyphenyl)ethane $2a^{13}$ was subjected to reductive coupling by the McMurry reaction following Grützmacher's procedure⁹ (Scheme 2). Although none of the desired [2.2]MCP-1-ene **3a** was observed, the dimer **6a** was obtained in 11% yield.

Interestingly, when pyridine was used in the present cyclization reaction, [2.2]MCP-1,2-diol **4a** was obtained in 10% yield, but the formation of the desired [2.2]MCP-1-ene **3a** was again not observed. This finding seems to be due to the much more strained structure of **3a** than that of diol **4a** containing the larger ring during formation of the unsaturated C=C linkage. Thus, during the McMurry reaction the intramolecular cyclization to afford **3a** might be quite difficult. Interestingly, when the reductive coupling reaction of **2a** was carried out in the presence of pyridine at room temperature, the yield of [2.2]MCP-1,2-diol **4a** increased from 10% to 30% along with trace amount of dimer **6a**. Results of the McMurry reaction for substrates **2a–e** are presented in Table 1.

DOI: 10.1039/b010075g

J. Chem. Soc., Perkin Trans. 1, 2001, 2089–2097 2089

This journal is $\ensuremath{\mathbb{C}}$ The Royal Society of Chemistry 2001

 Table 1
 McMurry reaction of 1,n-bis(5-acetyl-2-methoxyphenyl)alkanes 2

Run	Substrate	Number of methylene units, n	Conditions ^a	Products (% yield) ^b		
1	2a	2	A ^c	3a (0)	4a (0)	5a (0)
2	2a	2	В	3a (0)	4a (10)	5a (0)
3	2a	2	\mathbf{B}^{d}	3a (0)	4a (30)	5a (0)
4	2b	3	\mathbf{A}^{c}	3b (70)	4b (0)	5b (0)
5	2b	3	В	3b (80)	4 b (7)	5b (0)
6	2c	4	А	3c (69)	4c (13)	5c (0)
7	2c	4	В	3 c (83)	4c (15)	5c (0)
8	2d	5	А	3d (45)	4d (0)	5d (34)
9	2d	5	В	3d (77)	4d (8)	5d (4)
10	2e	6	А	3e (19)	4e (0)	5e (60)

^{*a*} Reaction conditions: A; The reaction was carried out in the absence of pyridine. B; The reaction was carried out in the presence of pyridine. ^{*b*} Isolated yields. ^{*c*} The dimers **6a** and **6b** were obtained in 11% and 2% yield, respectively. ^{*d*} The reaction was carried out at room temperature.



Scheme 2 (see Table 1). *Reagents and conditions:* i, TiCl₄–Zn–pyridine, THF, reflux for 60 h.

The structures of products, diol 4a and dimer 6a were determined on the basis of their elemental analyses and spectral data. Thus, Griffin et al. reported¹⁴ the structure of 1,2-dimethyl[2.2]MCP and assigned the exo-endo-arrangement. We have assigned the ¹H NMR signals of 4a in a similar fashion. In the ¹H NMR spectrum of 4a in CDCl₃ upfield shifts and the different chemical shifts for internal aromatic protons at δ 4.36 and 4.83 due to the ring current of the opposite aromatic ring were observed.^{15,16} These data strongly suggest that the structure of 4a is the anti-conformer. Furthermore, the two methyl groups show different chemical shifts at δ 1.30 and 1.83 each as a singlet. The four external aromatic protons were also observed as different chemical shifts at δ 6.81, 6.88, 7.13 and 7.40; the latter proton is in a strongly deshielding region of the oxygen atom of the endo-OH on the ethylene bridge. In contrast, one of the two internal aromatic protons was observed at lower field (δ 4.83) attributable to being in a strongly deshielding region of the oxygen atom of the exo-OH on the ethylene bridge. These data strongly support the suggestion that the two OH groups are in an endoand exo-arrangement and, therefore, anti-4a is found to be the trans-diol.

The mass spectral data for dimer **6a** (M^+ = 588) strongly support a cyclic dimeric structure. In the ¹H NMR spectrum of the tetramethoxy[2₄]MCP **6a**, protons of methyl groups, methoxy groups and ArCH₂CH₂Ar methylene protons each appear as a singlet at 27 °C. This behaviour indicates that the rate of conformational ring flipping of macrocycle **6a** is faster than the

NMR time-scale above this temperature. However, in dimer **6a** even at -60 °C in CDCl₃–CS₂ (1:3) the singlet signal of the ArCH₂CH₂Ar ethylene protons remains unsplit. These observations indicate the flexible structure of **6a**, similar to that of [2₄]MCP in spite of the introduction of two additional double bonds of the ethylene bridge.¹⁷

Similar McMurry cyclization of 1,3-bis(5-acetyl-2-methoxyphenyl)propane 2b carried out under the same reaction conditions afforded the desired [2.3]MCP-1-ene 3b in 70% yield along with a trace amount of dimer 6b. When pyridine was used in the present cyclization reaction, the yield of 3b increased to 80%. The same results were obtained in the case of the longer diacetvldiphenvlbutane 2c except for the formation of [4.2]MCP-11,12-diol 4c. Interestingly, the coupling reaction of 1,5-bis(5-acetyl-2-methoxyphenyl)pentane 2d with lowvalent titanium (TiCl₄/Zn) in the absence of pyridine led to the intramolecular cyclization affording 1,2-dimethyl[2.5]MCP-1ene 3d along with the [5.1]MCP 5d in 34% yield. The formation of the corresponding 1,2-diol 4d was not observed. This finding suggests that the pinacol rearrangement of 1,2-diol 4d catalyzed by TiCl₄ or acids generated from the McMurry reagent occurred during the cyclization reaction. In fact, when the same reaction was carried out in the presence of an excess of pyridine, the corresponding 1,2-diol 4d was obtained in 8% yield. In the case of 1,6-bis(5-acetyl-2-methoxyphenyl)hexane 2e the preference for formation of the rearrangement product 5e was observed. Thus, the [2.n]MCP-to-[n.1]MCP ratio of products is strongly governed by the number of methylene bridges. The portion of the rearrangement product increases with increasing length of the bridge. To the best of our knowledge the present rearrangements of [2.n]MCPs to the smallerring-size [n.1]MCPs by a ring contraction are the first such case in cyclophane chemistry. In fact Grützmacher et al. reported 18 a similar acid-catalyzed rearrangement of [2.1.2.1]paracyclophane-1,2-diol to afford the corresponding ketones by a pinacol rearrangement. No ring contraction was observed in spite of the much larger ring size than the present [2.n]MCP-1,2-diols (see Scheme 3).



Scheme 3 Reagents and conditions: i, H₂SO₄.

Table 2 Pinacol rearrangement of [2.n]MCP-1,2-diols 4c-e

	Run	Substrate	Number of methylene units, n	Conditions ^a	Products (% yield) ^b
	1	4c	4	Α	5c (45) 7c (5), 8c (7), 2c (8)
	2	4c	4	В	5c (33) ^c
	3	4d	5	А	5d (65)
	4	4d	5	В	5d (62)
	5	4e	6	А	5e (95)
	6	4 e	6	В	5e (96)
Reaction condit	tions: A; I ₂ –	HOAc, 1,4-dioxa	ne, reflux for 30 min. B; BF ₃ ·Et ₂ O, C	H ₂ Cl ₂ , room temper	ature for 30 min. ^b Isolated yields. ^c Intractable

mixture was also obtained.

As mentioned previously, synthesis of [n.1]MCPs have not been commonplace due to the lack of a general method such as that developed by Boekelheide for the [2.2]MCPs.¹⁹ As mentioned previously, Vögtle² reported the first synthesis of both [4.1]- and [5.1]-MCP by use of the sulfone pyrolysis method. Later, Lin et al. succeeded in preparing the lower [3.1]homologue by a photochemical method.⁴ However, the synthesis of [n.1]MCPs has not been established so far due to both the lowyield preparation and the unsuccessful attempts to prepare the lower [3.1]homologue except by photochemical synthesis. On the other hand, substituent effects on the pinacol rearrangement have been reported.^{10,20} Thus, there is substantial interest in pinacol rearrangement of [n.2] cyclophanes with glycol units at the ethylene bridge to afford either [n.1]cyclophanes by ring contraction or [n.2]cyclophane ketones. From the above results, the present pinacol rearrangement could be used for the preparation of novel [n.1]MCPs.

An attempted pinacol rearrangement of [4.2]MCP-11,12diol **4c** with I_2 -HOAc^{10b,21} in 1,4-dioxane under reflux for 3 h led to the desired [4.1]MCP **5c** in 45% yield along with the dehydration products **7c** and **8c** in 5 and 7% yield (Scheme 4,



Scheme 4 (see Table 2). Reagents and conditions: i, I_2 -HOAc, 1,4-dioxane, reflux for 30 min; BF₃·Et₂O, CH₂Cl₂ room temperature for 30 min.

Table 2). Furthermore, the ring cleavage at the 1,2-diol moiety occurred to form 1,4-bis(5-acetyl-2-methoxyphenyl)butane 2c in 8% yield under the conditions used. The same reaction carried out in the presence of BF₃·Et₂O afforded the rearrangement product 5c in 33% yield along with a mixture of intractable products. In contrast, in the case of treatment of [5.2]MCP-12,13-diol 4d and [6.2]MCP-13,14-diol 4e with I₂-HOAc in 1,4-dioxane the desired pinacol rearrangement products [5.1]MCP 5d and 5e were the main products, obtained in 65, 95% yield, respectively. No formation of dehydration product or ring-cleavage product was observed. Similar results were also obtained in the presence of BF₃·Et₂O as a catalyst. The yields of the rearrangement products 5 increase with the number of the methylene bridges. This result might be attributed to the decrease of ring strain in [*n*.1]MCPs.

The structures of pinacol rearrangement products **5c**–**e** and dehydration products **7c**, **8c** were determined on the basis of their elemental analyses and spectral data. In particular, in the case of [4.1]- and [2.4]-systems, for example, the mass spectral data for **5c** (M^+ = 338) and **7c**, **8c** (M^+ = 338 and 320) strongly support a cyclic structure. The reaction pathway in which the rearrangement products **5** would be formed *via* the protic-acid-catalyzed rearrangement of the 1,2-diols **4** is the same as that of the rearrangement of 4-methoxyacetophenone pinacol.¹⁰*b*

However, in the case of [2.2]MCP-1,2-diol **4a** and [3.2]MCP-10,11-diol **4b** with BF₃·Et₂O in dichloromethane the formation of the desired pinacol rearrangement products [2.1]MCP **5a** and [3.1]MCP **5b** was not observed under the reaction conditions used. Only the transannular cyclization products **9a** and **9b** were obtained, along with a large amount of a mixture of intractable products. When CF₃SO₃H was used in the present reaction, the transannular cyclization products **9a** and **9b** were obtained in almost quantitative yields (Scheme 5). These results



Scheme 5 Reagents and conditions: i, CF_3SO_3H , CH_2Cl_2 , room temperature for 30 min.

are consistent with the results obtained from the *anti*-[2.2]- and -[2.3]MCP-1-enes, which undergo transannular cyclization reactions under the acidic conditions or photoirradiation attributable to the proximity of the intra-annular positions and the release of the considerable strain energy.^{9a,9b,22}

Deacetylation of ketone 5c with molten potassium hydroxide gave the bridge-methyl-substituted [*n*.1]MCP 10c in 91% yield. Similar treatment of ketones 5d and 5e afforded the desired products 10d and 10e in 95 and 68% yield, respectively (Scheme 6).



Scheme 6 Reagents and conditions: i, KOH, 180 °C for 3 h.

Although the preparations of [n.1]MCPs were reported by various research groups,²⁻⁴ we have accomplished the first successful, convenient preparation of a series of [n.1]MCPs by



pinacol rearrangement *via* ring contraction of [n.2]MCPs **4** having a 1,2-diol unit on the ethylene bridge.

[2.*n*]MCP-1-enes adopt either a 'stair-case' *anti* conformation or a *syn* conformation with overlaying aromatic rings (Fig. 1).^{9b,15,23} Depending on the size of the bridges and on the presence of intra-annular substituents, the interconversion between the *syn* and *anti* conformers occurs by ring flipping.²⁴

The conformation of 1,2-dimethyl[2.n]MCP-1-enes 3 was readily apparent from their ¹H NMR spectrum. Thus, the internal aromatic proton shows an upfield shift (δ 5.69–6.77) due to the ring current of the opposite benzene ring.^{15,16} The ¹H NMR spectrum of 1,2-dimethyl[2.3]MCP-1-ene 3b at 27 °C showed a broad singlet of the intra-annular proton H_i at δ 5.69, apart from those at δ 6.72 and 7.03 for the other two protons on the aromatic rings. The methyl protons at the bridged double bond and the methoxy protons were observed each as a singlet at $\delta = 2.19$ and 3.83, respectively, and the protons of the trimethylene bridge generate a complicated signal pattern as expected for a rigid anti-[2.3]MCP-1-ene. The protons of the benzylic CH₂ group were observed as two multiplets centred at δ 1.95 and 2.92, which are further split by coupling with the protons of the central CH₂ group. This central CH₂ group was also observed as a multiplet centred at δ 1.69. This peak pattern ascribed to six chemically distinct protons of the propano bridge confirms the absence of an anti-anti interconversion which would exchange HA and HB of each CH₂ group. As the temperature of the solution of 3b in CDBr₃ is increased, the individual peaks of the benzyl protons merge and eventually a single peak is observed above 70 °C. This observation indicates that the rate of conformational ring flipping of 3b is faster than the NMR time-scale at this temperature. The energy barrier to the conformational ring flipping estimated from the coalescence temperature (T_c) is 15.6 kcal mol⁻¹. †

Similar findings were also observed in the ¹H NMR spectrum of 1,2-dimethyl[2.4]MCP-1-ene 3c at 27 °C, but the protons of the butane bridge give rise to two multiplets centered at δ 1.57 and 2.47, respectively, providing a fast interconversion of the two anti conformations of 3c by ring flipping. However, as the temperature of the solution in $CDCl_3-CS_2$ (1 : 3) is decreased, a single peak for the benzyl protons splits into a pair of doublets below -30 °C. The energy barrier to the conformational ring flipping estimated from the coalescence temperature (T_c) is 10.7 kcal mol⁻¹. This finding indicates a more flexible structure for 3c than of 3b, attributable to the larger cyclophane ring size. In spite of a decrease in temperature to -100 °C in CDCl₃-CS₂ (1:3), no change in the spectrum is observed for the [2.5]system 3d. The solution conformation of 1,2-dimethyl-[2.n]MCP-1-enes 3 is sensitive to the chain length of the bridge. The ring-inversion barriers determined by variable-temperature ¹H NMR decrease with increasing length of the bridges. Values for coalescence temperature and ring-flipping energy barriers are given in Table 3

Similarly, the conformation of [n.1]MCPs **5** and **10** was readily apparent from their ¹H NMR spectrum. For example, in the ¹H NMR spectrum of 11-acetyl-11-methyl-6,15-dimethoxy-[4.1]MCP **5c** in CDCl₃ upfield shifts and the different chemical

Table 3 Coalescence temperature (T_c) and ΔG_c^{\ddagger} for conformational ring inversion of [2.*n*]MCP-1-enes **3**, and [*n*.1]MCPs **5** and **10**

Compd.	Number of methylene units, <i>n</i>	<i>T</i> _c /°C ^{<i>a</i>}	$\Delta G_{ m c}^{\ddagger}$ (kcal mol ⁻¹) ^{<i>a</i>}
3b	3	70	15.6
3c	4	-30	10.7
3d	5	<-100	
5c	4	90	17.3
5d	5	<-100	
10c	4	90	17.3
10d	5	<-100	
^{<i>a</i>} T_c and ΔC	G_{c}^{\ddagger} were determined in CE	$OCl_3 - CS_2 (1:3)$	or CDBr ₃ , using

 T_c and ΔG_c were determined in $CDC1_3$ - CS_2 (1.5) of $CDB1_3$, using $SiMe_4$ as a reference.

shifts for internal aromatic protons at δ 6.33 and 6.56 due to the ring current of the opposite aromatic ring were observed.^{15,16} These data strongly suggest that the structure of 5c is the anti-conformer. Furthermore, the two methoxy groups show different chemical shifts at δ 3.80 and 3.81, each as a singlet. The four external aromatic protons were also observed as different chemical shifts at δ 6.66, 6.72, 6.91 and 7.21; the latter proton is in a strongly deshielding region of the oxygen atom of the acetyl group on the methylene bridge. Thus, 5c adopts a 'stair-case' anti conformation as known for other [2.3]MCP-1enes^{9b,23e} and [3.2]MCPs²⁴ having the same 11-membered ring. The ¹H NMR spectrum of 5c in CDCl₃ at room temperature exhibits a split pattern for the benzyl protons as two multiplets centred at δ 2.34 and 2.7. The central CH₂ group was also observed as two multiplets centred at δ 0.95 and 1.69. These findings suggest a rigid structure of [4.1]MCP 5c at this temperature. However, as the temperature of the solution in CDBr, is increased, a pair of multiplets of the benzyl protons merged into a single peak above 90 °C. The energy barrier to conformational ring flipping estimated from the coalescence temperature (T_c) is 17.3 kcal mol⁻¹. The energy barrier for the [4.1]-system 10c was also estimated to be the same [17.3 $(T_c = 90 \text{ °C})$ kcal mol⁻¹]. This energy is much lower than that of the rigid [2.2]MCP (>27 kcal mol⁻¹)^{25,26} and is similar to that for the homologous [3.2]MCPs (15.8–19.1 kcal mol⁻¹).²⁷ This finding indicates a more rigid structure for [4.1]MCP 5c and 10c (ca. 6.6 kcal mol⁻¹) than that for [2.4]MCP-1-ene 3c $(\Delta G_{\rm c}^{\ddagger} = 10.7 \text{ kcal mol}^{-1})$ attributable to the one methylene decrease. It was also found that the solution conformation of [n.1]MCPs is sensitive to the chain length of the bridges. The ring-inversion barriers determined by variable-temperature ¹H NMR dramatically decrease with increasing length of the bridge by one unit.

Usually, parent [n.2]MCPs in which intra-annular substituents are absent preferably adopt an *anti*-conformation.^{28,29} Similarly, in the case of the [4.1]MCP **5c** the *anti*-conformation was found to be favorable. The conformation of **5c** has also been confirmed by X-ray crystallographic analysis. Single colorless crystals of 6,14-dimethoxy-1,2-dimethyl[2.4]MCP-1ene **3c** and of 11-acetyl-11-methyl-6,15-dimethoxy[4.1]MCP **5c** suitable for X-ray crystallography were both obtained by recrystallization from methanol–chloroform (1 : 1).

The perspective ORTEP drawings of **3c** and **5c** are illustrated in Figs. 2 and 3, with the atom-numbering system. Compound **3c** crystallized in centrosymmetric orthorhombic space group *Pbcn* (No. 60) and is located about a two-fold axis, since this molecule has crystallographic C_2 symmetry. Therefore, the asymmetric unit contains one half of a molecule (Z = 4). Compound **5c** crystallized in monoclinic space group $P2_1/c$ (No. 14) (Z = 4).

The X-ray crystallography clearly shows that both conformations of **3c** and **5c** adopt the *anti* form in which two aromatic rings are in a non-planar chain form. In **3c** the selected bond lengths of C1–C2 and C2–C3 in the tetramethylene chains and

^{† 1} cal = 4.184 J.





Fig. 3 X-Ray structure of 11-acetyl-6,15-dimethoxy-11-methyl[4.1]-MCP **5c**. Thermal ellipsoids are drawn at the 50% probability level. For clarity all hydrogen atoms are omitted.

C7–C9 and C9–C10 in the ethylenic chains have suitable values, 1.530(4), 1.507(4), 1.503(4), and 1.509(5) Å, respectively, and the length of the double bond is 1.359(4) Å. The bond angles defined by C8–C7–C9 and C7–C9–C9' are 124.1(3) and 123.0(3)°, showing that **3c** displays a slightly distorted conformation. The two benzene rings of **3c** distort into a slight boat shape and slightly deviate from planarity; the dihedral angles of the plane defined by C3–C4–C6–C7 between the planes defined by C3–C4–C6–C7 and 1.58(3)°, respectively. The tetramethylene bridge chains do not display a fully extended *anti* conformation, probably to minimize the strain. Both methoxy groups on the benzene rings of **3c** point towards the out side, away from the tetramethylene bridge chain. This might contribute to avoiding steric crowding with hydrogens and carbons of the bridge chains.

Again, in **5c** both methoxy groups on the benzene rings, as in **3c**, point towards the outside, away from the tetramethylene bridge chain. In **5c** the two benzene rings also distort into a slight boat shape. The bond angle of the bridged methylene moiety, C13-C17-C4 is 106.6 (3)°, and the torsional twists of the two benzene rings (C1-C2-C3-C4-C5-C6 and C11-C12-

C13–C14–C15–C16) relative to the C13–C17–C4 plane are 33.2 (3)° and 44.2 (3)°, respectively.

The UV spectra of the [*n*.1]MCPs **10c**,**d** showed different absorption curves to that of the model acyclic compound, 1,1bis(4-methoxyphenyl)ethane **11**. A small bathochromic shift at 285 nm (log $\varepsilon_{max} = 3.55$) in comparison with that of **11** at 277 nm (log $\varepsilon_{max} = 3.62$) was observed for the [4.1]MCPs **10c**, which is ascribed to a transannular interaction between the two benzene rings and an increase in the strain of these systems.¹⁵ The same phenomenon is also observed in [5.1]MCP **10d** but the bathochromic shift is smaller than that of [4.1]MCPs **10c**.

Tetracyanoethylene (TCNE) complexes have often been used in studies on the relative π -base strength of various methylsubstituted benzenes.³⁰ The π -basicity of the donor molecules increases with an increase in the number of substituted methyl groups and/or stacking benzene rings and an increase in the face-to-face overlapping between aromatic nuclei.³¹ A solution of [4.1]MCP 10c and TCNE in CH₂Cl₂ present a purple color and the charge-transfer band at 606 nm (log $\varepsilon = 1.138$) was observed in its UV spectrum. This absorption is due to the formation of 1:1 charge-transfer complex among the electron donor, [4.1]MCP, and the electron acceptor, TCNE. The position of the absorption maximum and the shape of the absorption curve remain unchanged when a 4-12-fold excess of TCNE was added. However, the charge-transfer absorption band of the reference compound, 2,4-dimethylanisole 12, with TCNE was observed at 414 nm (log $\varepsilon = 2.073$). Such a red shift could be due to the through-space electronic interaction of the opposite uncomplexed benzene ring, which tends to work as a π -electron donor. In contrast to [4.1]MCP 10c, which exhibits the charge-transfer absorption band with TCNE at 606 nm (log $\varepsilon = 1.138$), a mixture of TCNE and [5.1]MCP **10d** exhibits an absorption peak at 510 nm (log $\varepsilon = 1.716$), while that of [6.1]MCP 10e is shifted to 501 nm (log $\varepsilon = 0.931$). Introduction of the one methylene inert to the tetramethylene bridge of 10c causes a larger blue shift as indicated by the 96 nm shift for the CT-band of 10d due to the decreased transannular π -electron donation from the non-complexed benzene ring to the complexed benzene ring.

In conclusion, a new synthesis of [2.n]MCP-1-enes and [2.n]MCP-1,2-diols by a McMurry cyclization has been developed. Protic acid- or Lewis acid-induced pinacol rearrangements of [2.n]MCP-1,2-diols can be applied to the synthesis of [n.1]MCPs. Further studies on the present novel ring contraction of [2.n]cyclophanes with glycol units at the ethylene bridge to afford [n.1]cyclophanes are now in progress.

Experimental

Mps (Yanagimoto MP-S1) and bps are uncorrected. NMR spectra were determined at 270 MHz with a Nippon Denshi JEOL FT-270 NMR spectrometer with SiMe₄ as an internal reference: *J*-values are given in Hz. IR spectra were measured for samples as KBr pellets or a liquid film on NaCl plates in a Nippon Denshi JIR-AQ2OM spectrophotometer. UV spectra were measured by a Shimadzu 240 spectrophotometer. Mass spectra were obtained on a Nippon Denshi JMS-01SA-2 spectrometer at 75 eV using a direct-inlet system through GLC.

Materials

The preparations of 1,n-bis(5-*tert*-butyl-2-methoxyphenyl)alkanes 1^{11} and 1,n-bis(5-acetyl-2-methoxyphenyl)alkanes 2a- $2c^{13}$ have been previously described. 1,1-Bis(4-methoxyphenyl)ethane 11 was prepared according to the literature procedure.¹⁰⁶

Acetylation of 1,*n*-bis(5-*tert*-butyl-2-methoxyphenyl)alkanes 1 with acetyl chloride in the presence of AlCl₃–MeNO₂

Typical procedure. To a solution of 1,5-bis(5-*tert*-butyl-2-methoxyphenyl)pentane **1d** (3.96 g, 10 mmol) and acetyl chlor-



ide (2.1 cm³, 30 mmol) in CH₂Cl₂ (80 cm³) was added a solution of aluminium chloride (5.94 g, 44.5 mmol) in nitromethane (10 cm³) at 0 °C. After the reaction mixture had been stirred at room temperature for 2 h, it was poured into ice-water (100 cm³). The organic layer was extracted with CH₂Cl₂ (50 $cm^3 \times 2$). The extract was washed with water (50 cm³ × 2), dried (Na₂SO₄), and concentrated. The residue was chromatographed over silica gel (Wako C-300, 300 g) with CHCl₃ as eluent to give crude 2d as a colorless solid. Recrystallization from hexanebenzene (1:1) gave 1,5-bis(5-acetyl-2-methoxyphenyl)pentane 2d (3.35 g, 80%) as prisms, mp 91–92 °C; v_{max}(KBr)/cm⁻¹ 1672 (C=O); δ_H(CDCl₃) 1.40–1.46 (2 H, m), 1.60–1.66 (4 H, m), 2.56 (6 H, s), 2.63 (4 H, t, J 7.8), 3.89 (6 H, s), 6.86 (2 H, d, J 8.8), 7.77 (2 H, d, J 2.4) and 7.82 (2 H, dd, J 8.8 and 2.4); m/z 368 (M⁺) (Found: C, 75.20; H, 7.75. C₂₃H₂₈O₄ requires C, 74.97; H, 7.66%).

Similarly, compound **2e** was prepared in 64% yield in the same manner as described above.

1,6-Bis(5-acetyl-2-methoxyphenyl)hexane 2e. This compound was obtained as prisms [from hexane–benzene (1 : 1)], mp 118–120 °C; ν_{max} (KBr)/cm⁻¹ 1667 (C=O); δ_{H} (CDCl₃) 1.35–1.42 (4 H, m), 1.51–1.64 (4 H, m), 2.55 (6 H, s), 2.62 (4 H, t, J 7.6), 3.88 (6 H, s), 6.87 (2 H, d, J 8.8), 7.77 (2 H, d, J 2.0) and 7.83 (2 H, dd, J 8.8 and 2.0); *m*/*z* 382 (M⁺) (Found: C, 75.52; H, 7.90. C₂₄H₃₀O₄ requires C, 75.36; H, 7.91%).

General procedure for the McMurry coupling reaction of 2

Typical procedure. The McMurry reagent was prepared from TiCl₄ [23.8 g (13.8 cm³), 125 mmol] and 18 g (275 mmol) of Zn powder in 500 cm³ of dry THF, under nitrogen. A solution of 1,3-bis(5-acetyl-2-methoxyphenyl)propane 2b (2.14 g, 6.3 mmol) and pyridine (22.8 cm³, 200 mmol) in dry THF (250 cm³) was added within 60 h to the black mixture of the McMurry reagent by using a high-dilution technique with continuous refluxing and stirring. The reaction mixture was refluxed for an additional 8 h, cooled to room temperature, and hydrolyzed with 10% aq. K_2CO_3 (200 cm³) at 0 °C. The reaction mixture was extracted with CH_2Cl_2 (200 cm³ × 3). The combined extracts were washed with water, dried with Na2SO4, and concentrated. The residue was chromatographed over silica gel (Wako C-300, 300 g) with hexane-benzene (1:1)benzene and CHCl₃ as eluents to give **3b** (1.55 g, 80%), **6b** (10 mg, 0.5%) and 4b (151 mg, 7%) each as a colorless solid. Recrystallization of crude 3b from methanol gave 6,13-dimethoxy-1,2-dimethyl-[2.3]metacyclophan-1-ene **3b** as prisms, mp 122–125 °C; $v_{\text{max}}(\text{KBr})/\text{cm}^{-1}$ 2911, 2836, 1598, 1498, 1458, 1252, 1237, 1142, 1029 and 803; δ_H(CDCl₃; 27 °C) 1.69 (2 H, br m), 1.95 (2 H, br m), 2.19 (6 H, s), 2.92 (2 H, br m), 3.83 (6 H, s), 5.69 (2 H, br s), 6.72 (2 H, br d) and 7.03 (2 H, br dd); $\delta_{\rm H}$ (CDCl₃-CS₂, 1 : 3; -40 °C) 1.59-1.77 (2 H, m), 1.90-2.05 (2 H, m), 2.20 (6 H, s), 2.86-2.98 (2 H, m), 3.85 (6 H, s), 5.68 (2 H, d, J 2.4), 6.73 (2 H, d, J 8.3) and 7.08 (2 H, dd, J 2.4 and 8.3); m/z 308 (M⁺) (Found: C, 81.88; H, 7.74. C₂₁H₂₄O₂ requires C, 81.78; H, 7.84%).

6,13,23,30-Tetramethoxy-1,2,18,19-tetramethyl[2.3.2.3]metacyclophane-1,18-diene **6b**. This compound was obtained as prisms [from hexane-benzene (1 : 1)], mp 234–236 °C; v_{max} (KBr)/cm⁻¹ 2949, 2832, 1606, 1500, 1463, 1244, 1033 and 890; $\delta_{\rm H}$ (CDCl₃) 1.34–1.47 (4 H, m), 2.14 (12 H, s), 2.22 (8 H, t, *J* 7.8), 3.72 (12 H, s), 6.63 (4 H, d, *J* 8.3), 6.65 (4 H, d, *J* 2.4) and 6.87 (4 H, dd, *J* 8.3 and 2.4); *m*/*z* 616 (M⁺) (Found: C, 81.84; H, 7.85. C₄₂H₄₈O₄ requires C, 81.78; H, 7.84%).

10,11-Dihydroxy-5,15-dimethoxy-10,11-dimethyl[3.2]metacyclophane **4b**. This compound was obtained as prisms [from hexane-benzene (2 : 1)], mp 216–218 °C (decomp.); v_{max} (KBr)/ cm⁻¹ 3400 (OH), 2941, 1606, 1505, 1250, 1108, 1032, 818; $\delta_{\rm H}$ (CDCl₃) 1.40 (3 H, s), 1.78 (3 H, s), 1.78–1.99 (2 H, m), 2.04– 2.18 (2 H, m), 2.30 (1 H, s, replaced by D₂O), 2.91–3.06 (2 H, m), 3.02 (1 H, br s, replaced by D₂O), 3.86 (3 H, s), 3.87 (3 H, s), 4.96 (1 H, d, J 2.4), 5.49 (1 H, d, J 2.4), 6.74 (1 H, d, J 8.8), 6.79 (1 H, d, J 8.8), 7.15 (1 H, dd, J 8.8 and 2.4) and 7.36 (1 H, dd, J 8.8 and 2.4); m/z 342 (M⁺) (Found: C, 74.64; H, 7.75. C₂₁H₂₆O₄ requires C, 74.63; H, 7.51%).

Similarly, compounds **6a**, **3c**–**3e**, **4a**, **4c**, **4d** and **5d**, **5e** were prepared in the same manner as described above. The yields are listed in Table 1.

1,2-Dihydroxy-6,12-dimethoxy-1,2-dimethyl[2.2]metacyclophane **4a**. This compound was obtained as prisms [from hexane–benzene (1 : 2)], mp 202 °C (decomp.); v_{max} (KBr)/cm⁻¹ 3505, 3381 (OH), 2960, 2923, 1598, 1502, 1252, 1136, 1081, 1023, 818, 618; $\delta_{\rm H}$ (CDCl₃) 1.30 (3 H, s), 1.54–1.72 (2 H, m), 1.83 (3 H, s), 2.08 (1 H, s, replaced by D₂O), 2.94 (1 H, s, replaced by D₂O), 3.47–3.58 (2 H, m), 3.88 (6 H, s), 4.36 (1 H, d, J 2.4), 4.83 (1 H, d, J 2.4), 6.81 (1 H, d, J 8.5), 6.88 (1 H, d, J 8.5), 7.13 (1 H, dd, J 8.5, 2.4) and 7.40 (1 H, dd, J 8.5 and 2.4); *m*/z 328 (M⁺) (Found: C, 73.34; H, 7.29. C₂₀H₂₄O₄ requires C, 73.15; H, 7.37%).

6,12,22,28-Tetramethoxy-1,2,17,18-tetramethyl-[2.2.2.2]metacyclophane-1,17-diene **6a**. This compound was obtained as prisms [from hexane–benzene (1 : 3)], mp 262–264 °C; v_{max} (KBr)/cm⁻¹ 2926, 2833, 1603, 1500, 1458, 1244, 1032 and 810; $\delta_{\rm H}$ (CDCl₃) 2.17 (12 H, s), 2.51 (8 H, s), 3.72 (12 H, s), 6.57 (4 H, d, J 8.8), 6.79 (4 H, dd, J 8.8 and 2.4) and 6.91 (4 H, d, J 2.4); *m*/z 588 (M⁺) (Found: C, 81.35; H, 7.55. C₄₀H₄₄O₄ requires C, 81.60; H, 7.53%).

6,14-Dimethoxy-1,2-dimethyl[2.4]metacyclophan-1-ene 3c. This compound was obtained as prisms (from methanol), mp 122–124 °C; v_{max} (KBr)/cm⁻¹ 2955, 2929, 2867, 2834, 1598, 1497, 1457, 1286, 1238, 1137, 1100, 1026 and 811; $\delta_{\rm H}$ (CDCl₃; 27 °C) 1.57 (4 H, br s), 2.20 (6 H, s), 2.47 (4 H, br s), 3.77 (6 H, s), 6.62 (2 H, d, J 2.4), 6.79 (2 H, d, J 8.4) and 7.08 (2 H, dd, J 8.4 and 2.4); $\delta_{\rm H}$ (CDCl₃–CS₂, 1 : 3; -70 °C) 0.60–0.87 (2 H, m), 1.32–1.55 (2 H, m), 1.90–2.19 (2 H, m), 2.15 (6 H, s), 2.62–2.84 (2 H, m), 3.76 (6 H, s), 6.51 (2 H, d, J 2.4), 6.73 (2 H, d, J 8.8) and 7.03 (2 H, dd, J 8.8 and 2.4); *m*/z 322 (M⁺) (Found: C, 82.17; H, 8.24. C₂₂H₂₆O₂ requires C, 81.95; H, 8.13%).

6,15-Dimethoxy-1,2-dimethyl[2.5]metacyclophan-1-ene 3d. This compound was obtained as prisms (from methanol), mp 105–107 °C; ν_{max} (KBr)/cm⁻¹ 2926, 1859, 1604, 1504, 1461, 1256, 1238, 1129, 1103, 1033 and 807; $\delta_{\rm H}$ (CDCl₃) 0.73–0.78 (2 H, m), 1.37–1.47 (4 H, m), 2.06 (6 H, s), 2.45–2.49 (4 H, m), 3.65 (6 H, s), 6.67 (2 H, d, J 8.8), 6.76 (2 H, d, J 2.4) and 6.92 (2 H, dd, J 8.8 and 2.4); *m*/z 336 (M⁺) (Found: C, 81.97; H, 8.46. C₂₃H₂₈O₂ requires C, 82.10; H, 8.39%).

6,16-Dimethoxy-1,2-dimethyl[2.6]metacyclophan-1-ene 3e. This compound was obtained as prisms (from methanol), mp 114–115 °C; v_{max} (KBr)/cm⁻¹ 2916, 2855, 1603, 1495, 1463, 1298, 1238, 1132, 1034 and 818; $\delta_{\rm H}$ (CDCl₃) 0.80–0.92 (4 H, m), 1.50–1.61 (4 H, m), 2.11 (6 H, s), 2.45–2.52 (4 H, m), 3.72 (6 H, s), 6.58 (2 H, d, J 8.8), 6.77 (2 H, d, J 2.4) and 6.80 (2 H, dd, J 8.8 and 2.4); *m*/*z* 350 (M⁺) (Found: C, 82.38; H, 8.68. C₂₄H₃₀O₂ requires C, 82.24; H, 8.63%).

11,12-Dihydroxy-6,16-dimethoxy-11,12-dimethyl[4.2]metacyclophane 4c. This compound was obtained as prisms [from hexane–benzene (2 : 1)], mp 186–189 °C (decomp.); v_{max} (KBr)/cm⁻¹ 3488 (OH); $\delta_{\rm H}$ (CDCl₃) 1.06–1.09 (2 H, m), 1.36 (6 H, s), 1.60–1.68 (2 H, m), 2.15–2.32 (2 H, m), 2.52 (2 H, s, replaced by D₂O), 2.81–2.99 (2 H, m), 3.83 (6 H, s), 6.04 (2 H, d, *J* 2.4), 6.85 (2 H, d, *J* 8.8) and 7.61 (2 H, dd, *J* 8.8 and 2.4); *m*/z 356 (M⁺) (Found: C, 73.94; H, 7.86. C₂₂H₂₈O₄ requires C, 74.13; H, 7.92%).

12,13-Dihydroxy-6,17-dimethoxy-12,13-dimethyl[5.2]metacyclophane 4d. This compound was obtained as prisms [from hexane-benzene (2 : 1)], mp 154–156 °C; v_{max} (KBr)/cm⁻¹ 3495 (OH); δ_{H} (CDCl₃) 1.11–1.16 (2 H, m), 1.34–1.43 (4 H, m), 1.54 (6 H, s), 2.06–2.10 (2 H, m), 2.40 (2 H, s, replaced by D₂O), 2.73–2.78 (2 H, m), 3.83 (6 H, s), 6.13 (2 H, d, J 8.8), 6.84 (2 H, d, J 2.4) and 7.57 (2 H, dd, J 8.8 and 2.4); *m*/z 370 (M⁺) (Found: C, 74.54; H, 8.19. C₂₃H₃₀O₄ requires C, 74.56; H, 8.16%). 12-Acetyl-7,16-dimethoxy-12-methyl[5.1]metacyclophane 5d. This compound was obtained as prisms (from hexane), mp 161–164 °C; ν_{max} (KBr)/cm⁻¹ 1708 (C=O); δ_{H} (CDCl₃) 1.20–1.26 (2 H, m), 1.40–1.50 (4 H, m), 1.84 (3 H, s), 2.19 (3 H, s), 2.48–2.54 (4 H, m), 3.83 (6 H, s), 6.72 (2 H, d, *J* 2.4), 6.78 (2 H, d, *J* 8.8) and 7.10 (2 H, dd, *J* 8.8 and 2.4); *m*/z 352 (M⁺) (Found: C, 78.07; H, 7.99. C₂₃H₂₈O₃ requires C, 78.38; H, 8.01%).

13-Acetyl-8,17-dimethoxy-13-methyl[6.1]metacyclophane 5e. This compound was obtained as pale yellow prisms (from hexane), mp 144–146 °C; ν_{max} (KBr)/cm⁻¹ 1711 (C=O); $\delta_{\rm H}$ (CDCl₃) 1.12–1.18 (4 H, m), 1.63–1.75 (4 H, m), 1.74 (3 H, s), 2.11 (3 H, s), 2.54–2.62 (4 H, m), 3.83 (6 H, s), 6.80 (2 H, d, J 8.8), 6.90 (2 H, d, J 2.9) and 7.08 (2 H, dd, J 8.8 and 2.9); m/z 366 (M⁺) (Found: C, 78.76; H, 8.25. C₂₄H₃₀O₃ requires C, 78.65; H, 8.25%).

Treatment of 4c with iodine-acetic acid in 1,4-dioxane

To a solution of 1,2-dihydroxy-6,14-dimethoxy-1,2-dimethyl-[2.4]metacyclophane **4c** (356 mg, 1 mmol) in a mixture of 1,4dioxane (20 cm³) and acetic acid (40 cm³) was added powdered iodine (100 mg, 0.39 mmol) at room temperature. After the reaction mixture had been stirred under reflux for 30 min, it was poured into ice–water (100 cm³). The organic layer was extracted with CH₂Cl₂ (50 cm³ × 2). The extract was washed successively with 10% aq. sodium thiosulfate (20 cm³ × 2) and water (50 cm³ × 2), dried (Na₂SO₄), and concentrated. The residue was chromatographed over silica gel (Wako C-300, 300 g) with hexane–benzene (1 : 1), benzene, benzene–CHCl₃, and CHCl₃ as eluent and recrystallization from methanol or hexane–benzene (1 : 1) to give **8c** (29 mg, 7%), **5c** (156 mg, 45%), **7c** (20 mg, 5%), and **2c** (30 mg, 8%) as colorless prisms, respectively.

6,16-Dimethoxy-11,12-dimethylene[4.2]metacyclophane 8c. This compound was obtained as prisms (from methanol), mp 101–102 °C; v_{max} (KBr)/cm⁻¹ 2929, 1500, 1244, 1119, 1029 and 902; δ_{H} (CDCl₃) 1.33–1.44 (4 H, m), 2.46–2.58 (4 H, m), 3.76 (6 H, s), 5.04 (2 H, d, *J* 2.2), 5.46 (2 H, d, *J* 2.2), 6.42 (2 H, d, *J* 2.0), 6.73 (2 H, d, *J* 8.3) and 6.99 (2 H, d, *J* 2.0 and 8.3); *m/z* 320 (M⁺) (Found: C, 82.04; H, 7.75. C₂₂H₂₄O₂ requires C, 82.46; H, 7.55%).

11-Hydroxy-6,16-dimethoxy-11-methyl-12-methylene[4.2]-

metacyclophane 7c. This compound was obtained as yellow oil; v_{max} (NaCl)/cm⁻¹ 3488 (OH); δ_{H} (CDCl₃) 1.07–1.44 (4 H, br m), 1.67 (3 H, br s), 2.18–2.75 (4 H, br m), 2.68 (1 H, s, replaced by D₂O), 3.79 (3 H, s), 3.84 (3 H, s), 4.86 (1 H, d, J 1.7), 5.44 (1 H, d, J 1.7), 5.57 (1 H, br d), 6.60 (1 H, br d), 6.77 (1 H, d, J 8.3), 6.91 (1 H, d, J 8.3), 6.99 (1 H, dd, J 8.3 and 2.0) and 7.42 (1 H, br d); *m*/*z* 338 (M⁺) (Found: C, 78.34; H, 7.67. C₂₂H₂₆O₃ requires C, 78.07; H, 7.74%).

11-Acetyl-6,15-dimethoxy-11-methyl[4.1]metacyclophane 5c. This compound was obtained as prisms [from hexane–benzene (1 : 1)], mp 126–128 °C; ν_{max} (KBr)/cm⁻¹ 1704 (C=O); δ_{H} (CDCl₃) 0.84–1.06 (2 H, m), 1.56–1.81 (2 H, m), 1.80 (3 H, s), 2.22 (3 H, s), 2.27–2.41 (2 H, m), 2.62–2.77 (2 H, m), 3.80 (3 H, s), 3.81 (3 H, s), 6.33 (1 H, br s), 6.56 (1 H, d, br s), 6.66 (1 H, d, J 8.3), 6.72 (1 H, d, J 8.3), 6.91 (1 H, dd, J 8.3 and 2.4) and 7.21 (1 H, dd, J 8.3 and 2.4); δ_{C} (CDCl₃) 24.15, 24.22, 24.35, 27.81, 55.04, 61.33, 107.34, 107.39, 123.87, 124.59, 126.78, 127.75, 133.92, 136.88, 137.87, 140.35, 156.62, 156.90 and 208.98; *m*/*z* 338 (M⁺) (Found: C, 78.28; H, 7.67. C₂₂H₂₆O₃ requires C, 78.07; H, 7.74%).

Similarly, compounds **5d** and **5e** were prepared in 65, 95% yields in the same manner as described above.

Treatment of 4c with BF₃-diethyl ether in dichloromethane

To a suspension of 4c (35.6 mg, 0.1 mmol) in CH_2Cl_2 (3 cm³)

was added BF₃·Et₂O (8.4 mg, 0.059 mmol) and the mixture was stirred at room temperature for 30 min. The reaction mixture was quenched by water (5 cm³), and extracted with CH₂Cl₂ (10 cm³ × 2). The combined extracts were washed successively with 5% aq. NaHCO₃ (10 cm³) and water (10 cm³ × 2), dried with Na₂SO₄, and concentrated to leave a residue, which was chromatographed over silica gel (Wako C-300, 300 g) with benzene as eluent, and recrystallized from hexane–benzene (1 : 1) to give **5c** (11 mg, 33%) as colorless prisms.

Similarly, compounds 4d and 4e were treated with BF_{3} -diethyl ether as described above to give 5d and 5e in 62, 96% yield, respectively.

Treatment of 4b with trifluoromethanesulfonic acid in dichloromethane

To a suspension of 4b (34 mg, 0.1 mmol) in CH₂Cl₂ (4 cm³) was added trifluoromethanesulfonic acid (0.02 cm³, 0.2 mmol) and the mixture was stirred at room temperature for 30 min. The cooled solution was quenched by water (5 cm³), and extracted with CH_2Cl_2 (10 cm³ × 2). The combined extracts were washed successively with 5% aq. NaHCO₃ (10 cm³) and water (10 cm³ \times 2), dried with Na₂SO₄, and concentrated to leave a residue, which was chromatographed over silica gel (Wako C-300, 300 g) with benzene as eluent and recrystallized from hexane-benzene (1:1) to give 9b (30 mg, 98%) as pale brown solid. Recrystallization from methanol to give 3,6dimethoxy-9,10-dimethyl-4,5-propanophenanthrene 9b as colorless prisms, mp 161–163 °C; v_{max}(KBr)/cm⁻¹ 2927, 1588, 1498, 1264, 1218, 1131, 1032 and 798; $\delta_{\rm H}$ (CDCl₃) 2.45–2.56 (2 H, m), 2.59 (6 H, s), 2.81-2.84 (4 H, m), 3.97 (6 H, s), 7.27 (2 H, d, J 8.8), 7.88 (2 H, d, J 8.8); m/z 306 (M⁺) [Calc. for C₂₁H₂₂O₂ (306.41): C, 82.32; H, 7.24. Found: C, 82.07; H, 7.20%].

Similarly, compound 2a was treated with trifluoromethanesulfonic acid as described above for 1 min to give 9a in 98% yield.

3,6-Dimethoxy-9,10-dimethyl-4,5-ethanophenanthrene 9a. This compound was obtained as pale brown prisms (from MeOH), mp 198–200 °C; δ_{H} (CDCl₃) 2.54 (6 H, s), 3.14 (4 H, s), 3.88 (6 H, s), 7.16 (2 H, d, *J* 9.3), 7.77 (2 H, d, *J* 9.3); *m/z* 292 (M⁺) [Calc. for C₂₀H₂₀O₂ (292.38): C, 82.16; H, 6.89. Found: C, 82.23; H, 6.73%].

Cleavage reaction of ketones 5 with KOH to give 10

Typical procedure. A mixture of 5c (33 mg, 0.1 mmol) and potassium hydroxide (1.6 g, 40 mmol) was heated at 180 °C for 3 h. The cooled melt was dissolved in water (10 cm³) and extracted with CH₂Cl₂ (20 cm³ \times 3). The combined extracts were washed with water, dried with Na₂SO₄, and concentrated. The residue was chromatographed over silica gel (Wako C-300, 300 g) with hexane-benzene (1:1) as eluent to give 10c (28 mg, 91%) as a colorless solid. Recrystallization of crude 10c from methanol gave 6,15-dimethoxy-11-methyl[4.1]metacyclophane **10c** as prisms, mp 148–149 °C; v_{max} (KBr)/cm⁻¹ 2862, 1498, 1254, 1235, 1110, 1034 and 806; $\delta_{\rm H}$ (CDCl₃) 0.74–0.94 (2 H, m), 1.50-1.69 (2 H, m), 1.63 (3 H, d, J 7.2), 2.22-2.34 (2 H, m), 2.62-2.82 (2 H, m), 3.75 (3 H, s), 3.76 (3 H, s), 4.12 (1 H, q, J 7.2), 6.00 (1 H, br s), 6.21 (1 H, br s), 6.68 (1 H, d, J 8.3), 6.73 (1 H, d, J 8.3), 7.11 (1 H, dd, J 8.3 and 2.4), 7.17 (1 H, dd, J 8.3 and 2.4); m/z 296 (M⁺) (Found: C, 81.22; H, 8.03. C₂₀H₂₄O₂ requires C, 81.04; H, 8.16%).

Similarly, compounds **10d** and **10e** were prepared in 95, 68% yield in the same manner as described above.

7,16-Dimethoxy-12-methyl[5.1]metacyclophane 10d. This compound was obtained as prisms (from MeOH), mp 105–106 °C; ν_{max} (KBr)/cm⁻¹ 2930, 1498, 1239, 1124, 1034 and 810; $\delta_{\rm H}$ (CDCl₃) 0.94–1.13 (2 H, m), 1.35–1.48 (4 H, m), 1.65 (3 H, d, *J* 7.3), 2.42–2.54 (4 H, m), 3.79 (6 H, s), 4.09 (1 H, q, *J* 7.3), 6.56 (2 H, d, *J* 2.2), 6.73 (2 H, d, *J* 8.3) and 7.12 (2 H, dd, *J* 8.3 and

 Table 4
 Crystallographic data and data-collection details for 6,14-dimethoxy-1,2-dimethyl[2.4]MCP-1-ene 3c and 11-acetyl-6,15-dimethoxy-11-methyl[4.1]MCP 5c

	3c	5c
Formula	C ₂₂ H ₂₆ O ₂	C ₂₂ H ₂₆ O ₃
FW	322.45	338.45
Size/mm	$0.35 \times 0.35 \times 0.10$	$0.30 \times 0.25 \times 0.20$
Crystal system	Orthorhombic	Monoclinic
Space group	<i>Pbcn</i> (No. 60)	$P2_1/c$ (No. 14)
alÅ	8.6418 (6)	14.8271 (13)
b/Å	10.318 (1)	12.8995 (1)
c/Å	19.776 (1)	9.5892 (5)
βl°		95.9810 (57)
V/Å ³	1763.4 (2)	1824.1 (2)
Ζ	4	4
$\rho_{\rm calc}/{\rm g~cm^{-3}}$	1.124	1.066
T/K	295	295
Radiation	$Cu-K_{a}$	Cu-K _a
λ/Å	1.541 84	1.541 84
μ/cm^{-1}	5.19	6.02
No. of reflections	2101	4061
Unique reflections	1813	3665
Observed reflections	1039	2489
R	0.059	0.083
R_{w}^{a}	0.077	0.121
<i>s</i> "	2.51	3.89
$^{a}\omega = 4(F_{o})^{2}/[(\sigma I_{o})2 + 0.0]$	$0016(F_{o})^{4}].$	

2.2); *m/z* 310 (M⁺) [Found: C, 81.31; H, 8.54. C₂₁H₂₆O₂ (310.44) requires C, 81.25; H, 8.44%].

8,17-Dimethoxy-13-methyl[6.1]metacyclophane 10e. This compound was obtained as prisms (from MeOH), mp 115–116 °C; v_{max} (KBr)/cm⁻¹ 2933, 1500, 1249, 1112, 1032 and 810; δ_{H} (CDCl₃) 0.96–1.05 (4 H, m), 1.52–1.68 (4 H, m), 1.61 (3 H, d, J 6.8), 2.50–2.54 (4 H, m), 3.82 (6 H, s), 4.10 (1 H, q, J 6.8), 6.57 (2 H, d, J 2.4), 6.79 (2 H, d, J 8.3) and 7.19 (2 H, dd, J 8.3 and 2.4); *m*/*z* 324 (M⁺) [Found: C, 81.53; H, 8.65. C₂₂H₂₈O₂ (324.47) requires C, 81.44; H, 8.7%].

Crystal data and refinement details for 6,14-dimethoxy-1,2dimethyl[2.4]metacyclophan-1-ene 3c ‡

The X-ray analysis was performed with the MolEN program package³² and the structure was solved uneventfully by direct methods (SIR 88).³³ Refinement was by full-matrix least squares and the 110 parameters refined were atomic coordinates, temperature factors (anisotropic for carbon atoms), scale factor, and secondary extinction coefficient. No corrections were made for absorption. The 12 independent non-hydrogen atoms were refined anisotropically. The 13 independent hydrogen atoms were located at calculated positions thermally fixed at $B_{iso} = 5.0$ Å and were included in refinement, but restrained to ride on the atoms to which they are bonded.

Crystal data and refinement details for 11-acetyl-6,15dimethoxy-11-methyl[4.1]metacyclophane 5c‡

The X-ray analysis was performed with the MolEN program package³² and the structure was solved uneventfully by direct methods (SIR 88).³³ Refinement was by full-matrix least squares and the 227 parameters refined were atomic coordinates, temperature factors (anisotropic for carbon atoms), scale factor, and secondary extinction coefficient. No corrections were made for absorption. The 25 independent non-hydrogen atoms were refined anisotropically. The 26 independent hydrogen atoms were located at calculated positions thermally fixed at $B_{iso} = 5.0$ Å² and were included in the refinement, but restrained to ride on the atoms to which they are bonded.

Crystallographic data and data-collection details for 3c and 5c are given in Table 4.

The refined non-hydrogen atomic coordinates, temperature factors (anisotropic for carbon atoms), scale factor, and secondary estimation coefficient are available, on request, from the Cambridge Crystallographic Data Centre.

References

- 1 For part 57 in the series, see T. Yamato, K. Fujita, T. Abe and H. Tsuzuki, New J. Chem., 2001, 25, 728.
- 2 M. Atzmüller and F. Vögtle, Chem. Ber., 1978, 111, 2547.
- 3 (a) N. Finch, C. W. Gemenden and B. P. Korzun, J. Org. Chem., 1976, 41, 2509; (b) N. Finch and C. W. Gemenden, J. Org. Chem., 1979, 44, 2804.
- 4 C. Lin, P. Singh, M. Maddox and E. F. Ullman, J. Am. Chem. Soc., 1980, 102, 3261.
- 5 (a) J. E. McMurry, M. P. Fleming, K. L. Kees and L. R. Krepski, J. Org. Chem., 1978, 43, 3255; (b) J. E. McMurry, Acc. Chem. Res., 1983, 16, 405; (c) J. E. McMurry, G. J. Haley, J. R. Matz, J. C. Clardy and G. V. Duyne, J. Am. Chem. Soc., 1984, 106, 5018; (d) J. E. McMurry, Chem. Rev., 1989, 89, 1513.
- 6 R. H. Mitchell and S. A. Weerawana, *Tetrahedron Lett.*, 1986, 27, 453.
- 7 D. Tanner and O. Wennerström, Acta Chem. Scand., Ser. B, 1983, 37, 693.
- 8 H. Hopf and C. Mlynek, J. Org. Chem., 1990, 55, 1361.
- 9 (a) H.-F. Grützmacher, E. Neumann, F. Ebmeyer, K. Albrecht and P. Schelenz, *Chem. Ber.*, 1989, **122**, 2291; (b) H.-F. Grützmacher and E. Neumann, *Chem. Ber.*, 1993, **126**, 1495; (c) H.-F. Grützmacher and G. Nolte, *Chem. Ber.*, 1994, **127**, 1157.
- 10 (a) W. E. Bachmann and J. W. Ferguson, J. Am. Chem. Soc., 1934, 56, 2081; (b) C. C. Price and G. P. Mueller, J. Am. Chem. Soc., 1944, 66, 634; (c) K. Sisido and H. Nozaki, J. Am. Chem. Soc., 1948, 70, 776.
- 11 (a) M. Tashiro, T. Yamato and G. Fukata, J. Org. Chem., 1978, 43, 1413; (b) T. Yamato, J. Matsumoto, K. Tokuhisa, M. Kajihara, K. Suehiro and M. Tashiro, Chem. Ber., 1992, 125, 2443.
- 12 (a) R. B. Moodie and K. Schofield, Acc. Chem. Res., 1976, 9, 287;
 (b) A. Fischer and R. Röderer, Can. J. Chem., 1976, 54, 3978;
 (c) A. Fischer and K. C. Teo, Can. J. Chem., 1978, 56, 258;
 (d) A. Fischer and K. C. Teo, Can. J. Chem., 1978, 56, 1758;
 (e) H. Suzuki, Synthesis, 1977, 217; (f) P. C. Myhre and M. Beug, J. Am. Chem. Soc., 1966, 88, 1568; (g) P. C. Myhre, M. Beug and L. L. James, J. Am. Chem. Soc., 1968, 90, 2105; (h) P. C. Myhre, M. Beug, M. Beug, K. S. Brown and B. Östman, J. Am. Chem. Soc., 1971, 93, 3452; (i) A. Fischer and R. Röderer, J. Chem. Soc., Chem. Commun., 1975, 798; (j) G. A. Olah and S. J. Kuhn, J. Am. Chem. Soc., 1964, 86, 1067; (k) W. Verboom, A. Durie, R. J. M. Egberink, Z. Asfari and D. N. Reinhoudt, J. Org. Chem., 1992, 57, 1313; (l) E. Kelderman, L. Derhaeg, G. J. T. Heesink, W. Verboom, J. F. J. Engbersen, N. F. van Hulst, A. Persoons and D. N. Reinhoudt, Angew. Chem., Int. Ed. Engl., 1992, 31, 1075.
- 13 T. Yamato, K. Maeda, H. Kamaimura, K. Noda and M. Tashiro, J. Chem. Res., 1995, (S) 310; (M) 1865.
- 14 R. W. Griffin, Jr, R. W. Baugman and C. E. Ramey, *Tetrahedron Lett.*, 1968, 5419.
- 15 (a) B. H. Smith, in Bridged Aromatic Compounds, Academic Press, New York, 1964; (b) F. Vögtle and P. Neumann, Angew. Chem., 1972, 84, 75; F. Vögtle and P. Neumann, Angew. Chem., Int. Ed. Engl., 1972, 11, 73; (c) F. Vögtle and P. Neumann, Synthesis, 1973, 85; (d) F. Vögtle and G. Höhner, Top. Curr. Chem., 1978, 74, 1; (e) P. M. Keehn and S. M. Rosenfield, Cyclophanes. Academic Press, New York, 1983, vol. 1; (f) F. Vögtle. Cyclophane Chemistry, Wiley, New York 1993.
- 16 (a) M. Tashiro and T. Yamato, J. Org. Chem., 1981, 46, 4556; (b) M. Tashiro and T. Yamato, J. Org. Chem., 1983, 48, 1461.
- 17 M. Tashiro, A. Tsuge, T. Sawada, T. Makishima, S. Horie, T. Arimura, S. Mataka and T. Yamato, J. Org. Chem., 1990, 55, 2404.
- 18 H.-F. Grützmacher, A. Mehdizadeh and A. Mülverstedt, *Chem. Ber.*, 1994, **127**, 1163.
- 19 V. Boekelheide, P. H. Anderson and T. A. Hylton, J. Am. Chem. Soc., 1974, 96, 1558.
- 20 (a) C. J. Collins, *Quart. Rev.*, 1960, **14**, 357; (b) W. E. Bachman and F. H. Moser, *J. Am. Chem. Soc.*, 1932, **54**, 1124; (c) M. R. Kegelman and E. V. Brown, *J. Am. Chem. Soc.*, 1954, **76**, 2711; (d) C. H. Beale and H. H. Hatt, *J. Am. Chem. Soc.*, 1932, **54**, 2405; (e) R. P. Zelinski and M. Jursich, *J. Am. Chem. Soc.*, 1956, **78**, 1015. See also Ref. 10a.

[‡] CCDC reference numbers 155635 and 155636. See http://www.rsc.org/ suppdata/p1/b0/b010075g/ for crystallographic files in .cif or other electronic format.

- 21 W. E. Bachman, Org. Synth., 1943, Coll. Vol. 2, 71.
- 21 (w) E. Bachman, Og. Symm. 1943, Coli. Vol. 2, 71.
 22 (a) N. L. Allinger, B. J. Gordon, H.-E. Hu and R. A. Ford, J. Org. Chem., 1967, 32, 2272; (b) T. Sato, E. Yamada, Y. Okamura, T. Amada and K. Hata, Bull. Chem. Soc. Jpn., 1965, 38, 1049; (c) M. Fujimoto, T. Sato and K. Hata, Bull. Chem. Soc. Jpn., 1967, 40, 600; (d) T. Sato, M. Wakabayashi, Y. Okamura, T. Amada and K. Hata, Bull. Chem. Soc. Jpn., 1967, 40, 2363; (e) T. Sato and K. Nishiyama, J. Chem. Soc., Chem. Commun., 1973, 220; (f) T. Sato, K. Nishiyama, S. Shimada and K. Hata, Bull. Chem. Soc. Jpn., 1971, 44, 2858; (g) S. Hayashi and T. Sato, Bull. Chem. Soc. Jpn., 1972, 45, 2360; (h) T. Sato and K. Nishiyama, J. Org. Chem., 1972, 37, 3254.
- 23 (a) M. Tashiro and T. Yamato, J. Am. Chem. Soc., 1982, 104, 3701;
 (b) T. Yamato, J. Matsumoto, S. Ide, K. Suehiro, K. Kobayashi and M. Tashiro, Chem. Ber., 1993, 126, 447; (c) T. Yamato, M. Sato, K. Noda, J. Matsumoto and M. Tashiro, J. Chem. Res., 1993, (S) 394 (M) 2601; (d) T. Yamato, J. Matsumoto, M. Sato, K. Noda and M. Tashiro, J. Chem. Soc., Perkin Trans. 1, 1995, 1299;
 (e) T. Yamato, K. Fujita, K. Futatsuki and H. Tsuzuki, Can. J. Chem., 2000, 78, 1089.
- 24 D. Krois and H. Lehner, Tetrahedron, 1982, 38, 3319.
- 25 (a) T. Sato, S. Akabori, M. Kainosho and K. Hata, Bull. Chem. Soc. Jpn., 1966, **39**, 856; (b) T. Sato, S. Akabori, M. Kainosho and K. Hata, Bull. Chem. Soc. Jpn., 1968, **41**, 218.
- 26 I. Gault, B. J. Price and I. O. Sutherland, Chem. Commun., 1967, 540.

- 27 R. W. Griffin, Jr. and R. A. Coburn, J. Am. Chem. Soc., 1967, 89, 4638.
- 28 (a) H. Förster and F. Vögtle, Angew. Chem., 1977, 89, 443; H. Förster and F. Vögtle, Angew. Chem., Int. Ed. Engl., 1977, 16, 429; (b) R. H. Mitchell, K. S. Weerawana and G. W. Bushnell, Tetrahedron Lett., 1984, 25, 907; (c) K. Böckman and F. Vögtle, Chem. Ber., 1981, 114, 1065; (d) M. F. Semmelhack, J. J. Harrisson, D. C. Young, Y. Guitierrez, S. Rafii and J. Clardy, J. Am. Chem. Soc., 1985, 107, 7508.
- 29 R. W. Hoffmann, Angew. Chem., 1992, 104, 1147; R. W. Hoffmann, Angew. Chem., Int. Ed. Engl., 1992, 31, 1124.
- 30 R. E. Merrifield and W. D. Phillips, J. Am. Chem. Soc., 1958, 80, 2778.
- 31 (a) R. Boschi and W. Schmidt, Angew. Chem., Int. Ed. Engl., 1973, 12, 402; (b) T. Sato and T. Takemura, J. Chem. Soc., Perkin Trans. 2, 1976, 1195; (c) S. Hayashi and T. Sato, Nippon Kagaku Zasshi, 1970, 91, 950; (d) D. J. Cram and R. H. Bauer, J. Am. Chem. Soc., 1959, 81, 5971; (e) L. A. Singer and D. J. Cram, J. Am. Chem. Soc., 1963, 85, 1080; (f) M. Sheehan and D. J. Cram, J. Am. Chem. Soc., 1969, 91, 3553; (g) M. Tashiro, T. Yamato, K. Kobayashi and T. Arimura, J. Org. Chem., 1987, 52, 3196; (h) T. Yamato, S. Ide, K. Tokuhisa and M. Tashiro J. Org. Chem., 1992, 57, 271. See also Refs. 22e, 22h.
- 32 MolEN: an International Structure Solution Procedure, Enraf-Nonius, Delft, The Netherlands, 1990.
- 33 M. C. Burla, M. Camalli, G. Cascarano, C. Giacovazzo, G. Polidori, R. Spagna and D. Viterbo, J. Appl. Crystallogr., 1989, 22, 389.