

View Article Online View Journal

ChemComm

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: X. Mou, Z. Xu, S. Wang, D. Zhu, J. Wang, W. Bao, S. Zhou, C. Yang and D. Zhang, *Chem. Commun.*, 2015, DOI: 10.1039/C5CC03979G.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/chemcomm

Journal Name



Au(I)-Catalyzed Rearrangement/Cyclization Cascade Toward theSynthesisof2-Substituted-1,4,5,6-tetrahydrocyclopenta[b]pyrrole^{†‡}

Accepted 00th January 20xx DOI: 10.1039/x0xx00000x

Received 00th January 20xx,

Xue-Qing Mou,^a Zheng-Liang Xu,^a Shao-Hua Wang,^{*ab} Dao-Yong Zhu,^a Jie Wang,^a Wen Bao,^a Shi-Jiang Zhou,^a Chao Yang^a and Di Zhang^a

www.rsc.org/

An Au(I)-catalyzed tandem reaction for the synthesis of 2phenyl-1,4,5,6-tetrahydrocyclopenta[b]pyrrole derivatives from 1-(1-hydroxy-3-phenylprop-2-yn-1-yl)cyclobutanol and primary amine or NH_4OAc has been developed to afford a series of polysubstituted pyrroles in moderate to good yields.

Since its first detection by F. F. Runge in 1834 and structure elucidation by von Bayer in 1870's, pyrrole has been considered as one of the most important heterocycles due to its broad utilities.¹⁻⁴ Up to now, pyrroles and its derivatives not only present in lot of biological contexts, but also represent as the key structure components of many bioactive molecules⁵⁻⁶ and some organic conducting materials,⁷⁻⁸ e.g., vitamin B12, atorvastatin and polypyrroles. Therefore, developing synthetic methodologies toward the construction of pyrrole skeleton is always of considerable interest to synthetic chemists and has led to several classical strategies like Hantzsch pyrrole synthesis,⁹ Knorr pyrrole synthesis,¹⁰ Paal-Knorr pyrrole synthesis,¹¹ Van Leusen reaction,¹² Barton-Zard reaction¹³ and Piloty-Robinson pyrrole synthesis.¹⁴ Especially, in recent years, under the promotion of transition metal catalyzed annulation reactions,15 a number of efficient methodologies toward the construction of pyrrole compounds have also been developed.¹⁶ Despite of the results achieved, it is still highly desirable to further explore alternative approaches toward the synthesis of such an important heterocycle.

As we all known, the Paal–Knorr pyrrole synthesis utilizing 1,4-diketones and amines as starting materials is one of the most commonly used method to construct multi-substituted pyrroles. Therefore, it is likely that a similar type of reaction could take place between a 1,4-diketone intermediate generated *in situ* and an amine within a tandem process.

This journal is C The Royal Society of Chemistry 20xx

Following this consideration and in connection with our interest in the transition metal catalyzed cyclization reactions and the development of synthetic methodologies toward the synthesis of heterocycles,^{17,18} we envisioned that a facile un efficient tandem reaction procedure to pyrrole structures from 1-(1-hydroxy-3-phenylprop-2-yn-1-yl)-cyclobutanol anr' primary amine could be achieved through a Meyer-Schuste Rearrangement¹⁹ / 1,2-migration²⁰ / Paal-Knorr cyclization cascade in the presence of proper catalyst, and would afford the products with a cyclopenta[b]pyrrole moiety, which is a common structural motif of many bioactive molecules, such as (+)-roseophilin.²¹ D-amino inhibitor²² ar 1 oxidase cyclooxygenase 2 inhibitors²³ (Scheme 1). To our knowledge, there is no report about such a reaction to give 1,4,5, tetrahydrocyclopenta[b]pyrrole via a tandem process, which is highly desirable in terms of reaction efficiency and ato a economy. Herein, we present a new and unique strategy to construct 1,4,5,6-tetrahydrocyclopenta[b]pyrrole via an Au() catalyzed tandem reaction (Scheme 1).



Scheme 1 Proposed synthetic strategy and related bioactive molecules.

In order to realize such a tandem reaction, we fir c investigated systematically a variety of reaction condition, such as Lewis acid catalysts, solvents, reaction temperature and time with 1-(1-hydroxy-3-phenylprop-2-yn-? yl)cyclobutanol (1a) and NH₄OAc (2a) as the model substrates. As summarized in Table 1, the desired reaction was attempte initially in the presence of Ph₃PAuCl (0.1 equiv) and AgOTf ($^{\circ}$

^{a.} School of Pharmacy & State Key Laboratory of Applied Organic Chemistry Lanzhou University, Lanzhou 730000, P. R. China. E-mail: wangshh@lzu.edu.cn;

^{b.} Key Laboratory of Drug Targeting and Drug Delivery System, Ministry of Education, Sichuan University, Chengdu, P. R. China.

[†]Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x ‡ Patent pending, Chinese patent application (No. 201510253806.6)

COMMUNICATION

equiv) in DCM at room temperature, as similar type of catalysts has been used in the Meyer-Schuster Rearrangement.²⁴ To our delight, the desired product 2phenyloctahydrocyclopenta[b]pyrrole (3a) was isolated in 26% yield (Table 1, entry 1). Inspired by this result, we next screened different solvents and reaction temperature. And it was found that nonpolar solvents usually gave better yields than polar ones. For example, when toluene, cyclohexane, nhexane, or *n*-heptane were employed, the reaction could afford the expected product in higher yields than other solvents, like THF, DCE, 1,4-dioxane and DCM (Table 1, entries 1-9). Among the nonpolar solvents screened, the use of nheptane could give the highest yield of 74% at 80 °C (Table 1, entry 9). And it was also observed that either decreasing or increasing temperature would cause a diminished yield of desired product (Table 1, entries 8-10). Next, different Lewis acid catalysts were further applied into the reaction, but none of them gave higher yields than the combination of Ph₃PAuCl and AgOTf (Table 1, entries 11 –21). Moreover, when Ph₃PAuCl or AgOTf alone was used as the catalyst, no desired product 3a was detected except the decomposition of the substrate 1a (Table 1, entries 18-19). Based on the information above, the loading of the catalyst, the reaction time were evaluated as well, and we were pleased to find that a lower amounts of catalyst (5 mol%) could catalyze the reaction well to give a yield of 75% by slightly prolonging the reaction time (Table 1, entry 22). Without catalyst, no conversion was observed (Table 1, entry 23), which indicated that the catalyst played a pivotal role in the reaction.

Table 1 Optimization of reaction conditions.^a

OH

	Ph			
	1a 2a		39	
			ou	h
Ent	Catalyst (mol%)	Solvent	Т	Yield
ry		00110111	(°C)	(%)
1	Ph₃PAuCl+AgOTf(10)	CH_2CI_2	30	26
2	Ph₃PAuCl+AgOTf(10)	DCE	60	30
3	Ph ₃ PAuCl+AgOTf(10)	THF	60	25
4	Ph ₃ PAuCl+AgOTf(10)	dioxane	60	35
5	Ph ₃ PAuCl+AgOTf(10)	<i>c</i> -hexane ^c	60	32
6	Ph ₃ PAuCl+AgOTf(10)	toluene	110	54
7	Ph₃PAuCl+AgOTf(10)	<i>n</i> -hexane	80	48
8	Ph₃PAuCl+AgOTf(10)	<i>n</i> -heptane	60	65
9	Ph ₃ PAuCl+AgOTf(10)	<i>n</i> -heptane	80	74
10	Ph ₃ PAuCl+AgOTf(10)	<i>n</i> -heptane	100	69
11	Ph ₃ PAuCl+CF ₃ CO ₂ Ag(10)	<i>n</i> -heptane	80	trace
12	AuCl+AgOTf(10)	<i>n</i> -heptane	80	trace
13	AuCl+ $CF_3CO_2Ag(10)$	<i>n</i> -heptane	80	n.d. ^d
14	Ph ₃ PAuCl+AgSbF ₆ (10)	<i>n</i> -heptane	80	42
15	IPrAuCl+AgOTf(10)	<i>n</i> -heptane	80	54
16	IPrAuCl+AgSbF ₆ (10)	<i>n</i> -heptane	80	46
17	SPhosAuCl+AgOTf(10)	<i>n</i> -heptane	80	23
18	Ph ₃ PAuCl(10)	<i>n</i> -heptane	80	n.d. ^d
19	AgOTf(10)	<i>n</i> -heptane	80	n.d. ^d
20	Cu(OTf) ₂ (10)	<i>n</i> -heptane	80	n.d. ^d

2 | J. Name., 2012, 00, 1-3

yl)cyclobutanol (0.5 mmol), ammonium acetate (5 equiv.), catalyst in solvent (1 mL) at 80 °C under Ar for 4 h. b Isolated yield. c Hexane = cyclohexane. d n.d. = not detected. e The reaction time was 6 h.

With the optimized reaction conditions in hand (Table 1 entry 22), we subsequently investigated the substrate scope c. such a transformation. Firstly, a series of amines wer subjected to the tandem reaction with compound 1a. A shown in Table 2, in most cases, the amines tested could proceed smoothly to transform into the polysubstitute ' pyrrole products in moderate yields (Table 2, 3a-3p). Amana them ammonium acetate as the amine source usually gave higher yield than aliphatic and aromatic amines. Among aromatic amine investigated, the electronic effect of the substituents on phenyl ring was not clearly observed. The presence of electronic neutral or deficient substituent group like bromo, chloro and nitro could give the expected products in moderate yield leading to additional possibilities for further derivatization (Table 2, 3h-3k). Besides, benzylamine and (4methoxyphenyl)methanamine also could go through to a reaction to give the desired products in moderate yield, whic , might simplify the synthesis process in case the pyrrol nitrogen needs pre-protection (Table 2, 3I-3m). Additionally when cyclic alkyl amine (2f, 2n), alkoxy substituted alkyl amin. (20) and propargyl amine (2s) were subjected to the tander reaction with compound 1a, they all afforded the desire. products. Encouraged by these results, we next applied the tandem reaction process to examine the scope of propargyl. alcohols. It was found that introducing an additional electron donating group like methyl and methoxy group on 🔤 aromatic ring of substrate 1a were well-tolerated, although they gave slightly lower yield than that of 1a (for example, product 3g and 3r, 3p and 3s). Also, replacing the phenyl grou , of compound 1a with a steric hindered tert-butyl group prove to be efficient under standard conditions, and therefore revealed that the aromatic ring of propargylic alcohri substrate 1 is not a prerequisite in this reaction. With respec to R² component in propargylic alcohols substrate 1, phenul was also compatible to give corresponding product, albeit in low yield. It should be noted that when the substrates with cyclopentanol moiety instead of cyclobutanol was applied to this tandem reaction, no desired products were obtair id, which might be attributed to the low reactivity of such a fi membered unit during the 1,2-migration step.

Table 2 Gold-catalyzed tandem synthesis of polysubstitute pyrrole.^a



This journal is C The Royal Society of Chemistry 20xx

Journal Name

Published on 12 June 2015. Downloaded by University of Birmingham on 15/06/2015 00:39:31

Journal Name

3d 56% 3a 74% **3b** 51% 3c 64% 3f 65% **3g** 58% **3h** 61% 3e 61% NO₂ **3i** 58% **3j** 57% 3k 60% 31 62% **3p** 57% 3m 54% 3n 22% 3o 43% 3q 52% 3r 43% **3s** 49% 3t 42% **3u** 68% 3v 59% 3w 35%

Reaction conditions: without other notifications, all of the reactions were performed with substrate 1 (0.5 mmol), R³NH₂ (3 equiv) or NH₄OAc (5 equiv), PPh₃AuCl (0.025 equiv), AgOTf (0.025 equiv) in *n*-heptane (1 mL) at 80 °C under Ar for 6h.

Based on above experimental results and related studies in this field, a plausible mechanism for this reaction was proposed with substrate 1a as an example (Scheme 2). Firstly, in the presence of Au(I) catalyst, a Meyer-Schuster rearrangement reaction would take place to give α,β unsaturated ketone intermediate I, which could go through an Au(I)-catalyzed 1,2-migration process to give 1,4-dicarbonyl intermediate II. Finally, a Paal-Knorr cyclization between intermediate II and amine would lead to the final product with a 1,4,5,6-tetrahydrocyclopenta[b]pyrrole skeleton.

RNH₂, PPh₃AuCl, AgOTf ò⊦ *n*-heptane, 80 °C юн 1a Meyer-schuster Paal-Knorr RNH_2 rearrangement Pyrrole Synthesis Au(I) HC но 1.2-rearrangement

Scheme 2 Plausible reaction mechanism.

COMMUNICATION

In summary, an Au(I)-catalyzed tandem reaction vie in xiol kine Meyer-Schuster rearrangement / 1,22migration 50,039a2 Knorr cyclization cascade has been successfully developed Through the use of this methodology, a series polysubstituted pyrroles were obtained in moderate to goc yields, which might provide a simple and versatile procedure for the synthesis of 1,4,5,6-tetrahydrocyclopenta[b]pyrro derivatives as well as related bioactivity screening.

Acknowledgements

This work was supported by the NSFC (No. 21202073 and 21472077), the fundamental research funds for the centr , universities (Izujbky-2014-k20, Izujbky-2015-k12), and the Openir Project of Key Laboratory of Drug Targeting and Drug Deliver System, Ministry of Education (Sichuan University).

Notes and references

3

- T. Eicher, Journal für Praktische Chemie/Chemiker-Zeitung, 1 1998, 340, 487.
- C. T. Walsh, S. Garneau-Tsodikova and A. R. Howard-Jone 2 Nat. Prod. Rep., 2006, 23, 517.
 - C. Schmuck and D. Rupprecht, Synthesis, 2007, 3095.
- H. Fan, J. Peng, M. T. Hamann and J.-F. Hu, Chem. Rev., 2008, 4 108, 264.
- 5 (a) A. Gossauer, Pyrrole. In Houben-Weyl; Thieme: Stuttgart. 1994; E6a/1, p 556; (b) A. Fürstner and H. Weintritt, J. Ar Chem. Soc., 1998, 120, 2817; (c) D. L. Boger, C. W. Boyce, N A. Labroli, C. A. Sehon and Q. Jin, J. Am. Chem. Soc., 1999 121, 54; (d) B. Sayah, N. Pelloux-Léon and Y. Vallée, J. Or Chem., 2000, 65, 2824; (e) J.-H. Liu, Q.-C. Yang, T. C. W. Mak and H. N. C. Wong, J. Org. Chem., 2000, 65, 3587.
- 6 G. W. Gribble, In Comprehensive Heterocyclic Chemistry II; A. R. Katritzky, C. W. Rees, E. F. V. Scriven, Eds.; Pergamon Pres Oxford, 1996; Vol. 2, p 207.
- 7 L. Groenendaal, E. W. Meijer and J. A. J. M. Vekemans, in Electronic Materials: The Oligomer Approach, Wiley-VC Verlag GmbH, 2007, pp. 235.
- (a) V. M. Domingo, C. Alemán, E. Brillas and L. Juliá, J. Org. 8 Chem., 2001, 66, 4058; (b) A. G. MacDiarmid, Synth. Met., 1997, 84, 27; (c) K. Tamao, S. Ohno and S. Yamaguchi, Chen Commun., 1996, 1873.
- 9 (a) A. Hantzsch, Chem. Ber., 1890, 23, 1474-1476; (b) F. Feis Chem. Ber., 1902, 35, 1537.
- 10 A. H. Corwin. Heterocyclic Compounds. 1950, 1, p. 287
- 11 (a) C. Paal, Chem. Ber., 1884, 17, 2756-2767; (b) L. Knor Chem. Ber., 1884, 17, 2863.
- 12 O. H. Oldenziel, D. Van Leusen and A. M. Van Leusen, J. Org. Chem., 1977, 42, 3114.
- 13 J. J. Li, Heterocyclic Chemistry in Drug Discovery. New Yor. Wiley, 2013.
- 14 (a) O. Piloty, Chem. Ber., 1910, 43, 489; (b) G. M. Robir on and R. Robinson, J. Chem. Soc., Trans., 1918, 113, 639.
- 15 For selected reviews, (a) J. Ye and S. Ma, Acc. Chem. Res., 2014, 47, 989; (b) C. Obradors and A. M. Echavarren, Cher Commun., 2014, 50, 16-28; (c) C. Obradors and A. N Echavarren, Acc. Chem. Res., 2014, 47, 902; (d) R. Shintan Bull. Chem. Soc. Jpn., 2012, 85, 931; (e) D. L. J. Broere and E. Ruijter, Synthesis-Stuttgart, 2012, 44, 2639; (f) Y. Deng, A. A. Persson and J.-E. Backvall, Chemistry-a European Journa, 2012, 18, 11498; (g) M. M. Heravi and E. Hashemi, Monats Chem., 2012, 143, 861; (h) K. Tanaka and Y. Tajima, Eur. Org. Chem., 2012, 3715; (i) Y. Yamamoto, Chem. Rev., 2012,



COMMUNICATION

112, 4736; (j) B. Chattopadhyay and V. Gevorgyan, *Angew. Chem. Int. Ed.*, 2012, **51**, 862; (k) A. Marinetti, H. Jullien and A. Voituriez, *Chem. Soc. Rev.*, 2012, **41**, 4884; (l) C. Aubert, L. Fensterbank, P. Garcia, M. Malacria and A. Simonneau, *Chem. Rev.*, 2011, **111**, 1954; (m) Z.-X. Yu, Y. Wang and Y. Wang, *Chemistry-an Asian Journal*, 2010, **5**, 1072; (n) J. Montgomery, *Acc. Chem. Res.*, 2000, **33**, 467; (o) A. Padwa, *Molecules*, 2001, **6**, 1; (p) B. M. Trost and M. J. Krische, *Synlett*, 1998, 1; (q) M. Malacria, *Chem. Rev.*, 1996, **96**, 289.

- 16 For selected examples after 2010, (a) Q. Li, A. Fan, Z. Lu, Y. Cui, W. Lin and Y. Jia Org. Lett., 2010, 12, 4066; (b) Q. Li, J. Jiang, A. Fan, Y. Cui and Y. Jia Org. Lett., 2011, 13, 312; (c) B. Li, N. Wang, Y. Liang, S. Xu and B. Wang, Org. Lett., 2013, 15, 136; (d) B. V. S. Reddy, M. R. Reddy, Y. G. Rao, J. S. Yadav and B. Sridhar, Org. Lett., 2013, 15, 464; (e) L. Wang and L. Ackermann, Org. Lett., 2013, 15, 176; (f) Z. Chen, B. Lu, Z. Ding, K. Gao and N. Yoshikai, Org. Lett., 2013, 15, 1966; (g) T. Miura, K. Hiraga, T. Biyajima, T. Nakamuro and M. Murakami, Org. Lett., 2013, 15, 3298; (h) F. Shi, R. Y. Zhu, X. Liang and S. J. Tu, Adv. Synth. Catal., 2013, 355, 2447; (i) G. Lonzi and L. A. Lopez, Adv. Synth. Catal., 2013, 355, 1948; (j) S. Peng, L. Wang, J. Huang, S. Sun, H. Guo and J. Wang, Adv. Synth. Catal., 2013, 355, 2550; (k) J. Liu, Z. Fang, Q. Zhang, Q. Liu and X. Bi, Angew. Chem. Int. Ed., 2013, 52, 6953; (I) M. Gao, C. He, H. Chen, R. Bai, B. Cheng and A. Lei, Angew. Chem. Int. Ed., 2013, 52, 6958; (m) V. Estevez, M. Villacampa and J. C. Menendez, Chem. Commun., 2013, 49, 591; (n) L. Meng, K. Wu, C. Liu and A. Lei, Chem. Commun., 2013, 49, 5853; (o) C. Wang, C. Dong, L. Kong, Y. Li and Y. Li, Chem. Commun., 2014, 50, 2164; (p) J. C. Liu, X. T. Zhang, H. Peng, H. F. Jiang and B. L. Yin, Adv. Synth. Catal., 2015, 357, 727; (q) C.-E. Kim, S. Park, D. Eom, B. Seo and P. H. Lee, Org. Lett., 2014, 16, 1900; (r) T. Li, X. Xin, C. Wang, D. Wang, F. Wu, X. Li and B. Wan, Org. Lett., 2014, 16, 4806; (s) H. Ueda, M. Yamaguchi, H. Kameya, K. Sugimoto and H. Tokuyama, Org. Lett., 2014, 16, 4948; (t) S. Gujarathi, X. G. Liu, L. Song, H. Hendrickson and G. R. Zheng, Tetrahedron, 2014, 70, 5267; (u) Y. Tokimizu, M. Wieteck, M. Rudolph, S. Oishi, N. Fujii, A. S. K. Hashmi and H. Ohno, Org. Lett., 2015, 17, 604.
- 17 Y.-Q. Zhang, D.-Y. Zhu, B.-S. Li, Y.-Q. Tu, J.-X. Liu, Y. Lu and S.-H. Wang, *J. Org. Chem.*, 2012, **77**, 4167.
- (a) D.-Y. Zhu, Z. Zhang, X.-Q. Mou, Y.-Q. Tu, F.-M. Zhang, J.-B. Peng, S.-H. Wang and S.-Y. Zhang, *Adv. Synth. Catal.*, 2015, 357, 747-752; (b) L. Xu, S.-H. Wang, H. Li, *Chin. J. Org. Chem.*, DOI 10.6023/cjoc201501022.
- 19 K. H. Meyer and K. Schuster, Chem. Ber., 1922, 55, 819.
- 20 E. Zhang, C.-A. Fan, Y.-Q. Tu, F.-M. Zhang and Y.-L. Song, J. Am. Chem. Soc., 2009, **131**, 14626.
- 21 A. Fürstner, Angew. Chem. Int. Ed., 2003, 42, 3582.
- M. L. R. Heffernan, Q. K. Fang, R. J. Foglesong, S. C. Hopkins,
 C. O. Ogbu, M. Soukri, K. L. Spear, U.S. Pat. Appl.
 Publ., US 20100029737, 2010.
- 23 B. Zhong, M. Li, P. R. China Pat., CN1955163A, 2007.
- 24 R. S. Ramón, N. Marion and S. P. Nolan, *Tetrahedron*, 2009, 65, 1767.

Journal Name

View Article Online DOI: 10.1039/C5CC03979G