



## Boradipyrrromethenecyanines derived from conformationally restricted nuclei

Viktor P. Yakubovskiy, Mykola P. Shandura, Yuriy P. Kovtun\*

*Institute of Organic Chemistry, National Academy of Sciences of Ukraine, 5 Murmanska Street, 02660 Kyiv, Ukraine*

### ARTICLE INFO

#### Article history:

Received 13 October 2009

Received in revised form

19 January 2010

Accepted 20 January 2010

Available online 2 February 2010

#### Keywords:

BODIPY

Boradiaza-*s*-indacene

Polymethine dye

Merocyanine

Long-wavelength dye

### ABSTRACT

A series of *meso*-polymethine-substituted 4,4-difluoro-4-bora-3a,4a-diaza-*s*-indacenes were synthesized based on the reaction of *meso*-methyl borondipyrrromethene with a number of hemicyanine derivatives. The dyes obtained exhibited a weak short-wavelength absorption, a strong long-wavelength absorption and weak fluorescence. Upon protonation, the long-wavelength band disappeared while the intensity of the short-wavelength band increased markedly. The properties of the dyes were closely related to those merocyanine dyes rather than borondipyrrromethenes.

© 2010 Elsevier Ltd. All rights reserved.

### 1. Introduction

The dyes derived from borondipyrrromethene (BODIPY) have attracted considerable attention over the past two decades on account of their excellent thermal, chemical and photochemical stability, high molar absorption coefficient, high fluorescence quantum yield, general insensitivity to solvent polarity and pH, large two-photon cross-section for multiphoton excitation, lack of ionic charge and good solubility [1–4]. However, most BODIPY compounds have absorption maxima below 600 nm. As long-wavelength dyes are important [5,6] there are many synthetic approaches for modifying the BODIPY system so as to red shift their absorption maxima.

One of the most promising approaches to the modification of BODIPY is its peripheral functionalization with conjugated chromophores. A wealth of studies have been devoted to the synthesis of  $\alpha$ -substituted BODIPYs **A** [7–11] including derivatives **B** with dimethylamino auxochromes [12–17] (see Scheme 1). The styryl moieties at the  $\alpha$ -positions cause a noticeable bathochromic shift in  $\lambda_{\text{max}}$  of the dye. Introduction of additional 4-dialkylamino substituents into **A** (structure **B**) results in even more pronounced spectral changes, with bathochromic shifts around 80 nm. Under acidic conditions only the diethylamino groups are protonated, leading to the optical properties that are very similar to those of **A**, where Ar = Ph. As known, in cyanine dyes the chain carbon atoms


are the very positions which can be protonated, unlike terminal groups [18,19]. Therefore, in case of **B** dialkylamino groups cannot be referred to as true “end groups” as they are considered in the ideology of polymethine dyes. Thus, these NR<sub>2</sub> groups in structures **B** provide considerable spectral effects but do not change the nature of colour of the BODIPY core. Derivatives **C** *meso*-functionalized with *p*-dimethylaminophenyl groups [20–25] show much the same absorption as the unsubstituted BODIPY nucleus and, like **B** compounds, drastically reduced fluorescence.

We have recently established that *meso*-substitution of BODIPY with more complex heterocycle-terminated polymethine chains results in the formation of merocyanine dyes (see compounds **5a–7a** in Scheme 3) and thus qualitatively change the very nature of the BODIPY absorption [26]. These dyes are mainly characterized by two electronic transitions, one long-wavelength polymethinic and the other short-wavelength localized in the borondipyrrromethene moiety. In terms of conventional cyanine nomenclature, we have introduced the term boradipyrrromethenecyanines to describe these new substituted BODIPY systems.

The present work is directed towards the synthesis of dyes structurally similar to **5a–7a** and containing the bridged borondipyrrromethene residue **4** as one end group. It is known that the bridging of  $\alpha$ -aryl substituents causes significant bathochromic shifts in the absorption of BODIPYs as well as an increase in their molar absorption coefficients and fluorescence quantum yields [27,28]. Thus, our concern is with the effects caused by these structural changes on the spectra of the corresponding boradipyrrromethene merocyanines.

\* Corresponding author.

E-mail address: [kovtun@ioch.kiev.ua](mailto:kovtun@ioch.kiev.ua) (Y.P. Kovtun).



Scheme 1.

## 2. Results and discussion

Compound **4** was synthesized similarly to its analogue **4a** [26]. Pyrrole **1** [29] was condensed with triethyl orthoacetate in the presence of *p*-toluenesulfonic acid to give the tosylate salt of dipyrromethene **2** (see Scheme 2).

Unlike the previously described unbridged analogue [26], this salt appears to be quite stable in solutions; the  $^1\text{H}$  NMR spectrum in  $\text{DMSO}-d_6$  exhibits no signs of dissociation. One can therefore assume that the ethylene bridging of the dipyrromethene moiety makes this unit more basic. The absorption spectrum of **2** in dichloromethane likewise suggests its stability, whereas a significantly reduced molar absorption coefficient in more polar DMF points to the partial dissociation of the salt (see Table 1). When treated with triethylamine in ethanol, salt **2** is converted to compound **3** characterized by the  $^1\text{H}$  NMR-detectable prototropic tautomerism of ethylene **3a** and *meso*-methyl dipyrromethene form **3b**. The former is observed in highly polar  $\text{DMSO}-d_6$ , and the latter in weakly polar  $\text{CDCl}_3$ . Salt **2** was boiled in chloroform with boron trifluoride etherate in the presence of Hünig's base to produce **4** in high yield (90%); **4** is also obtainable from **3a** under the same conditions.

The *meso*-methyl group of compound **4** appears to be quite reactive in condensations with (2-acetanilidovinyl) heterocyclic derivatives (see Scheme 3). A series of dyes **5–8** containing, respectively, 2-indolenyl, 2-benzothiazolyl, 2-quinolyl, and 4-quinolyl as the second heterocyclic end residue (in the order of increasing basicity) were obtained. The corresponding cyanine condensations proceeded quite smoothly to furnish the dyes desired in good yields (45–65%). It should be noted that even dye **8** with the very basic 4-quinolyl end nucleus was prepared successfully, whereas its unbridged analogue proved inaccessible via the conventional procedure [26].

As expected, compound **4** is more deeply coloured than its unbridged counterpart **4a**; a noticeable batho- and hyper-chromic


effect in absorption is accompanied by a significant increase in the fluorescence quantum yield (see Scheme 2). At the same time, the spectral properties of dyes **5–8** are much like those of analogous compounds **5a–7a** (see Fig. 1 and Table 1). They also exhibit two absorption bands: long-wavelength having the polymethinic origin and short-wavelength assigned to the electronic transition in the borondipyrromethene moiety. The latter is red-shifted due to bridging, so that the two bands are closer to each other than for dyes **5a–7a**. In DMF solutions, the absorption of merocyanines **5–7** is shifted only 20–33 nm to longer wavelengths and a little more intense relative to respective dyes **5a–7a** (see Table 1). This unexpectedly slight difference in spectral behaviour between bridged and unbridged dyes is attributable to the competition between the batho- and hyper-chromic effect of bridging, on the one hand, and the opposite trend caused by the less polymethinic character of the bridged dyes, on the other hand. Indeed, bridging makes the borondipyrromethene nucleus, as well as its boron-free precursor, less electron-withdrawing; at the same time, it is known that any decrease in the donor-acceptor strength of end groups brings the corresponding neutral merocyanines further from the so-called cyanine limit [30] and hence blue-shifts their absorption.


The borondipyrromethene local absorption peak which appears for indolenine dye **5** as a shoulder of the polymethinic band becomes increasingly separate for dyes **6–8** as the basicity of the second end group rises. The band separation is caused by an increasing red shift of the long-wavelength polymethinic absorption as a result of the approach to the cyanine limit with rising donor-acceptor difference of dye end groups. Accordingly, the long-wavelength and short-wavelength bands of dye **8** (much weaker than those of the other dyes in the series) are practically resolved from each other.

For all the dyes obtained, just as for their unbridged analogues **5a–7a**, protonation with strong acid results in a complete disappearance of the long-wavelength band and an intensity increase of the short-wavelength band. This effect is exemplified by the absorption of protonated dye **6** in Fig. 1.

As already mentioned, dipyrromethene **2** and hence its boron compound **4** are weaker electron-acceptors than their unbridged analogues. Accordingly, the  $^1\text{H}$  NMR signal of the methyl group in compound **4** is shifted upfield from that in **4a** (cf. the respective chemical shifts of 2.64 and 2.76 ppm). Likewise, dyes **5–7** exhibit upfield shifts (of about 0.2 ppm) for the proton resonances of the polymethine chain in comparison to respective analogues **5a–7a**.

Solvatochromism of the merocyanines obtained is illustrated by the absorption of dye **5** which demonstrates batho- and

Scheme 2. Reagents: (i)  $\text{CH}_3(\text{OEt})_3$ ,  $\text{TosOH} \cdot \text{H}_2\text{O}$ ; (ii)  $\text{Et}_3\text{N}$ ,  $\text{EtOH}$ ; (iii)  $\text{BF}_3 \cdot \text{Et}_2\text{O}$ ,  $\text{Et}(i\text{-Pr})_2\text{N}$ .



Scheme 3.

hyper-chromic effects with increasing solvent polarity (see Fig. 2). Solvent effects decrease as the basicity of the second heterocyclic nucleus rises: dyes **5–8** show the respective red shifts of 38, 21, 10, and 9 nm on going from dichloromethane to DMF. This tendency is consistent with the fact that the bridged dyes with the second end group of high basicity are closer to the so-called cyanine limit and hence less solvent-sensitive.

The comparative analysis of the two series shows that for the bridged dyes with weakly and moderately basic end residues **5** and **6** solvatochromic red shifts are larger than those for the unbridged **5a** and **6a** (Table 1). This suggests that dyes **5–6** are further from the cyanine limit and hence more solvatochromic than **5a–6a** due to the decreased electron-withdrawing ability of the bridged borondipyrromethene end nucleus. As the basicity of the second heterocyclic nucleus rises further, the solvatochromic effects for bridged and unbridged dyes are equalized (**7** and **7a**).

Compound **4**, being a typical BODIPY, exhibits a narrow fluorescence band with a quantum yield of 0.86. Polymethine-substituted dyes **5–8** as well as their analogues **5a–7a** are weakly fluorescent, particularly in polar solvents; the protonated dyes are likewise non-fluorescent.

### 3. Experimental

Electronic absorption spectra were recorded on a Shimadzu UV-3100 spectrophotometer.  $^1\text{H}$  (300 MHz, 25 °C,  $\text{Si}(\text{CH}_3)_4$  as internal standard) and  $^{19}\text{F}$  NMR (188 MHz,  $\text{CFCl}_3$  as internal standard) spectra were obtained with a Varian VXR-300 instrument. LC/MS spectra were recorded using a liquid chromatography/mass spectrometric system consisting of an Agilent 1100 Series high-performance liquid chromatograph equipped with a diode-matrix and an Agilent LC/MSD SL mass-selective detector. Fluorescence spectra were recorded on Solar CM 2203 fluorescence spectrophotometer. Relative fluorescence quantum yields ( $\phi$ ) were determined relative to indodicarbocyanine [31] (for compound **4**) and pentamethyne dioxaborinate [32] (for compounds **5–8**).

#### 3.1. Compound 2

A mixture of pyrrole **1** (1.69 g, 10 mmol), *p*-toluenesulfonic acid (1 g, 5.25 mmol), and triethyl orthoacetate (5.5 mL, 30 mmol) was stirred at room temperature for 20 min. After diluting with ethyl acetate (30 mL) the reaction mixture was allowed to stand for 20 min and the product was filtered off. Yield 1.78 g, 66%. M.p.

**Table 1**  
Optical properties of prepared compounds.

Dye	$\text{CH}_2\text{Cl}_2$			DMF		
	$\lambda_{\text{abs}}$ , nm ( $\epsilon \cdot 10^{-3}$ , $\text{M}^{-1} \text{cm}^{-1}$ )	fwhm <sup>a</sup> , $\text{cm}^{-1}$	$\lambda_{\text{em}}$ , nm ( $\phi_f$ )	$\lambda_{\text{abs}}$ , nm ( $\epsilon \cdot 10^{-3}$ , $\text{M}^{-1} \text{cm}^{-1}$ )	fwhm <sup>a</sup> , $\text{cm}^{-1}$	$\lambda_{\text{em}}$ , nm ( $\phi_f$ )
<b>2</b>	606 (108)	875		603 (47)	967	
<b>4</b>	621 (134)	625	631 (0.86)	617 (113)	685	
<b>5</b>	664 (103)	2334	723 (0.011)	702 (107)	2762	n.d. <sup>b</sup>
<b>6</b>	699 (119)	2569	741 (0.005)	724 (123)	1627	n.d. <sup>b</sup>
<b>7</b>	741 (145)	1170	n.d. <sup>b</sup>	751 (140)	1026	n.d. <sup>b</sup>
<b>8</b>	797 (49)	1163	n.d. <sup>b</sup>	806 (56)	1056	n.d. <sup>b</sup>
<b>5a</b> <sup>c</sup>	661 (108)	1631	712 (0.006)	682 (116)	1634	736 (<0.001)
<b>6a</b> <sup>c</sup>	679 (110)	2289	726 (0.005)	693 (105)	1558	729 (<0.001)
<b>7a</b> <sup>c</sup>	710 (125)	1432	745 (0.004)	718 (128)	1334	754 (<0.001)

<sup>a</sup> Full width at half-maximum height.

<sup>b</sup> Too weak to be reliably detectable.

<sup>c</sup> The data taken from Ref. [26].




Fig. 1. Absorption spectra of dyes **4** and **5–8** in DMF ( $C = 1 \times 10^{-5}$  mol/l).

174–176 °C.  $^1\text{H}$  NMR (DMSO, 300 MHz):  $\delta = 2.30$  (s, 1H,  $\text{CH}_3$ ) 2.88–2.94 (m, 7H), 3.02 (q, 4H,  $\text{CH}_2$ ), 7.12–7.19 (m, 3H), 7.46–7.57 (m, 9H), 8.19 (d,  $J = 3.6$  Hz, 2H, ArH), 12.68 (s, 1H, NH). LC-MS:  $m/z$  536 ( $[\text{M} + \text{H}]^+$ ). Anal. calcd. for  $\text{C}_{33}\text{H}_{30}\text{N}_2\text{SO}_3$ : C, 74.16; H, 5.62; N, 5.24. Found: C, 74.37; H, 5.4; N, 7.51.

### 3.2. Compound 3

To a suspension of salt **2** (1.07 g, 2 mmol) in ethanol (10 mL), triethylamine (0.4 g, 4 mmol) was added, followed by stirring the reaction mixture at room temperature for 30 min. The resulting precipitate was filtered off and washed with ethanol. Yield 0.54 g, 74%.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300 MHz):  $\delta = 2.54$  (s,  $\text{CH}_3$ ), 2.87 (q,  $J = 7$  Hz, 4H,  $\text{CH}_2$ ), 3.01 (q,  $J = 7$  Hz, 4H,  $\text{CH}_2$ ), 6.89 (s, 2H, pyrrole H), 7.32–7.42 (m, 6H), 7.86 (d,  $J = 7.5$  Hz, 2H, ArH) **4b**;  $^1\text{H}$  NMR (DMSO, 300 MHz):  $\delta = 2.64$  (q,  $J = 7$  Hz, 4H,  $\text{CH}_2$ ), 2.85 (q,  $J = 7$  Hz, 4H,  $\text{CH}_2$ ), 5.34 (s, 2H, C=6;  $\text{CH}_2$ ), 6.15 (s, 2H, pyrrole H), 7.09–7.19 (m, 6H), 7.69 (d,  $J = 7$  Hz, 2H, ArH) **4a**. LC-MS:  $m/z$  364 ( $[\text{M} + \text{H}]^+$ ). Anal. calcd. for  $\text{C}_{26}\text{H}_{22}\text{N}_2$ : C, 86.19; H, 6.08; N, 7.73. Found: C, 86.27; H, 5.92; N, 7.5.




Fig. 2. Solvatochromism of dye **5** ( $C = 1 \times 10^{-5}$  mol/l). 1, hexane ( $\lambda_{\text{max}} = 626$  nm,  $\epsilon = 92,000$ ); 2, toluene ( $\lambda_{\text{max}} = 650$  nm,  $\epsilon = 98,000$ ); 3,  $\text{CH}_2\text{Cl}_2$  ( $\lambda_{\text{max}} = 664$  nm,  $\epsilon = 103,000$ ); 4, DMF ( $\lambda_{\text{max}} = 702$  nm,  $\epsilon = 107,000$ ).

### 3.3. Compound 4

To a stirred solution of salt **2** (1.5 g, 2.8 mmol) in dry chloroform (50 mL), first boron trifluoride etherate (5 mL, 40 mmol) and then *N,N*-diisopropylethylamine (2 mL, 12 mmol) were added. After the reaction mixture was refluxed with stirring for 1 h, it was cooled and washed with water ( $3 \times 50$  mL), and the organic layer was dried over  $\text{Na}_2\text{SO}_4$ . Chloroform was evaporated and the residue was chromatographed on silica gel (using a 3:1 mixture of hexane and ethyl acetate as eluent). Yield 1.03 g, 89%. M.p.  $>250$  °C.  $^1\text{H}$  NMR (DMSO, 300 MHz):  $\delta = 2.64$  (s, 3H,  $\text{CH}_3$ ), 2.73 (q,  $J = 6.3$  Hz, 4H,  $\text{CH}_2$ ), 2.90 (q,  $J = 6.3$  Hz, 4H,  $\text{CH}_2$ ), 7.38–7.43 (m, 8H), 8.53 (d,  $J = 7$  Hz, ArH).  $^{19}\text{F}$  NMR (DMSO, 188 MHz):  $\delta = -136.43$  (q). LC-MS:  $m/z$  392 ( $[\text{M} - 19(\text{F})]^+$ ). Anal. calcd. for  $\text{C}_{26}\text{H}_{21}\text{BF}_2\text{N}_2$ : C, 76.1; H, 5.12; N, 6.83. Found: C, 76.25; H, 5.32; N, 6.9.

### 3.4. General procedure for the synthesis of dyes 5–7

To a mixture of compound **4** (0.1 g, 0.25 mmol) and the corresponding hemicyanine (0.3 mmol) in acetic anhydride (1 mL), triethylamine (0.04 g, 0.4 mmol) was added. The reaction mixture was boiled for 2–3 min and allowed to stand at room temperature overnight. The precipitated product was filtered off.

### 3.5. Dye 5

Yield 0.08 g, 52%. M.p.  $>250$  °C.  $^1\text{H}$  NMR (DMSO, 300 MHz):  $\delta = 1.63$  (s, 6H,  $\text{CH}_3$ ), 2.72 (q,  $J = 7$  Hz, 4H,  $\text{CH}_2$ ), 2.86 (q,  $J = 7$  Hz, 4H,  $\text{CH}_2$ ), 3.48 (s, 3H,  $\text{NCH}_3$ ), 6.24 (d,  $J = 12.6$  Hz, 1H, CH), 7.07–7.36 (m, 12H), 7.48 (d,  $J = 8.1$  Hz, 1H, HetH), 8.17 (t,  $J = 12.9$  Hz, 1H, CH), 8.46 (d,  $J = 7.2$  Hz, 2H, ArH).  $^{19}\text{F}$  NMR (DMSO, 188 MHz):  $\delta = -134.44$  (q). LC-MS:  $m/z$  594 ( $[\text{M} + \text{H}]^+$ ). Anal. calcd. for  $\text{C}_{39}\text{H}_{34}\text{BF}_2\text{N}_3$ : C, 78.92; H, 5.73; N, 7.08. Found: C, 79.07; H, 5.4; N, 7.1.

### 3.6. Dye 6

Yield 0.09 g, 63%. M.p.  $>250$  °C.  $^1\text{H}$  NMR (DMSO, 300 MHz):  $\delta = 2.7$  (q,  $J = 7.2$  Hz, 4H,  $\text{CH}_2$ ), 2.84 (q,  $J = 7$  Hz, 4H,  $\text{CH}_2$ ), 3.79 (s, 3H,  $\text{NCH}_3$ ), 6.67 (d,  $J = 12.6$  Hz, 1H CH), 7.06 (s, 2H, pyrrole H), 7.17–7.35 (m, 8H), 7.52 (t,  $J = 8.1$  Hz, 1H, HetH), 7.66 (d,  $J = 8.4$  Hz, 1H, HetH), 7.86–7.96 (m, 2H), 8.41 (d,  $J = 8.4$  Hz, 2H, ArH).  $^{19}\text{F}$  NMR (DMSO, 188 MHz):  $\delta = -133.94$ – $134.4$  (m). LC-MS:  $m/z$  584 ( $[\text{M} + \text{H}]^+$ ). Anal. calcd. for  $\text{C}_{36}\text{H}_{28}\text{BF}_2\text{N}_3\text{S}$ : C, 74.1; H, 4.8; N, 7.2. Found: C, 74.27; H, 5.05; N, 7.1.

### 3.7. Dye 7

Yield 0.09 g, 64%. M.p.  $>250$  °C.  $^1\text{H}$  NMR (DMSO, 300 MHz):  $\delta = 2.69$  (q,  $J = 7.2$  Hz, 4H,  $\text{CH}_2$ ), 2.82 (q,  $J = 7.2$  Hz, 4H,  $\text{CH}_2$ ), 3.96 (s, 3H,  $\text{NCH}_3$ ), 6.74 (d,  $J = 12.3$  Hz, 1H, CH), 7.08 (s, 2H, pyrrole H), 7.13–7.19 (m, 3H), 7.26–7.33 (m, 4H), 7.46 (t,  $J = 7.2$  Hz, 1H, HetH), 7.73–8.05 (m, 5H), 8.36–8.43 (m, 3H).  $^{19}\text{F}$  NMR (DMSO, 188 MHz):  $\delta = -133.77$ – $134.05$  (m). LC-MS:  $m/z$  580 ( $[\text{M} + \text{H}]^+$ ). Anal. calcd. for  $\text{C}_{38}\text{H}_{31}\text{BF}_2\text{N}_3$ : C, 78.9; H, 5.4; N, 7.26. Found: C, 79.05; H, 5.24; N, 7.11.

### 3.8. Dye 8

To a mixture of compound **4** (0.1 g, 0.25 mmol) and 4-(2-acetanilidovinyl)-1-methylquinolinium tosylate (0.12 g, 0.3 mmol) in pyridine (1 mL), triethylamine (0.04 g, 0.4 mmol) was added. The reaction mixture was heated to boiling and boiled for 2–3 min, followed by cooling to room temperature. After evaporation of pyridine in vacuo, the residue was chromatographed on silica gel (using dichloromethane as eluent). Yield 0.06 g, 43%. M.p.  $>250$  °C.  $^1\text{H}$  NMR (DMSO, 300 MHz):  $\delta = 2.69$  (q,  $J = 6.9$  Hz, 4H,  $\text{CH}_2$ ),

2.82 (q,  $J = 6.9$  Hz, 4H,  $CH_2$ ), 4.07 (s, 3H,  $NCH_3$ ), 6.7 (s, 2H, pyrrole  $H$ ), 7.11–7.38 (m, 8H), 7.61 (d,  $J = 7.2$  Hz, 1H,  $HetH$ ), 7.68–7.73 (m, 1H), 7.93 (d,  $J = 3.3$  Hz, 2H,  $HetH$ ), 8.25 (d,  $J = 6.9$  Hz, 1H,  $HetH$ ), 8.37–8.52 (m, 4H).  $^{19}F$  NMR (DMSO, 188 MHz):  $\delta = -133.61$ – $-133.91$  (m). LC-MS:  $m/z$  580 ( $[M + H]^+$ ). Anal. calcd. for  $C_{38}H_{31}BF_2N_3$ : C, 78.9; H, 5.4; N, 7.26. Found: C, 79.1; H, 5.3; N, 7.15.

#### 4. Conclusions

Merocyanines **5**–**8** derived from a conformationally restricted (bridged) borondipyrromethene nucleus represent long-wavelength polymethine dyes and much resemble, by spectral behaviour, their analogues **5a**–**7a** with flexible conformations of phenyl substituents. Absorption spectra of dyes **5**–**8** also contain two bands, one assigned to the polymethinic transition and the other to the local transition in the borondipyrromethene moiety. It has been shown that bridging mainly affects the location of the short-wavelength band and slightly affects the long-wavelength band. Like conformationally free dyes **5a**–**7a**, merocyanines **5**–**7** exhibit weak fluorescence (only in nonpolar media).

#### References

- [1] Loudet A, Burgess K. BODIPY dyes and their derivatives: syntheses and spectroscopic properties. *Chemical Reviews* 2007;107:4891–932.
- [2] Ulrich G, Zissel R, Harriman A. The chemistry of fluorescent bodipy dyes: versatility unsurpassed. *Angewandte Chemie International Edition* 2008;47:1184–201.
- [3] Wood TE, Thompson A. Advances in the chemistry of dipyrins and their complexes. *Chemical Reviews* 2007;107:1831.
- [4] Haugland RP. The handbook. A guide to fluorescence probes and labeling technologies. 10th ed. Eugene, OR: Invitrogen Corp.; 2005.
- [5] Fabian J, Nakazumi H, Matsuoka M. Near-infrared absorbing dyes. *Chemical Reviews* 1992;92:1197–226.
- [6] Dähne S, Resch-Genger U, Wolfbeis OS, editors. Near-infrared dyes for high technology application. Dordrecht: Kluwer Academic; 1998.
- [7] Rurack K, Kollmannsberger M, Daub J. A highly efficient sensor molecule emitting in the near infrared (NIR): 3,5-distyryl-8-(p-dimethylaminophenyl)-di-fluoroboradiaza-*s*-indacene. *New Journal of Chemistry* 2001;25:289–92.
- [8] Coskun A, Akkaya EU. Difluorobora-*s*-diazaindacene dyes as highly selective dosimetric reagents for fluoride anions. *Tetrahedron Letters* 2004;45:4947–9.
- [9] Rohand T, Qin W, Boens N, Dehaen W. Palladium-catalyzed coupling reactions for the functionalization of BODIPY dyes with fluorescence spanning the visible spectrum. *European Journal of Organic Chemistry*; 2006:4658–63.
- [10] Ekmekci Z, Yilmaz MD, Akkaya EU. A monostyryl-boradiazaindacene (BODIPY) derivative as colorimetric and fluorescent probe for cyanide ions. *Organic Letters* 2008;10:461–4.
- [11] Qin W, Rohand T, Dehaen W, Clifford JN, Driesen K, Beljonne D, et al. Borondipyrromethene analogs with phenyl, styryl, and ethynylphenyl substituents: synthesis, photophysics, electrochemistry, and quantum-chemical calculations. *Journal of Physical Chemistry Part A* 2007;111:8588–97.
- [12] Rurack K, Kollmannsberger M, Daub J. Molecular switching in the near-infrared (NIR) with a functionalized boron-dipyrromethene dye. *Angewandte Chemie International Edition* 2001;40:385–7.
- [13] Yu YH, Descalzo AB, Shen Z, Rohr H, Li Q, Wang YW, et al. Mono- and Di(dimethylamino)styryl-substituted borondipyrromethene and borondiindomethene dyes with intense near-infrared fluorescence. *Chemistry - An Asian Journal* 2006;1-2:176–87.
- [14] Saki N, Dinc T, Akkaya EU. Excimer emission and energy transfer in cofacial boradiazaindacene (BODIPY) dimers built on a xanthene scaffold. *Tetrahedron* 2006;62:2721–5.
- [15] Peng X, Du J, Fan J, Wang J, Wu Y, Zhao J, et al. A selective fluorescent sensor for imaging  $Cd^{2+}$  in living cells. *Journal of the American Chemical Society* 2007;129:1500–1.
- [16] Deniz E, Isbasar GC, Bozdemir OA, Yildirim IT, Siemarczuk A, Akkaya EU. Bidirectional switching of near IR emitting boradiazaindacene fluorophores. *Organic Letters*; 2008:3401–3.
- [17] Baruah M, Qin W, Flors C, Hofkens J, Vallee RAL, Beljonne D, et al. Solvent and pH dependent fluorescent properties of a dimethylaminostyryl borondipyrromethene dye in solution. *Journal of Physical Chemistry Part A* 2006;110:5998–6009.
- [18] James TH. The theory of the photographic process. New York: Macmillan; 1973.
- [19] Tolmachev AI, MYu Kornilov, Karaban EF. Direction of protonation in polymethine dyes. *Theoretical and Experimental Chemistry* 1976;12:644.
- [20] Werner T, Huber C, Heint S, Kollmannsberger M, Daub J, Wolfbeis O. Novel optical pH-sensor based on a boradiaza-indacene derivative. *Fresenius' Journal of Analytical Chemistry* 1997;359:150.
- [21] Kollmannsberger M, Gareis T, Heils S, Brue J, Daub J. Electrogenerated chemiluminescence and proton-dependent switching of fluorescence: functionalized difluoroboradiaza-*s*-indacene. *Angewandte Chemie International Edition* 1997;36:1333–5.
- [22] Kollmannsberger M, Rurack K, Resch-Genger U, Daub J. Ultrafast charge transfer in amino-substituted boron dipyrromethene dyes and its inhibition by cation complexation: a new design concept for highly sensitive fluorescent probes. *Journal of Physical Chemistry Part A* 1998;102:10211–20.
- [23] Rurack K, Kollmannsberger M, Resch-Genger U, Daub J. A selective and sensitive fluoroionophore for Hg(II), Ag(I), and Cu(II) with virtually decoupled fluorophore and receptor units. *Journal of the American Chemical Society* 2000;112:968.
- [24] Shen Z, Rohn H, Rurack K, Uno H, Spieles M, Schulz B, et al. Boron-diindomethene (BDI) dyes and their tetrahydrobicyclo precursors – en route to a new class of highly emissive fluorophores for the red spectral range. *Chemistry - A European Journal* 2004;10:4853.
- [25] Qin W, Baruah M, Van der Auweraer M, De Schryver FC, Boens N. Photophysical properties of borondipyrromethene analogues in solution. *Journal of Physical Chemistry Part A* 2005;109:7371–84.
- [26] Yakubovskiy V, Shandura M, Yu Kovtun. Boradipyrromethenecyanines. *European Journal of Organic Chemistry*; 2009:3237–43.
- [27] Chen J, Reibenspies J, Derecskei-Kovacs A, Burgess K. Through-space  $^{13}C$ – $^{19}F$  coupling can reveal conformations of modified BODIPY dyes. *Chemical Communications*; 1999:2501–2.
- [28] Chen J, Burghart A, Derecskei-Kovacs A, Burgess K. 4,4-Difluoro-4-bora-3a,4a-diaza-*s*-indacene (BODIPY) dyes modified for extended conjugation and restricted bond rotations. *Journal of Organic Chemistry* 2000;65:2900–6.
- [29] Severin T, Bruck B. Umsetzungen mit 1-nitro-2-dimethylamino-ethylen. *Chemische Berichte* 1965;98:3847–53.
- [30] Rettig W, Dekhtyar M. Merocyanines: polyene-polymethine transition in donor-acceptor-substituted stilbenes and polyenes. *Chemical Physics* 2003;293:75–90.
- [31] Mostovnikov V, Rubinov A, Alperovich M, Avdeeva V, Levkoev I, Loiko M. Dependence of the luminescence and generation properties of solutions of polymethine dyes on their structure. *Journal of Applied Spectroscopy* 1974;20:31–5.
- [32] Gerasov A, Shandura M, Kovtun Y. Series of polymethine dyes derived from 2,2-difluoro-1,3,2-(2 *H*)-dioxaborine of 3-acetyl-7-diethylamino-4-hydroxycoumarin. *Dyes and Pigments* 2008;77:598–607.