

Solid-phase synthesis of aryl vinyl ethers based on polystyrene-supported β -phenylselenoethanol

Jia-Li Zhang^a, Shou-Ri Sheng^{b*}, Xue Liu^b and Shu-Ying Lin^b

^aDepartment of Chemistry and Chemical Engineering, East China Jiaotong University, Nanchang 330013, P. R. China


^bCollege of Chemistry and Chemical Engineering, Jiangxi Normal University, Nanchang 330027, P. R. China

A novel facile solid-phase organic synthesis of aryl vinyl ethers by reaction of polystyrene-supported β -phenylselenoethanol with phenols under Mitsunobu conditions and subsequent oxidation-elimination with 30% hydrogen peroxide has been developed. The advantages of this method include straightforward operation, lack of odour, good yield and high purity of the products.

Keywords: solid-phase organic synthesis, polystyrene-supported β -phenylselenoethanol, aryl vinyl ether

Vinyl ethers are valuable intermediates that can be used in a wide array of chemical transformations.¹ More specifically, aryl vinyl ethers with unsubstituted vinyl moiety, have wide synthetic applications and are employed as key intermediates for the generation of new polymeric materials,² as dienophiles for cycloaddition reactions such as [2 + 2],³ [2 + 4]⁴ and 1,3-dipolar cycloadditions,⁵ in cyclopropanations,⁶ in hydroformylations,⁷ and in natural product analogue synthesis.⁸ Aryl vinyl ethers are usually prepared according to the following procedures: the dehydrohalogenation of aryl 2-haloethyl ethers,⁹ the addition of phenols to acetylene¹⁰ and the copper (II)-promoted coupling of arylboronic acids with phenols.¹¹ Recently, the use of vinyl acetate in an iridium-catalysed reaction with phenols,¹² and the transformations with copper (II) acetate mediated coupling of 2, 4, 6-trivinyl-cyclotriphosphazane as a vinylboronic acid equivalent,¹³ and tributyl(vinyl)tin¹⁴ with phenols have also been reported. However, most of these methods involved difficulties such as harsh reactions, laborious manipulation and low overall yields, or in some cases, reactions are unsuitable for sensitive substrates, vigorous toxic compounds are used or some reagents are not readily available. It is well known that phenylseleno group is readily converted to a leaving group giving access to carbon–carbon double bond *via* oxidation followed by β -elimination under extremely mild conditions.¹⁵ Moreover, the polymeric selenium reagents¹⁶ have been now developed for solid-phase organic synthesis (SPOS) with a combined advantage of decrease volatility and simplification of product work-up. In continuation of our interest in solid-phase organoselenium chemistry,¹⁷ describe here a new simple and efficient SPOS approach to aryl vinyl ethers based on a novel polystyrene-supported β -phenylselenoethanol reagent (Scheme 1).

Polymer-supported β -phenylselenoethanol (**3**) was readily prepared by treatment of a THF-swollen suspension of cross-linked (1%) polystyrene-bound selenium bromide (**1**)¹⁶ with LiBH_4 , followed by treatment with 2-chloroethanol. The IR spectrum of resin **3** showed a large hydroxyl absorption at 3400 cm^{-1} , and band at 1060 cm^{-1} (C–O). Resin **3** can be stored at room temperature for a long time without diminution of capacity or the liberation of disagreeable odors. With the resin **3** in hand, the etherification reaction was investigated from resin **3** with phenol (**4a**) under Mitsunobu reaction conditions [triphenylphosphine/diethyl azodicarboxylate] in 4-methylmorpholine to afford polystyrene-supported 2-phenoxyethyl selenide (**5a**) efficiently, which could not be reliably analysed with FT-IR. Hence we carried out next cleavage reaction directly after washing the resin **5a** using solvents. Treatment of resin **5a** with 30% hydrogen peroxide at 0°C and then at room temperature afforded the corresponding phenyl vinyl ether (**6a**) in good yields (90%) and with good purities of crude material (95%). The residual resin, polystyrene-supported phenylseleninic acid, was obtained as a by-product, whose IR data were identical to the previously reported data.¹⁸ The polystyrene-supported phenylseleninic acid could be converted to polymer-supported selenium lithium for recycle by treatment of it with $\text{KI}/\text{Na}_2\text{S}_2\text{O}_3$,^{19,20} followed by bromine.¹⁶ For example, phenyl vinyl ether (**6a**) was obtained in 88% yield under the same reaction condition using the recovered selenium lithium resin (second run), and in 85% yield after second recycle (*i.e.* third run). It was shown that recycling 2–3 times led to a gradual deterioration of the resin. After successfully preparation of **6a**, extension of this method to the synthesis of other analogues in good yields and good purities was investigated (Table 1). As seen from the Table 1, for substrates phenols, with substitution of an



Scheme 1

* Correspondent. E-mail: shengsr@jxnu.edu.cn

Table 1 The yields and purities of aryl vinyl ethers (**6a–6m**)

Entry	R (Phenol)	Product	Yield ^a /%	Purity ^b /%
1	H (4a)	6a	90	95
2	3-CH ₃ (4b)	6b	89	94
3	4-CH ₃ O (4c)	6c	90	97
4	4- <i>t</i> -C ₄ H ₉ (4d)	6d	88	95
5	4-C ₆ H ₅ (4e)	6e	86	96
6	4-Cl (4f)	6f	83	95
7	4-Br (4g)	6g	88	95
8	2-Br (4h)	6h	86	94
9	4-NO ₂ (4i)	6i	90	96
10	4-CN (4j)	6j	88	95
11	4-CO ₂ CH ₃ (4k)	6k	86	94
12	4-NHCOCH ₃ (4l)	6l	88	96
13	1-Naphthol (4m)	6m	84	95

^aOverall yields based on polystyrene-supported selenium bromide **1** (1.18 mmol Br/g).

^bDetermined by HPLC of crude cleavage product ($\lambda = 254$ nm).

electron-withdrawing group or an electron-donating group on the aromatic ring resulted in no obvious effect on the reaction yields.

In summary, we have developed a novel, efficient and convenient method for the SPOS of aryl vinyl ethers employing polymer-supported β -phenylselenoethanol. The advantages of this method include straightforward operation, lack of odour, good yield and high purity of the products.

Experimental

Melting points were uncorrected. ¹H NMR (400 MHz) and ¹³C NMR (100 MHz) spectra were recorded on a Bruker Avance (400 MHz) spectrometer, using CDCl₃ as the solvent and TMS as internal standard. FT-IR spectra were taken on a Perkin-Elmer SP One FT-IR spectrophotometer. HPLC analysis was performed on Agilent 1100 automated system having a PDA detector using a gradient with CH₃CN/H₂O (1 mL min⁻¹) on a RP-18e column (150 × 4.6 mm). Polystyrene (H 1000, 100–200 mesh, cross-linked with 1% divinylbenzene) for preparation of selenium bromide resin¹⁶ was purchased from Nankai University, and the other starting materials were purchased from commercial suppliers and used without further purification. THF was distilled from sodium benzophenone immediately prior to use.

Preparation of polystyrene-supported β -phenylselenoethanol (**3**)

Under a nitrogen atmosphere, to deep red coloured polystyrene-supported selenium bromide **1** (1.0 g, 1.18 mmol Br/g, the loading of functional Br was analysed by elementary analysis) swollen in THF (10 mL) for 30 min was added LiBH₄ (3.0 mmol). After 1 h with shaking at room temperature, 2-chloroethanol (3.0 mmol) in 2 mL of THF was added slowly and the mixture was shaken for 6 h. The resin was collected on a filter and washed successively with H₂O (2 × 20 mL), THF (3 × 5 mL) and CH₂Cl₂ (3 × 5 mL), and then dried under vacuum overnight to afford 950 mg of resin **3**, which was calculated to have a loading of 1.20 mmol g⁻¹, assuming the etherification reaction went to completion. IR (KBr): $\nu = 3400, 3052, 2925, 1595, 1475, 1060, 940, 732, 688$ cm⁻¹.

Preparation of aryl vinyl ethers (**6a–m**); general procedure

The resin **3** (1.0 g, 1.20 mmol) was swollen in THF (10 mL) at room temperature for 30 min and then a solution of triphenylphosphine (1.57 g, 6.0 mmol) and phenols (**4a–m**) (5.0 mmol) in 4-methylmorpholine (5 mL) was added. Neat diethyl azodicarboxylate (790 μ L, 5.0 mmol) was added in small portions over a period of 20 min at room temperature. After the suspension was shaken for 12 h at room temperature, the resin (**5a–m**) was filtered and subsequently washed with THF (3 × 5 mL), DMSO (2 × 10 mL), THF (3 × 5 mL), water (2 × 10 mL), MeOH (2 × 10 mL) and CH₂Cl₂ (3 × 5 mL). To a suspension of the swollen resin **5a–m** in CH₂Cl₂ (10 mL) and 0.5 mL of 30% H₂O₂ (5.8 mmol) was added at 0 °C. The suspension was shaken at 0 °C for 0.5 h and then at room temperature for 1.0 h, the residual resin was collected by filtration and washed with CH₂Cl₂ (2 × 10 mL). The filtrate was treated with saturated NaHCO₃ (20 mL) and washed with water, dried over magnesium sulfate and evaporated to give crude products **6a–m** with 94–97% purity determined by HPLC, which were further purified by column chromatography on

silica gel using chloroform/hexane (10:90) as eluent to give pure products **6a–m** for ¹H NMR and IR analyses.

Phenyl vinyl ether (6a): Colourless oil. (lit.¹⁴ oil); ¹H NMR: $\delta = 7.15$ – 7.02 (m, 5 H), 6.58 (dd, $J = 14.0, 6.0$ Hz, 1 H), 4.70 (dd, $J = 14.0, 1.5$ Hz, 1 H), 4.34 (dd, $J = 6.0, 1.5$ Hz, 1 H); ¹³C NMR: $\delta = 154.0, 144.4, 133.1, 120.5, 115.3, 95.3$; IR (neat): $\nu = 3045, 1640, 1623, 1600, 1495, 1230, 1212, 1165, 1155, 1145, 956, 942$ cm⁻¹.

3-Methylphenyl vinyl ether (6b): Colourless oil (lit.¹⁰ oil); ¹H NMR: $\delta = 6.83$ – 7.20 (m, 4 H), 6.50 (dd, $J = 14.2, 6.5$ Hz, 1 H), 4.32 (dd, $J = 14.2, 1.8$ Hz, 1 H), 4.04 (dd, $J = 6.5, 1.8$ Hz, 1 H), 2.30 (s, 3 H); ¹³C NMR: $\delta = 154.6, 140.4, 132.1, 123.7, 122.6, 119.5, 115.3, 95.6, 21.5$; IR (neat): $\nu = 3050, 1640, 1622, 1600, 1500, 1380, 1230, 1160, 1149, 960, 822$ cm⁻¹.

4-Methoxyphenyl vinyl ether (6c): Colourless oil (lit.²¹ oil); ¹H NMR: $\delta = 6.95$ (d, $J = 8.2$ Hz, 2 H), 6.86 (d, $J = 8.2$ Hz, 2 H), 6.58 (dd, $J = 14.1, 6.3$ Hz, 1 H), 4.65 (dd, $J = 14.1, 2.0$ Hz, 1 H), 4.35 (dd, $J = 6.3, 2.0$ Hz, 1 H), 3.78 (s, 3 H); ¹³C NMR: $\delta = 155.4, 145.8, 134.1, 120.5, 118.0, 115.5, 95.5, 55.2$; IR (neat): $\nu = 3045, 1638, 1620, 1600, 1495, 1379, 1230, 1162, 1145, 958, 825$ cm⁻¹.

4-*t*-Butylphenyl vinyl ether (6d): Colourless oil (lit.⁹ oil); ¹H NMR: $\delta = 6.80$ (d, $J = 8.2$ Hz, 2 H), 7.18 (d, $J = 8.2$ Hz, 2 H), 6.51 (dd, $J = 14.0, 6.2$ Hz, 1 H), 4.28 (dd, $J = 14.0, 1.6$ Hz, 1 H), 4.24 (dd, $J = 6.2, 1.6$ Hz, 1 H), 1.31 (s, 9 H); ¹³C NMR: $\delta = 157.1, 145.5, 133.7, 120.0, 117.3, 95.8, 38.0, 28.5$; IR (neat): $\nu = 3045, 2940, 1640, 1600, 1600, 1500, 1378, 1240, 1180, 1149, 825$ cm⁻¹.

4-Phenylphenyl vinyl ether (6e): White solid, m.p. 72–73 °C. (lit.¹⁴ m.p. 71–72 °C); ¹H NMR: $\delta = 7.66$ – 7.34 (m, 7 H), 7.12–7.06 (m, 2 H), 6.80 (dd, $J = 13.5, 6.0$ Hz, 1 H), 4.72 (dd, $J = 13.5, 1.5$ Hz, 1 H), 4.47 (dd, $J = 6.0, 1.5$ Hz, 1 H); ¹³C NMR: $\delta = 156.6, 148.2, 140.6, 136.3, 129.0, 128.5, 127.2, 126.8, 117.5, 95.5$; IR (KBr): $\nu = 3050, 3021, 1636, 1595, 1509, 1476, 1240, 1136, 826, 755$ cm⁻¹.

4-Chlorophenyl vinyl ether (6f): Colourless oil (lit.²² oil); ¹H NMR: $\delta = 7.43$ (d, $J = 8.4$ Hz, 2 H), 7.01 (d, $J = 8.4$ Hz, 2 H), 6.63 (dd, $J = 13.7, 6.1$ Hz, 1 H), 4.80 (dd, $J = 13.7, 1.8$ Hz, 1 H), 4.51 (dd, $J = 6.1, 1.8$ Hz, 1 H); ¹³C NMR: $\delta = 156.2, 148.1, 132.8, 119.1, 116.1, 95.8$; IR (neat): $\nu = 3048, 1635, 1595, 1475, 1232, 1165, 1142, 1058, 1002, 955, 834$ cm⁻¹.

4-Bromophenyl vinyl ether (6g): Colourless oil (lit.¹⁴ oil); ¹H NMR: $\delta = 7.40$ (d, $J = 8.5$ Hz, 2 H), 6.95 (d, $J = 8.5$ Hz, 2 H), 6.60 (dd, $J = 13.8, 6.1$ Hz, 1 H), 4.78 (dd, $J = 13.8, 1.8$ Hz, 1 H), 4.48 (dd, $J = 6.1, 1.8$ Hz, 1 H); ¹³C NMR: $\delta = 155.9, 147.7, 132.6, 118.5, 115.7, 96.0$; IR (neat): $\nu = 3050, 1636, 1578, 1475, 1230, 1160, 1140, 1060, 1000, 950, 820$ cm⁻¹.

2-Bromophenyl vinyl ether (6h): Colourless oil (lit.¹³ oil); ¹H NMR: $\delta = 7.67$ – 7.65 (m, 1 H), 7.42–7.36 (m, 1 H), 7.21–7.18 (m, 1 H), 7.10–7.06 (m, 1 H), 6.80 (dd, $J = 6.3, 13.5$ Hz, 1 H), 4.68 (dd, $J = 1.8, 13.5$ Hz, 1 H), 4.53 (dd, $J = 1.8, 6.3$ Hz, 1 H); ¹³C NMR: $\delta = 153.5, 148.7, 133.7, 129.7, 125.3, 118.5, 113.6, 96.2$; IR (neat): $\nu = 3043, 1640, 1595, 1472, 1232, 1164, 1142, 1063, 1005, 953, 765$ cm⁻¹.

4-Nitrophenyl vinyl ether (6i): Colourless oil (lit.¹⁴ oil); ¹H NMR: $\delta = 8.25$ (d, $J = 8.9$ Hz, 2 H), 7.10 (d, $J = 8.9$ Hz, 2 H), 6.68 (dd, $J = 13.6, 6.0$ Hz, 1 H), 5.01 (dd, $J = 13.6, 1.9$ Hz, 1 H), 4.70 (dd, $J = 6.0, 1.9$ Hz, 1 H); ¹³C NMR: $\delta = 161.3, 145.5, 142.8, 125.7, 116.3, 99.1$; IR (neat): $\nu = 3060, 1638, 1600, 1580, 1498, 1481, 1330, 1230, 1160, 1120, 1100, 945, 840$ cm⁻¹.

4-Cyanophenyl vinyl ether (6j): Colourless oil (lit.¹⁴ oil); ¹H NMR: $\delta = 7.70$ (d, $J = 8.6$ Hz, 2 H), 7.10 (d, $J = 8.6$ Hz, 2 H), 6.66 (dd, $J = 13.7, 6.1$ Hz, 1 H), 4.98 (dd, $J = 13.7, 2.0$ Hz, 1 H), 4.68 (dd, $J = 6.1, 2.0$ Hz, 1 H); ¹³C NMR: $\delta = 159.8, 145.8, 134.1, 118.6, 117.1, 106.1, 98.6$; IR (neat): $\nu = 3050, 2200, 1635, 1595, 1492, 1300, 1235, 1160, 1125, 950, 824$ cm⁻¹.

4-Methoxycarbonylphenyl vinyl ether (6k): Colourless oil (lit.¹⁴ oil); ¹H NMR: $\delta = 8.00$ (d, $J = 8.3$ Hz, 2 H), 7.15 (d, $J = 8.3$ Hz, 2 H), 6.88 (dd, $J = 13.6, 6.0$ Hz, 1 H), 4.85 (dd, $J = 13.6, 1.6$ Hz, 1 H), 4.55 (dd, $J = 6.0, 1.6$ Hz, 1 H), 3.84 (s, 3 H); ¹³C NMR: $\delta = 166.5, 160.2, 146.6, 131.5, 124.5, 116.1, 97.3, 51.8$; IR (neat): $\nu = 3050, 2985, 2940, 1710, 1635, 1596, 1498, 1425, 1300, 1272, 1235, 1156, 1132, 1100, 840$ cm⁻¹.

4-Acetaminophenyl vinyl ether (6l): White solid, m.p. 102–103 °C. (lit.¹⁴ m.p. 103–103.5 °C); ¹H NMR: $\delta = 7.40$ – 7.50 (m, 2 H), 7.26 (br s, 1 H), 6.95–7.05 (m, 2 H), 6.63 (dd, $J = 13.7, 6.1$ Hz, 1 H), 4.75 (dd, $J = 13.7, 1.7$ Hz, 1 H), 4.55 (dd, $J = 6.1, 1.7$ Hz, 1 H), 2.18 (s, 3 H); ¹³C NMR: $\delta = 169.2, 153.2, 148.5, 133.4, 122.1, 117.3, 94.6, 24.1$; IR (KBr): $\nu = 3258, 3188, 3130, 3055, 1650, 1600, 1495, 1300, 1235, 1210, 1162, 1145, 940, 830$ cm⁻¹.

1-Naphthyl vinyl ether (6m): White solid, m.p. 31–32 °C. (lit.²³ m.p. 32 °C); ¹H NMR: $\delta = 7.00$ – 7.50 (m, 7 H), 6.71 (dd, $J = 14.1, 6.0$ Hz, 1 H), 4.81 (dd, $J = 14.1, 1.6$ Hz, 1 H), 4.45 (dd, $J = 6.0, 1.6$ Hz,

¹H); ¹³C NMR: δ = 152.7, 144.9, 133.9, 132.6, 128.9, 128.7, 128.3, 126.7, 125.9, 123.5, 115.8, 95.7; IR (KBr): ν = 3050, 1630, 1600, 1495, 1255, 1226, 1172, 1152, 1142, 942 cm⁻¹.

We gratefully acknowledge financial support from the Natural Science Foundation of Jiangxi Province of P. R. China (No. 2007GZH839 and No. 2008GZH0029) and the Research Program of Jiangxi Province Department of Education (GJJ08165).

Received 26 December 2008; accepted 6 March 2009

Paper 08/0352 doi: 10.3184/030823409X447682

Published online: 20 May 2009

References

- 1 B.M. Trost and I. Fleming, (eds.), *Comprehensive organic synthesis*, Pergamon: Oxford, Vol. 9, 1991.
- 2 K. Kojima, M. Sawamoto and T. Higashimura, *Macromolecules*, 1989, **22**, 1552.
- 3 H.A.A. El-Nabi, *Tetrahedron*, 1997, **53**, 1813.
- 4 I.E. Markó, G.R. Evans and J.-P. Declercq, *Tetrahedron*, 1994, **50**, 4557.
- 5 S.N. Savinov and D.J. Austin, *J. Chem. Soc., Chem. Commun.*, 1999, 1813.
- 6 P.E. Maligres, M.M. Waters, J. Lee, R.A. Reamer, D. Askin, M.S. Ashwood and M. Cameron, *J. Org. Chem.*, 2002, **67**, 1093.
- 7 A. Nait Ajjou and H. Alper, *J. Am. Chem. Soc.*, 1998, **120**, 1466.
- 8 K.A. Ahrendt, R.G. Bergman and J.A. Ellman, *Org. Lett.*, 2003, **5**, 1301.
- 9 K. Mizuno, Y. Kimura and Y. Otsuji, *Synthesis*, 1979, 688.
- 10 W. Reppe, *Liebigs Ann. Chem.*, 1956, **601**, 8.
- 11 D.A. Evans, J.L. Katz and T.R. West, *Tetrahedron Lett.*, 1998, **39**, 2937.
- 12 Y. Okimoto, S. Sakaguchi and Y. Ishii, *J. Am. Chem. Soc.*, 2002, **124**, 1590.
- 13 N.F. McKinley and D.F. O'Shea, *J. Org. Chem.*, 2004, **69**, 5087.
- 14 M. Blouin and R. Frenette, *J. Org. Chem.*, 2001, **66**, 9043.
- 15 H.J. Reich, *Acc. Chem. Res.*, 1979, **12**, 22.
- 16 K.C. Nicolaou, J. Pastor, S. Barluenga and N. Winssinger, *Chem. Commun.*, 1998, 1947.
- 17 Q.-S. Hu, S.-R. Sheng, S.-Y. Lin, M.-H. Wei, Q. Xin and X.-L. Liu, *J. Chem. Res(S)*, 2007, 74.
- 18 G. Zundel, *Angew. Chem., Int. Ed. Engl.*, 1969, **8**, 499.
- 19 X. Huang and W.-M. Xu, *Tetrahedron Lett.*, 2002, **43**, 5495.
- 20 F. Ferranti and D. De Filippo, *J. Chem. Soc (B)*, 1971, 1925.
- 21 S. Matysiak, H.-P. Fitznar, R. Schnell and W. Pfeleiderer, *Helv. Chim. Acta.*, 1998, **81**, 1545.
- 22 S. Kuwata, Y. Shigemitsu and Y. Odaira, *J. Org. Chem.*, 1973, **38**, 3803.
- 23 H. Christol, H.-J. Cristau and M. Soleiman, *Synthesis*, 1975, 736.