Tetrahedron Letters 50 (2009) 1226-1228

Contents lists available at ScienceDirect

Tetrahedron Letters

journal homepage: www.elsevier.com/locate/tetlet

# Preparation and oxidative coupling of bis[o-(hydrosilyl)phenyl]cuprates and bis[o-(fluorosilyl)phenyl]cuprates

lyl)biphenyls and 2,2'-bis(fluorosilyl)biphenyls.

Atsushi Kawachi\*, Takuya Teranishi, Yohsuke Yamamoto

Department of Chemistry, Graduate School of Science, Hiroshima University, Higashi-Hiroshima 739-8526, Japan

ABSTRACT

### ARTICLE INFO

### Article history Received 23 November 2008 Revised 22 December 2008 Accepted 8 January 2009 Available online 10 January 2009

Keywords: (o-Silvlphenvl)lithium Bis(o-silylphenyl)cuprate Oxidative coupling Biphenyl

1. Introduction

Recently, we reported the preparation of [o-(hydrosilyl)phenyl]lithiums 1 (R = Me (a), Ph (b)) and [o-(fluorosilyl)phenyl]lithiums 2 (R = Me (**a**), Ph (**b**)) by Br-Li exchange of corresponding aryl bromides **3** and **4**, respectively, with *tert*-butyllithium (Scheme 1).<sup>1</sup> Phenyllithiums 1 and 2 reacted with halosilanes and haloboranes to form silicon-functionalized o-disilylbenzenes<sup>1b</sup> and o-(boryl)silylbenzenes.<sup>1c</sup> Then, we turned our attention to exchange of the counter cation (Li<sup>+</sup>) in 1 and 2 with other metals in order to explore the synthetic utility of the o-silvlphenyl anions. Here, we report the conversion of 1 and 2 into bis(o-silylphenyl)cuprates 5 and 6 and the oxidative coupling of 5 and 6, forming silicon-functionalized 2,2'-disilylbiphenyls **7** and  $\mathbf{8}^2$ . The aim of this work is to confirm whether the hydrosilyl and fluorosilyl groups are compatible with intramolecular phenylcuprates and survive oxidative coupling at the ortho position.

First, we investigated the preparation of bis(o-silylphenyl)cuprates 5 and 6 and the oxidative coupling of them according to Lipshutz's procedure (Table 1 and Scheme 1).<sup>3</sup> [o-(Dimethylsilyl) phenyl]lithium (1a) was prepared from phenyl bromide 3a with tert-BuLi in Et<sub>2</sub>O at -78 °C (Scheme 1).<sup>1</sup> Treatment of **1a** with copper cyanide (0.5 equiv) at the same temperature afforded bis[o-(dimethylsilyl)phenyl](cyano)cuprate **5a**.<sup>4</sup> Introduction of gaseous oxygen into the solution of 5a at -78 °C or -110 °C yielded 2,2'bis(dimethylsilyl)biphenyl (7a) in moderate yields (Table 1, entries 1 and 2). Whereas treatment of 5a with 3 equiv of p-dinitroben $zene^5$  as an oxidant at  $-78 \circ C$  afforded the comparable yield of 7a (Table 1, entry 3), the yield of 7a was increased (74% yield) when 5 equiv of *p*-dinitrobenzene was used (Table 1, entry 4). The same treatment performed at -110 °C, however, resulted in a lower yield (43% yield) (Table 1, entry 5). In contrast, bis[o-(fluorodimethylsilyl)phenyl]cuprate 6a afforded only a trace amount of **8a**, and dimerized product **9a** was obtained as a major product  $(8a/9a = 6/94 \text{ in }^{1}\text{H NMR spectra})$  under the same reaction conditions (Scheme 1). The highly electrophilic fluorodimethylsilyl group in **6a** was easily attacked by the highly nucleophilic phenyl-

Bis[o-(hydrosilyl)phenyl]cuprates and bis[o-(fluorosilyl)phenyl]cuprates were prepared by reacting

[o-(hydrosilyl)phenyl]lithiums and [o-(fluorosilyl)phenyl]lithiums, respectively, with copper salts, such

as CuCN and Cu(OPiv)<sub>2</sub>. The phenylcuprates underwent oxidative coupling to afford 2,2'-bis(hydrosi-



(a) for 3a and 4a: tert-BuLi (x 1)/ Et<sub>2</sub>O/-78 °C, 2 h (b) for 3b and 4b: tert-BuLi (x 2)/ THF/ -78 °C, 1 h

Scheme 1. Preparation and reaction of 5 and 6.

39%





© 2009 Elsevier Ltd. All rights reserved.

<sup>\*</sup> Corresponding author. Tel.: +81 82 424 7431; fax: +81 82 424 0723. E-mail address: kawachi@sci.hiroshima-u.ac.jp (A. Kawachi).

<sup>0040-4039/\$ -</sup> see front matter © 2009 Elsevier Ltd. All rights reserved. doi:10.1016/j.tetlet.2009.01.009

 Table 1

 Oxidative coupling of bis[o-(dimethylsilyl)phenyl](cyano)cuprate 5a

Entry	Oxidant	Temp (°C)	Time (h)	Yield of <b>7a</b> (%)
1	02	78	0.5	48
2	02	-110	0.5	54
3	<i>p</i> -Dinitrobenzene (×3)	-78	1	45
4	<i>p</i> -Dinitrobenzene ( $\times$ 5)	-78	1	74
5	<i>p</i> -Dinitrobenzene (×5)	-110	1	43

cuprate moiety in another molecule of **6a** even at the low temperature (Scheme 2).

Lipshutz's procedure was also applicable to phenyl derivatives **1b** and **2b**. The reaction of phenyl bromides **3b** and **4b** with *tert*butyllithium (2 equiv) in THF at -78 °C provided phenyllithiums **1b** and **2b**,<sup>6</sup> which were treated with copper cyanide (0.5 equiv) to give phenylcuprates **5b** and **6b**, respectively (Scheme 1). Subsequent oxidation with *p*-dinitrobenzene (5 equiv) at -78 °C for 1 h produced 2,2'-bis(diphenylsilyl)biphenyls **7b** in 52% yield and **8b** in 39% yield. Corresponding dimerized product **9b** was not obtained perhaps because sterically demanding phenyl groups on the silicon retarded the nucleophilic substitution reaction in **6b** (Scheme 2).

In order to achieve oxidative coupling of the o-(fluorodimethylsilyl)phenyl anion, we examined several combinations of 2a with copper(II) salts  $(MX_2)$ ,<sup>7,8</sup> in which bis[o-(fluorosilyl)phenyl]cuprate **10a–M** (**M** = Cu) was postulated as an intermediate, as shown in Scheme 3. It was expected that organocuprate **10a–M** (**M** = Cu) had lower reactivity toward the fluorosilyl group than cyanocuprate 6a, retarding the formation of 9a. In each reaction, phenyllithium **2a** was treated with MX<sub>2</sub> in Et<sub>2</sub>O at -78 °C for 1 h,<sup>8</sup> and then the reaction mixture was allowed to warm to ambient temperature. The product ratio was determined by <sup>1</sup>H NMR spectroscopy, and the results are summarized in Table 2. Copper(II) dichloride<sup>8a</sup> exclusively provided 9a (8a/9a = 5/95) (Table 2, entry 1), and copper(II) bis(triflate)  $(Cu(OTf)_2)^{8b}$  also preferred the formation of **9a** to that of **8a** (ca. 10% yield) (**8a/9a** = 22/78) (Table 2, entry 2). In contrast, copper(II) bis(pivalate) (Cu(OPiv)<sub>2</sub>)<sup>8c,9</sup> afforded **8a** (ca. 40% yield) in higher yield than 9a (8a/9a = 92/8) (Table 2, entry 3). The other metal(II) salt, cobalt(II) dichloride,<sup>8d</sup> facilitated the dimerization rather than the formation of **8a** via **10a–M** (**M** = Co), giving 9a (8a/9a = 3/97) (Table 2, entry 4). We also treated 2a with tris(acetylacetonate)iron(III) (Fe(acac)<sub>3</sub>)<sup>8e,10</sup> (1.1 equiv) in Et<sub>2</sub>O at -78 °C for 1 h, which preferred 8a (ca. 30% yield) to 9a (8a/9a = 90/10), but a small amount of disiloxane 11 was formed together. The reason for the successful result in the case of Cu (OPiv)<sub>2</sub> remains unclear, but it may be partly due to the high solubility of copper(II) pivalate in Et<sub>2</sub>O, which leads to the complete conversion of 2a into 10a-M (M = Cu) at the low temperature. Finally, as the optimized conditions, we demonstrated that treatment of **2a** with  $Cu(OPiv)_2$  (3 equiv) in Et<sub>2</sub>O at -60 °C for 24 h provided biphenyl 8a in 66% yield together with a small amount of **9a** (**8a/9a** = 93/7) (Scheme 4).

In summary, we prepared bis[o-(hydrosilyl)phenyl]cuprates **5** and bis[o-(fluorosilyl)phenyl]cuprates **6**, which lead to correspond-



Scheme 2. Dimerization of 6



Scheme 3. Preparation and reaction of 10a.

 Table 2

 Oxidative coupling of bis[o-(fluorodimethylsilyl)phenyl]metals 10a-M (M = Cu, Co)

Entry	MX <sub>2</sub>	Temp (°C)	Time (h)	8a:9a
1	$CuCl_2(\times 2)$	-78	1	5:95
2	$Cu(OTf)_2(\times 1)$	-78	1	22:78
3	$Cu(OPiv)_2 (\times 3)$	-78	1	92:8
4	$CoCl_2$ (×0.5)	-78	1	3:97



Scheme 4. Oxidative coupling of 2a using Cu(OPiv)2.

ing silicon-functionalized 2,2'-disilylbiphenyls **7** and **8** by oxidative coupling. The present reactions demonstrate for the first time that the hydrosilyl and fluorosilyl functionalities are compatible with organocuprates at low temperatures and survive oxidative coupling at the *ortho* position.

## 2. Experimental

Preparation of 5a and its oxidative coupling: A solution of tert-BuLi in pentane (1.53 mol/L, 14.0 mL, 21.4 mmol) was added to a solution of **3a** (4.70 g, 21.8 mmol) in Et<sub>2</sub>O (40 mL) at -78 °C over 8 min. The reaction mixture was stirred at that temperature for 2 h. The resulting solution of **1a** was transferred via a Teflon tube to a suspension of CuCN (902 mg, 10.1 mmol) in Et<sub>2</sub>O (20 mL) at -78 °C over 7 min. The reaction mixture was allowed to warm to ambient temperature with vigorous stirring until the solution became homogeneous (for ca. 16 min). Then the resulting solution of **5a** was cooled again to -78 °C. A suspension of *p*-dinitrobenzene (17.0 g, 101 mmol) in Et<sub>2</sub>O (100 mL) was added to the solution of 5a at -78 °C over 15 min. The reaction mixture was stirred at the same temperature for 1 h, and was then allowed to warm to room temperature. The reaction mixture was filtered and the filtrate was concentrated in vacuo. The resulting residue was diluted with hexane (100 mL) and filtered. The filtrate was concentrated in vacuo to give a brown oil (2.57 g). The oil was subjected to column chromatography on silica gel (100 mL) using hexane as eluent ( $R_f = 0.6$ ) to give **7a** (2.19 g, 74% yield) as a colorless oil. Compound **7a**: <sup>1</sup>H NMR  $(CDCl_3, \delta)$ : 0.01 (d, J = 4 Hz, 6H), 0.03 (d, J = 4 Hz, 6H), 4.09 (sept, I = 4 Hz, 2H), 7.21–7.23 (m, 2H), 7.33-7.40 (m, 4H), 7.58–7.60 (m, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>,  $\delta$ ): -3.05, -3.02, 126.66, 128.45, 129.64, 134.65, 136.47, 149.64. <sup>29</sup>Si{<sup>1</sup>H} NMR (CDCl<sub>3</sub>,  $\delta$ ): -18.7 (d of sept,  ${}^{1}J_{\text{Si-H}} = 191 \text{ Hz}, {}^{2}J_{\text{Si-H}} = 6 \text{ Hz}$ ). Anal. Calcd for C<sub>16</sub>H<sub>22</sub>Si<sub>2</sub>: C, 71.04; H, 8.20. Found: C, 70.79; H, 8.24.

Preparation of **5b** and its oxidative coupling: A solution of tert-BuLi in pentane (1.58 mol/L, 1.3 mL, 2.0 mmol) was added to a solution of **3b** (339 mg, 1.0 mmol) in THF (2.0 mL) at -78 °C over 2 min. The reaction mixture was stirred at that temperature for

1 h to give a solution of 1b. A suspension of CuCN (45 mg, 0.50 mmol) in THF (1.0 mL) was transferred via a Teflon tube to the solution of **1b** at  $-78 \,^{\circ}$ C over 3 min. The reaction mixture was allowed to warm to ambient temperature with vigorous stirring until the solution became homogeneous (for ca. 3 min). The resulting solution of **5b** was cooled again to -78 °C. A suspension of p-dinitrobenzene (840 mg, 5.0 mmol) in THF (3.0 mL) was added to the solution of  $\mathbf{5b}$  at  $-78~^\circ\text{C}$  over 2 min. The reaction mixture was stirred at the same temperature for 1 h, and was then allowed to warm to room temperature. The volume of the reaction mixture was reduced to ca. 3 mL by evaporation. The concentrated reaction mixture was diluted with hexane (10 mL) and filtered. The filtrate was concentrated in vacuo to give a brown solid (353 mg). The solid was subjected to column chromatography on silica gel (15 mL) using hexane/AcOEt (20:1) as eluent ( $R_f = 0.23$ ) to give a white solid (247 mg), which was recrystallized from CH<sub>2</sub>Cl<sub>2</sub>/hexane to afford **7b** (134 mg, 52% yield) as colorless crystals. Compound **7b**: mp: 138–139 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, δ): 4.94 (s, 2H), 6.94 (d, I = 7 Hz, 2H), 7.16 (t, I = 7 Hz, 4H), 7.23–7.46 (m, 22H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, *δ*): 126.51, 127.77, 127.80, 128.83, 129.38, 129.40, 130.21, 132.66, 133.84, 134.26, 135.67, 135.85, 136.24, 149.87. <sup>29</sup>Si{<sup>1</sup>H} NMR (CDCl<sub>3</sub>,  $\delta$ ): -22.9 (d, <sup>1</sup>J<sub>Si-H</sub> = 206 Hz, <sup>4</sup>J<sub>Si-H</sub> = 5 Hz). Anal. Calcd for C<sub>36</sub>H<sub>30</sub>Si<sub>2</sub>: C, 83.34; H, 5.83. Found: C, 83.21; H, 5.75.

Preparation of **6b** and its oxidative coupling: A solution of tert-BuLi in pentane (1.58 mol/L, 1.3 mL, 2.0 mmol) was added to a solution of **4b** (360 mg, 1.0 mmol) in THF (2.0 mL) at -78 °C over 3 min. The reaction mixture was stirred at that temperature for 1 h to give a solution of 2b. A suspension of CuCN (45 mg, 0.50 mmol) in THF (1.0 mL) was transferred via a Teflon tube to the solution of 2b at -78 °C over 3 min. The reaction mixture was stirred at -78 °C for 12 min to give a homogeneous solution of **6b**. A suspension of *p*-dinitrobenzene (850 mg, 5.0 mmol) in THF (3.0 mL) was added to the solution of **6b** at -78 °C over 2 min. The reaction mixture was stirred at the same temperature for 1 h, and was then allowed to warm to room temperature. The volume of the reaction mixture was reduced to ca. 3 mL by evaporation. The concentrated reaction mixture was diluted with hexane (10 mL) and filtered. The filtrate was concentrated in vacuo to give a brown solid (365 mg). This solid was washed with hexane to yield pure 8b (108 mg, 39% yield) as colorless crystals. Compound **8b**: **m**p: 139–140 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>,  $\delta$ ): 6.91 (d, I = 8 Hz, 2H), 7.14 (ddd, J = 8 Hz, 8 Hz, and 1 Hz, 2H), 7.23-7.34 (m, 18H), 7.38-7.43 (m, 4H), 7.48 (dd, I = 8 Hz and 1 Hz, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>,  $\delta$ ): 126.40, 127.71, 129.69, 130.08, 130.36 (d,  ${}^{3}I_{C-F} = 14 \text{ Hz}$ ), 132.15  $(d, {}^{2}J_{C-F} = 16 \text{ Hz}), 133.33 (d, {}^{2}J_{C-F} = 17 \text{ Hz}), 133.71 (d, {}^{2}J_{C-F$ 17 Hz), 134.46, 134.48, 135.10, 135.97, 136.00, 149.30. <sup>19</sup>F NMR (CDCl<sub>3</sub>,  $\delta$ ): -165.67 (s). <sup>29</sup>Si{<sup>1</sup>H} NMR (CDCl<sub>3</sub>,  $\delta$ ): -4.7 (d, <sup>1</sup>J<sub>Si</sub>-<sub>F</sub> = 283 Hz). Anal. Calcd for C<sub>36</sub>H<sub>28</sub>F<sub>2</sub>Si<sub>2</sub>: C, 77.94; H, 5.09. Found: C, 77.54; H, 5.30.

Preparation of **10a** and its oxidative coupling: A solution of tert-BuLi in pentane (1.53 mol/L, 0.35 mL, 0.54 mmol) was added to a solution of **4a** (124 mg, 0.53 mmol) in Et<sub>2</sub>O (10 mL) at -78 °C over 3 min. The reaction mixture was stirred at that temperature for 2 h. A suspension of  $Cu(OPiv)_2$  (421 mg, 1.58 mmol) in  $Et_2O$  (1.0 mL) was added to the resulting solution of 1a at -78 °C over 5 min. The reaction mixture was stirred at -60 °C for 24 h, and was then allowed to warm to room temperature. The reaction mixture was diluted with hexane (10 mL) and filtered. The filtrate was concentrated in vacuo. The resulting residue was diluted with hexane (10 mL) and filtered. The filtrate was concentrated in vacuo to give a greenish oil (71 mg). The oil was subjected to bulb-to-bulb distillation (115-125 °C (bath temperature)/0.1 mmHg) to yield 8a with small amount of 9a (total 58 mg, 8a:9a = 93:7, 66% yield of 8a) as a colorless oil. Compound **8a**: <sup>1</sup>H NMR (CDCl<sub>3</sub>,  $\delta$ ): 0.00 (d, <sup>3</sup> $J_{F-H}$  = 8 Hz, 6H), 0.15 (d,  ${}^{3}J_{F-H}$  = 8 Hz, 6H), 7.28–7.31 (m, 2H), 7.45–7.48 (m, 4H), 7.75–7.78 (m, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>,  $\delta$ ): -0.73 (d, <sup>2</sup> $J_{C-F}$  = 16 Hz), 0.90 (d,  ${}^{2}J_{C-F}$  = 16 Hz), 127.05, 129.44, 129.82, 134.11 (d,  ${}^{3}J_{C-F} = 3 \text{ Hz}$ ), 135.67 (d,  ${}^{2}J_{C-F} = 15 \text{ Hz}$ ), 148.54 (d,  ${}^{3}J_{C-F} = 3 \text{ Hz}$ ). <sup>19</sup>F NMR (CDCl<sub>3</sub>,  $\delta$ ): -158.19 (sept,  ${}^{3}J_{F-H} = 8 \text{ Hz}$ ). <sup>29</sup>Si{<sup>1</sup>H} NMR  $(CDCl_3, \delta)$ : 20.5 (d,  ${}^{1}J_{Si-F}$  = 280 Hz). Anal. Calcd for  $C_{16}H_{20}F_2Si_2$ : C, 62.70; H, 6.58. Found: C, 62.41; H, 6.85.

# Acknowledgments

This work was supported by a Grant-in-Aid for Scientific Research on Priority Area 'Advanced Molecular Transformations of Carbon Resources' (No. 19020048) from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

### **References and notes**

- (a) Kawachi, A.; Zaima, M.; Tani, A.; Yamamoto, Y. Chem. Lett. 2007, 36, 362; (b) Kawachi, A.; Tani, A.; Machida, K.; Yamamoto, Y. Organometallics 2007, 26, 4697; (c) Kawachi, A.; Tani, A.; Shimada, J.; Yamamoto, Y. J. Am. Chem. Soc. 2008, 130, 4222.
- Examples of Preparation of 2,2'-disilylbiphenyls: (a) Kira, M.; Sakamoto, K.; Sakurai, H. J. Am. Chem. Soc. **1983**, 105, 7469; (b) Gross, U.; Kaufmann, D. Chem. Ber. **1987**, 120, 991; Binaphthyl analogs were also reported: (c) Fabris, F.; Lucchi, O. D. J. Organomet. Chem. **1996**, 509, 15; (d) Hoshi, T.; Shionoiri, H.; Katano, M.; Suzuki, T.; Hagiwara, H. Tetrahedron: Asymmetry **2002**, 13, 2167. and references cited therein.
- Lipshutz, B. H.; Siegmann, K.; Garcia, E.; Kayer, F. J. Am. Chem. Soc. 1993, 115, 9276.
- 4. These species are recognized as 'cyano-Gilman cuprates'.
- 5. Tamao, K.; Yamaguchi, S.; Shiro, M. J. Am. Chem. Soc. **1994**, 116, 11715.
- Because of the low solubility of phenyl derivatives 1b and 2b in Et<sub>2</sub>O, they were prepared in THF: two equivalents of *tert*-BuLi were required to consume the phenyl bromides completely.
- 7. In oxidative coupling of organometallic compounds using copper(II) salts, it is postulated that the copper(II) salts serve as oxidants, see: Jukes, A. E. Adv. Organomet. Chem **1974**, *12*, 215. and also Ref. 8a.
- The stoichiometries were based on those given in the referenced papers, and we believe that these stoichiometries are not far from the optimized ones: CuCl<sub>2</sub>: (a) Wingerter, S.; Gornitzka, H.; Bertrand, G.; Stalke, D. *Eur. J. Inorg. Chem.* **1999**, 173; Cu(OTf)<sub>2</sub>: (b) Kobayashi, Y.; Taguchi, T.; Tokuno, E. *Tetrahedron Lett.* **1977**, *42*, 3741; Cu(OPiv)<sub>2</sub>: (c) Muci, A. R.; Campos, K. R.; Evans, D. A. *J. Am. Chem. Soc.* **1995**, *117*, 9075; CoCl<sub>2</sub>: (d) Koenig, K. E.; Lein, G. M.; Stuckler, P.; Kaneda, T.; Cram, D. J. *J. Am. Chem. Soc.* **1979**, *101*, 3553; Fe(acaC)<sub>3</sub>: (e) Tsuji, H.; Komatsu, S.; Kanda, T.; Umehara, T.; Saeki, T.; Tamao, K. Chem. Lett. **2006**, *35*, 758.
- Muto, Y.; Hirashima, N.; Tokii, T.; Kato, M.; Suzuki, I. Bull. Chem. Soc. Jpn. 1986, 59, 3762.
- 10. Charles, R. G.; Pawlikowski, M. A. J. Phys. Chem. 1958, 62, 440.