## **Concerted** [2+2] Cycloaddition of Alkenes to a Ruthenium–Phosphorus Double Bond\*\*

Eric J. Derrah, Dimitrios A. Pantazis, Robert McDonald, and Lisa Rosenberg\*

We recently described a terminal phosphido complex of ruthenium containing a  $Ru=PR_2 \pi$  bond, and demonstrated its high activity in 1,2-addition reactions of polar and non-polar substrates.<sup>[1]</sup> Herein we report the [2+2] cycloaddition reactions of this planar phosphido complex with both activated and simple alkenes to yield metallaphosphacyclobutanes, and we provide evidence for a concerted cycloaddition pathway in these P–C bond-forming reactions (Scheme 1 a). Metal-mediated P–C bond-forming reactions



**Scheme 1.** Types of metal-mediated P-C bond formation involving primary or secondary phosphine reagents.

represent a direct and potentially stereoselective route to useful ligands or reagents,<sup>[2]</sup> but the development and understanding of metal-catalyzed phosphination and hydrophosphination by primary or secondary phosphines

[\*] E. J. Derrah, Prof. L. Rosenberg Department of Chemistry, University of Victoria P.O. Box 3065, Victoria, BC V8W3V6 (Canada) Fax: (+1) 250-721-7147 E-mail: lisarose@uvic.ca Homepage: http://web.uvic.ca/~lisarose/ Dr. D. A. Pantazis Institute for Physical and Theoretical Chemistry University of Bonn, 53115 Bonn (Germany) Dr. R. McDonald X-ray Crystallography Laboratory, Department of Chemistry University of Alberta, Edmonton, AB T6G 2G2 (Canada)
[\*\*] We are grateful to the NSERC of Canada for financial support.
Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201000356.

Angew. Chem. Int. Ed. 2010, 49, 3367-3370

lags behind that of the corresponding amination chemistry.<sup>[3]</sup> Among late-metal examples of such catalysis, the critical P-C bond-forming step has been attributed either to nucleophilic attack of free phosphine onto an unsaturated substrate that has been activated through binding with the metal (e.g. Scheme 1 b),<sup>[4]</sup> or to attack of the strongly nucleophilic P of a phosphido intermediate (M-PR2) onto a substrate containing an electrophilic carbon atom (e.g. Scheme 1 c).<sup>[5]</sup> Lanthanideor calcium-catalyzed hydrophosphination of alkynes and alkenes,<sup>[6-8]</sup> which involve the insertion of an unsaturated substrate into the metal-phosphorus bond of a phosphido intermediate (Scheme 1d), include the only examples of metal-mediated hydrophosphination of simple alkenes.<sup>[7b,8]</sup> Our results point to an alternative mechanism for hydrophosphination catalyzed by late-metals that will allow the incorporation of a much wider range of alkene substrates (electron rich and electron deficient) than are currently viable in this transition metal mediated process.<sup>[9]</sup>

The five-coordinate half-sandwich complex  $[Ru(\eta^5-indenyl)(PR_2)(PPh_3)]$  (**1a**,**b**) reacts rapidly with acrylonitrile to give two isomers of the metallacyclic product **2a**,**b**, in which the nitrile group on the ruthenium-bound  $\alpha$ -carbon atom is oriented either *syn* or *anti* with respect to the Ru–indenyl bond (Scheme 2). In solution, these complexes show diag-



Scheme 2.

nostic upfield <sup>31</sup>P{<sup>1</sup>H} NMR signals due to the phosphorus atom in the metallacycle (e.g., *syn*-**2a**:  $\delta = -13.3$  ppm), and upfield <sup>13</sup>C{<sup>1</sup>H} NMR signals for the  $\alpha$ -carbon atom attached to Ru (e.g., *syn*-**2a**:  $\delta = -28.3$  ppm). <sup>13</sup>C{<sup>1</sup>H}DEPT-135 NMR analysis of a mixture of *syn*- and *anti*-**2a** confirmed their identical regiochemistry, showing that the  $\alpha$ -carbon atom in both isomers ( $\delta = -29.1$  ppm in <sup>13</sup>C NMR spectrum for *anti*-**2a**) contains one proton and not two. Unequivocal assignment of the stereochemistry of *syn*-**2b** was obtained from solution NMR studies of a purified sample, which allowed identification of the signal resulting from the lone proton on the  $\alpha$ -carbon atom, and showed its proximity to the PPh<sub>3</sub> aryl protons.<sup>[10]</sup> X-ray crystallography confirmed the analogous *syn* structure of the major isomer of **2a** in the solid state (Figure 1).<sup>[11]</sup> We detected no trace of the other regioisomers



## Communications



*Figure 1.* Molecular structure of syn-**2**a.<sup>[11]</sup> The thermal ellipsoids are drawn at the 20% probability level.

of complexes **2a,b**, in which the nitrile is bound to the carbon atom  $\beta$  to Ru in the metallacycle. Despite the high basicity of P in **1a,b**,<sup>[1a]</sup> we did not observe base-catalyzed acrylonitrile polymerization in these reactions,<sup>[12]</sup> nor did we see telomerization resulting from additional insertions into the Ru–C bond.<sup>[5c]</sup> Finally, we previously showed that **1a,b** in solution exists in equilibrium with a small amount of phospha-alkene/ hydride isomer (e.g., for **1b**, [Ru(P{=C(CH\_3)\_2}iPr)(H)( $\eta^5$ indenyl)(PPh<sub>3</sub>)]).<sup>[1]</sup> Apparently conversion of this structural isomer into **1** occurs more rapidly than its reaction with alkenes, since we neither see it in any of the product mixtures described in this paper, nor do we see any products attributable to its cycloaddition or insertion reactions.

Simple alkenes also undergo [2+2] cycloaddition reactions at the Ru–P  $\pi$  bond in **1a,b** (Scheme 3). Solutions of **1** placed under 1 atm of ethylene rapidly and quantitatively



Scheme 3.

transform into  $[Ru(\eta^5-indenyl)(\kappa^2-CH_2CH_2PR_2)(PPh_3)]$ (**3a,b**) at room temperature, as determined by <sup>31</sup>P{<sup>1</sup>H} NMR analysis. Similar products were obtained from the addition of 1-hexene (**4a,b**) and the strained internal alkene norbornene (**5a,b**); the latter gave exclusive addition at the *exo* face (see the Supporting Information and Figure 2). For 1:1 reactions of these more sterically demanding substrates long reaction times were required, which resulted in competing decomposition to the orthometalated complexes **6a,b**.<sup>[1b]</sup> The use of excess alkene gave much less decomposition and shorter



**Figure 2.** Molecular structure of exo-5a.<sup>[19]</sup> The thermal ellipsoids are drawn at the 20% probability level

reaction times (although still longer than that for ethylene; Scheme 3), a qualitative indication of the rate dependence of cycloaddition upon the concentration of alkene. The sensitivity of this reaction to the steric bulk of the incoming alkene is also highlighted by the complete absence of cycloaddition when one equivalent of cyclooctene was added to 1a,b, even after four weeks, by which time the samples had mostly decomposed into 6a,b.

The participation of simple alkenes in these reactions, and the sensitivity of rates to the size of the alkene, point to a possible concerted [2+2] cycloaddition mechanism, as opposed to the stepwise Michael-addition mechanism responsible for other late-metal phosphido-mediated P-C bondforming reactions.<sup>[13]</sup> Furthermore, we observe no change in the rate of the cycloaddition of 1-hexene to 1a,b when this reaction is carried out in a 1:1 mixture of [D<sub>8</sub>]toluene/THF, relative to that in pure [D<sub>8</sub>]toluene.<sup>[14]</sup> If a stepwise reaction were occurring, we would expect the rate to increase in the more polar solvent mixture because of the stabilization of a zwitterionic intermediate.<sup>[15]</sup> More compelling evidence for the concerted nature of these [2+2] cycloadditions comes from the reactions of *cis*- and *trans*-[D<sub>2</sub>]ethylene with **1a**,**b**. The barrier to rotation around the terminal CHD<sup>-</sup> group in a putative zwitterionic intermediate would be sufficiently low to give significant isotopic scrambling (i.e., four isomers: two cis and two trans) if a stepwise mechanism were important.<sup>[16,17]</sup> The reactions proceed rapidly at room temperature, giving quantitative conversion into yellow solutions of  $[D_2]$ -**3a,b** that each contain just two major isotopomers in a 1:1 ratio, as determined by <sup>31</sup>P{<sup>1</sup>H} NMR analysis (Figure 3).<sup>[18]</sup> These results indicate that stereochemistry at the alkene is conserved, and provide strong evidence for the importance of a concerted cycloaddition mechanism in this P-C bondforming reaction.

Inspection of the molecular structure of *syn*-**2a** (Figure 1) suggests that the diastereoselectivity observed for products **2** and **4** results from crowding at Ru in these metallacycles: the



**Figure 3.** 202.46 MHz <sup>31</sup>P{<sup>1</sup>H} NMR spectra showing a total of four distinct products from the addition of a) *trans*-[D<sub>2</sub>]ethylene and b) *cis*-[D<sub>2</sub>]ethylene to **1a** in [D<sub>8</sub>]toluene. [Ru] = Ru( $\eta^{5}$ -indenyl)(PPh3). Downfield signals represent PPh<sub>3</sub>, and upfield signals represent PCy<sub>2</sub>.

proton on the metallacycle  $\alpha$ -carbon atom points down, toward the bulky PPh<sub>3</sub> ligand in the syn-isomer, whereas for anti-2 or anti-4 the more sterically demanding nitrile or butyl group would point down. However, DFT calculations indicate that the optimized structures of syn- and anti-2a are isoenergetic at zero Kelvin, although thermodynamic corrections lead to a  $\Delta G^{\circ}$  value that is 5.3 kcalmol<sup>-1</sup> lower for syn-2a than that for anti-2a. Interestingly, when PPh<sub>3</sub> is replaced with PH<sub>3</sub> in these calculations, which should eliminate the anticipated steric preference for the syn isomer, the optimized structure of the resulting syn isomer is  $1.5 \text{ kcal mol}^{-1}$  lower in energy than the anti isomer, and thermodynamic corrections give a  $\Delta G^{\circ}$  value for the syn isomer that is 4.6 kcal mol<sup>-1</sup> lower than that for the anti isomer. Therefore, differences in steric crowding in the two diastereomers are small, and their ground-state stabilities are comparable. This suggests that transition-state effects, possibly electronic in origin,<sup>[20]</sup> dictate the observed stereoselectivity (i.e., these are kinetic product distributions).<sup>[21]</sup> We continue to search computationally for the relevant transition states in these concerted cycloaddition reactions to explain both the diastereo- and regioselectivity observed.

There are surprisingly few examples of discrete complexes containing metal-heteroatom bonds that undergo the insertion of simple alkenes, despite the wide range of metalcatalyzed reactions of alkenes for which the proposed mechanisms invoke such chemistry.<sup>[22]</sup> The cycloaddition reactions reported herein include the first examples of intermolecular insertion of simple alkenes into a metal-phosphorus bond, and are distinct from the lanthanide- and calcium-catalyzed hydrophosphination described above in that the additions are occurring at a M–P double bond.<sup>[23]</sup> Currently we are investigating the conditions required to protonolyze the Ru–C bond in these and related metallacyclic ruthenium complexes to achieve a complete hydrophosphination cycle via this metathesis-like chemistry.

## **Experimental Section**

Full synthetic, spectroscopic, and computational details, as well as X-ray crystallographic experimental details for the structures of **2a** and **5a**, are available in the Supporting Information. CCDC 761575 (**2a**) and 761576 (**5a**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam. ac.uk/data\_request/cif.<sup>[11,19]</sup>

Received: January 20, 2010 Published online: March 31, 2010

**Keywords:** alkenes · cycloadditions · metallacycles · P ligands · ruthenium

- a) E. J. Derrah, K. E. Giesbrecht, R. McDonald, L. Rosenberg, Organometallics 2008, 27, 5025; b) E. J. Derrah, D. A. Pantazis, R. McDonald, L. Rosenberg, Organometallics 2007, 26, 1473.
- 2] D. S. Glueck, Chem. Eur. J. 2008, 14, 7108.
- [3] For examples see the following, and references therein; a) J. F. Hartwig, *Nature* 2008, 455, 314; b) T. E. Muller, K. C. Hultzsch, M. Yus, F. Foubelo, M. Tada, *Chem. Rev.* 2008, 108, 3795.
- [4] a) P. Butti, R. Rochat, A. D. Sadow, A. Togni, Angew. Chem.
  2008, 120, 4956; Angew. Chem. Int. Ed. 2008, 47, 4878; b) M. O. Shulyupin, I. G. Trostyanskaya, V. A. Kazankova, I. P. Beletskaya, Russ. J. Org. Chem. 2006, 42, 17; c) A. D. Sadow, A. Togni, J. Am. Chem. Soc. 2005, 127, 17012; d) F. Jérôme, F. Monnier, H. Lawicka, S. Derien, P. H. Dixneuf, Chem. Commun. 2003, 696.
- [5] a) V. S. Chan, M. Chiu, R. G. Bergman, F. D. Toste, J. Am. Chem. Soc. 2009, 131, 6021; b) D. S. Glueck, Synlett 2007, 2627; c) C. Scriban, D. S. Glueck, L. N. Zakharov, W. S. Kassel, A. G. DiPasquale, J. A. Golen, A. L. Rheingold, Organometallics 2006, 25, 5757; d) M. O. Shulyupin, M. A. Kazankova, I. P. Beletskaya, Org. Lett. 2002, 4, 761; e) M. A. Kazankova, M. O. Shulyupin, I. P. Beletskaya, Synlett 2003, 2155. In reference [5d] the authors proposed a mechanism which includes alkene insertion into a M–H bond, but they later revise this interpretation (Ref. [5e]) and describe the mechanism as a Michael-type addition onto activated aryl alkenes, noting that these reactions do not proceed for simple alkenes.
- [6] K. Takaki, K. Komeyama, D. Kobayashi, T. Kawabata, K. Takehira, J. Alloys Compd. 2006, 408, 432.
- [7] a) A. M. Kawaoka, M. R. Douglass, T. J. Marks, Organometallics 2003, 22, 4630; b) A. Motta, I. L. Fragala, T. J. Marks, Organometallics 2005, 24, 4995.
- [8] M. R. Crimmin, A. G. M. Barrett, M. S. Hill, P. B. Hitchcock, P. A. Procopiou, Organometallics 2007, 26, 2953.
- [9] D. K. Wicht, D. S. Glueck in *Catalytic Heterofunctionalization*. From Hydroamination to Hydrozirconation (Ed.: A. Togni, H. Grutzmacher), Wiley-VCH, Weinheim, 2001, p. 143.
- [10] See the Supporting Information for details. Similar data obtained for 2a was complicated by overlap of the peaks of interest with those from the Cy groups.
- [11] Crystallographic data for **2a**:  $C_{42}H_{47}NP_2Ru$ ,  $M_r = 728.82$ ; crystal dimensions (mm)0.57 × 0.28 × 0.28; monocclinic space group  $P2_1/n$  (an alternate setting of  $P2_1/c$  [No. 14]); a = 13.1935(8), b = 16.9872(11), c = 15.6856(10) Å;  $\beta = 91.1278(9)^\circ$ ; V = 3514.8(4) Å<sup>3</sup>; Z = 4;  $\rho_{calcd} = 1.377$  g cm<sup>-3</sup>;  $\mu = 0.568$  mm<sup>-1</sup>;  $\lambda = 0.71073$  Å;  $T = -80^{\circ}$ C;  $2\theta_{max} = 52.80^{\circ}$ ; total data collected = 27646;  $R_1 = 0.0229$  (6571 observed reflections with  $F_o^2 \ge 2\sigma (F_o^2)$ );  $wR_2 = 0.0632$  for 415 variables and all 7203 unique reflections; residual electron density = 0.532 and -0.247 e Å<sup>-3</sup>.
- [12] F. Schaper, S. R. Foley, R. F. Jordan, J. Am. Chem. Soc. 2004, 126, 2114.

Angew. Chem. Int. Ed. 2010, 49, 3367-3370

## Communications

- [13] See for example Ref. [5c] and a) W. Malisch, B. Klupfel, D. Schumacher, M. Nieger, J. Organomet. Chem. 2002, 661, 95.
- [14] We were limited in our choice of polar solvent because of the reactivity of **1a**,**b** with nitriles as well as halide-containing or protic solvents (see Ref. [1a]).
- [15] Solvent effects on rate are particularly pronounced for zwitterionic intermediates in [2+2] additions, which are generally less successful in achieving conformations that minimize the coulombic separation than are those in, for example, [2+4] cycloadditions; a) R. Huisgen, Acc. Chem. Res. 1977, 10, 117; b) R. Huisgen, Pure Appl. Chem. 1980, 52, 2283.
- [16] Examples of the utility of [D<sub>2</sub>]ethylene in diagnosing mechanisms of cycloaddition reactions include: a) P. D. Bartlett, R. A. Minns, K. Hummel, S. P. Elliott, C. M. Sharts, G. M. Cohen, J. Y. Fukunaga, J. Am. Chem. Soc. 1972, 94, 2898; b) K. N. Houk, Y. T. Lin, F. K. Brown, J. Am. Chem. Soc. 1986, 108, 554; c) H. B. Liu, R. J. Hamers, J. Am. Chem. Soc. 1997, 119, 7593; d) P. P. Nicholas, R. T. Carroll, J. Org. Chem. 1968, 33, 2345.
- [17] We also examined the addition of *cis* and *trans*-2-butene to **1 a,b**, but these internal alkenes reacted very slowly, allowing competing orthometalation of **1 a,b** (see Ref. [1b]). The somewhat ambiguous results of these experiments (see the Supporting Information) point to a stereochemical preference, in this sterically congested system, for the formation of *cisoid* metallacycles, which is avoided through the use of *cis* and *trans*-[D<sub>2</sub>]ethylene.
- [18] In the <sup>1</sup>H NMR spectra of *cis-* and *trans-*[D<sub>2</sub>]-**3**a,b, peaks resulting from the metallacycle protons show reduced multiplicities and intensities relative to those in the spectra of [D<sub>0</sub>]-**3**a,b. <sup>2</sup>H NMR spectra confirmed the equal distribution of deuterium throughout all four positions on the metallacycle.
- [19] Crystal data for **5a**: C<sub>46</sub>H<sub>54</sub>P<sub>2</sub>Ru,  $M_r = 769.90$ ; crystal dimensions (mm)0.54 × 0.40 × 0.22; monocclinic space group  $P2_1/n$  (an alternate setting of  $P2_1/c$  [No. 14]); a = 11.6081(7), b = 17.4813(11), c = 19.5737(12) Å;  $\beta = 105.6999(8)^{\circ}$ ; V = 3823.8(4) Å<sup>3</sup>; Z = 4;  $\rho_{calcd} = 1.337$  gcm<sup>-3</sup>;  $\mu = 0.525$  mm<sup>-1</sup>;  $\lambda = 0.71073$  Å; T = -80 °C;  $2\theta_{max} = 54.98^{\circ}$ ; total data collected = 33061;  $R_1 = 0.0272$  (7822 observed reflections with  $F_o^2 \ge 2\sigma(F_o^2)$ );  $wR_2 = 0.0728$  for 442 variables and all 8721 unique reflections; residual electron density = 0.622 and -0.264 e Å<sup>-3</sup>.
- [20] Addition of excess ethylvinylether, an electron-rich terminal alkene of intermediate bulk, to **1a,b** gave quantitative

[2+2] cycloaddition within 30 min at RT (a rate intermediate between acrylonitrile and 1-hexene), but gave the *syn* and *anti* isomers (**7a,b**) in a 1:1 ratio. It is not yet clear whether this selectivity arises from transition state effects in the original cycloaddition (i.e. a purely kinetic phenomenon) or whether the electron-rich nature of the resulting metallacycle hastens equilibration of these isomers (DFT calculations again indicate that these *syn* and *anti* isomers are isoenergetic).

- [21] We have seen no evidence for the reversibility of these [2+2] cycloaddition reactions (i.e., no free alkene appears in solution samples of the purified metallacyclic products).  ${}^{31}P{}^{1}H$  NMR analysis of sealed samples of **2a,b** showed a slight increase in the relative amount of the *anti* isomers (from <5% to ca. 10%) over a year at RT, consistent with extremely slow equilibration, presumably via reversible dissociation of the phosphine end of the metallacycle, which would allow the requisite bond rotation(s) and inversion at Ru to give epimerization at the  $\alpha$ -carbon atom.
- [22] J. W. Tye, J. F. Hartwig, J. Am. Chem. Soc. 2009, 131, 14703.
- [23] Other relevant cycloadditions at M-P double bonds include the following. [2+1] cycloaddition of alkenes and alkynes at metal phosphinidenes (M=PR): a) M. L. G. Borst, N. van der Riet, R. H. Lemmens, F. J. J. de Kanter, M. Schakel, A. W. Ehlers, A. M. Mills, M. Lutz, A. L. Spek, K. Lammertsma, Chem. Eur. J. 2005, 11, 3631; b) M. L. G. Borst, R. E. Bulo, D. J. Gibney, Y. Alem, F. J. J. de Kanter, A. W. Ehlers, M. Schakel, M. Lutz, A. L. Spek, K. Lammertsma, J. Am. Chem. Soc. 2005, 127, 16985; c) A. Marinetti, F. Mathey, Organometallics 1984, 3, 456. This last study gives evidence pointing to a concerted mechanism for this route to metal-bound phosphiranes. [2+2] cycloaddition of alkynes with a zirconium phosphinidene complex to give metallacyclobutene complexes: d) T. L. Breen, D. W. Stephan, Organometallics 1996, 15, 5729. Stoichiometric [2+2] cycloaddition of isocyanates and isothiocyanates with group 6 phosphenium complexes  $(M = P^{(\delta+)}R_2)$ : e) W. Malisch, K. Grun, O. Fey, C. A. El Baky, J. Organomet. Chem. 2000, 595, 285; f) W. Malisch, C. Hahner, K. Grun, J. Reising, R. Goddard, C. Kruger, Inorg. Chim. Acta 1996, 244, 147; g) H. Pfister, W. Malisch, J. Organomet. Chem. 1992, 439, C11. The mechanism of these cycloadditions (Refs. [23e-g]) has not been studied.