

Preparation and Novel One-electron Redox Reactions of a New Stable 4,7-Diisopropylbenzo[1,2-*d*][1,2,3]trisenole and its Radical Cation Salt

Satoshi Ogawa,^a Takamasa Kikuchi,^a Shigeya Niizuma^b and Ryu Sato^{*a}


^a Department of Applied Chemistry and Molecular Science, Faculty of Engineering, Iwate University, Morioka 020, Japan

^b College of Humanities and Social Science, Iwate University, Morioka 020, Japan

Stable benzene-fused trisenide compounds, 4,7-diisopropylbenzo[1,2-*d*][1,2,3]trisenole **2** and its radical cation salt, 4,7-diisopropylbenzo[1,2-*d*][1,2,3]trisenolium hexafluorophosphate **3**, are prepared and undergo novel one-electron redox reactions.

1,2,3-Trisenoles may form an unusual 7π radical cation framework^{1,2} by oxidation. Only three examples of trisenoles have been isolated due to the inherent lack of stability.³ Our interest in the construction of reversible one-electron redox systems by using benzotrichalcogenoles⁴ has prompted us to synthesize a novel stable benzotrichalcogenole and its radical cation. We report here the first synthesis of 4,7-diisopropylbenzo[1,2-*d*][1,2,3]trisenole **2** the 7π radical cation, 4,7-diisopropylbenzo[1,2-*d*][1,2,3]trisenolium hexafluorophosphate **3** and their novel one-electron redox reactions.

The trisenole **2** was synthesized as follows (Scheme 1). The synthetic equivalent of unstable 1,4-diisopropyl-2,3-benzenediselenol, 4,7-diisopropyl-2,2-dimethyl-1,3,2-benzo-



Scheme 1 Reagents: i, Br_2 -Fe, CCl_4 ; ii (a) Mg, Et_2O (b) Se, (c) $\text{H}^+/\text{H}_2\text{O}$, (d) $[\text{O}]$; iii, (a) LiAlH_4 , (b) $\text{H}^+/\text{H}_2\text{O}$, (c) NaH, (d) Me_3SiCl ; iv, (a) tetramethylethylenediamine (tmen), Bu^nLi (4 equiv.), hexane, (b) Se, (c) Me_2SnCl_2 ; v, SeCl_4 , THF; vi, (a) SeOCl_2 , THF, (b) TMSOTf, THF, (c) Sml_2 , THF





Fig. 1 Cyclic voltammogram of **2** (2 mmol dm^{-3}) in MeCN at 20 °C, 0.1 mol dm^{-3} NBu_4Cl supporting electrolyte, glassy-carbon working electrode, ferrocene external standard, potential vs $\text{Ag}/0.01 \text{ mol dm}^{-3}$ AgNO_3 reference; scan rate, 100 mV s^{-1}

diselenastannole **1**,[†] was prepared by a sequence of bromination,⁵ selenation,⁶ trimethylsilyl protection,⁷ ortho lithiation⁸ and dimethyltin protection.⁹ Introduction of a selenium atom at the 2-position was performed by the following two methods. Stannole **1** (321 mg, 0.69 mmol) was treated with selenium tetrachloride (152 mg, 0.69 mmol) in THF at -78°C under an Ar atmosphere. The mixture was stirred at -78°C for 20 min and then at room temp. for 1 h and on work-up, the crude products were purified by column chromatography (silica gel, eluent hexane) to give **2** in 50% yield. Alternatively, sequential treatment of stannole **1** (93 mg, 0.20 mmol) in THF (5 ml) under an Ar atmosphere at -78°C with selenyl chloride (0.015 ml, 0.2 mmol) in THF (5 ml), trimethylsilyl trifluoromethanesulfonate (TMSOTf) (0.036 ml, 0.2 mmol) in THF (5 ml), 0.1 mol dm^{-3} samarium(II) iodide¹⁰ (4.0 ml, 0.4 mmol) in THF followed by the same work-up and purification gave **2** in 93% yield.

Cyclic voltammetry of **2** in MeCN at 20 °C under Ar exhibited a well-defined reversible one-electron oxidation wave at $E_{1/2} = 0.56 \text{ V}$ versus Ag/Ag^+ (Fig. 1). This result implies that **2** provides a stable radical cation even at room temperature.

The novel trisenolium radical cation salt **3** was isolated in the one-electron oxidation of **2** with 1 equiv. of NOPF_6 as a one-electron oxidant in ether-acetonitrile (Scheme 2). The



Scheme 2 Reagents: i, NOPF_6 , MeCN-diethylether; ii, Sml_2 in THF




Fig. 2 ESR spectra of **3** at 16 °C (a) and -70°C (b) in THF

dark blue salt **3** was stable and the structure of **3** in solution was analysed by ^{31}P NMR and ESR spectroscopy. The salt **3** dissolved readily in THF to give a red–purple solution. The ESR spectrum of the solution showed the presence of a singlet peak ($g = 2.0583$ G) attributable to a triselenolium radical cation. Valuable-temperature ESR studies on this species afforded decreasing of the intensities of the signal at low temperature. At -70°C the signal almost disappeared and the colour of the solution changed to dark blue, presumably as a result of partial dimerization² of the radicals (Fig. 2). Interestingly, the salt **3** undergoes one-electron reduction to give **2** quantitatively by treatment with samarium(II) iodide (Scheme 2).¹¹

Thus, the facile interconversion in the redox reaction of **2** and **3** are ascribed to the destabilization of the distorted neutral triselenide framework by lone pair–lone pair repulsion and the stabilization of the oxidized radical cation by 7π -aromaticity.

This work was supported by a Grant-in-Aid for Scientific Research (No. 0554015) from the Ministry of Education, Science and Culture, Japan.

Received, 21st March 1994; Com. 4/01658K

Footnotes

† Compound **1**: Colourless plates (hexane); mp $160.5\text{--}162.0^\circ\text{C}$ [Found: C, 35.85; H, 4.78. $\text{C}_{14}\text{H}_{22}\text{Se}_2\text{Sn}$ requires C, 36.01; H, 4.75%]; ν_{max} (KBr)/ cm^{-1} 2963, 2867, 1872, 1694, 1463, 1385, 1363, 1183, 1158, 902 and 816; δ_{H} (400 MHz, CDCl_3) 1.06 (6H, s, Me), 1.27 (12H, d, J 6.8, Me_2CH), 3.51 (2H, spt., J 6.8, Me_2CH) and 7.04 (2H, s, ArH); δ_{C} (100 MHz, CDCl_3) 1.0, 23.4, 37.6, 122.9, 138.5 and 148.3; δ_{Se} (76 MHz, CDCl_3 , relative to neat SeMe_2) 83.2 [$J(^{77}\text{Se}^{117}\text{Sn})$ 1061, $J(^{77}\text{Se}^{119}\text{Sn})$ 1110]; δ_{Sn} (149 MHz, CDCl_3 , relative to neat SnMe_4) 62.9 [$J(^{119}\text{Sn}^{77}\text{Se})$ 1110]; m/z 440 (M^+ , ^{80}Se , ^{120}Sn).

‡ Compound **2**: Reddish brown oil [Found: (M^+ , 100%, $^{80}\text{Se}^{80}\text{Se}^{78}\text{Se}$) 397.8743 and (84.5%, $^{80}\text{Se}^{80}\text{Se}^{80}\text{Se}$) 399.8682. $\text{C}_{12}\text{H}_{16}\text{Se}_3$ requires (M^+ , 100%, $^{80}\text{Se}^{80}\text{Se}^{78}\text{Se}$) 397.8761 and (86.2%, $^{80}\text{Se}^{80}\text{Se}^{80}\text{Se}$) 399.8753]; ν_{max} (neat)/ cm^{-1} 2958, 2865, 1885, 1754, 1462, 1385, 1367, 1323, 1160, 1104, 906 and 820; δ_{H} (400 MHz, CDCl_3) 1.24 (12H, d, J 6.8, Me_2CH), 3.02 (2H, d, J 6.8, Me_2CH) and 7.05 (2H, s, ArH); δ_{C} (100 MHz, CDCl_3) 23.4, 38.7, 125.0, 141.5 and 146.3; δ_{Se} (76 MHz, CDCl_3 , relative to neat SeMe_2) 440.9 [$J(^{77}\text{Se}^{77}\text{Se})$ 258] and 547.0 [$J(^{77}\text{Se}^{77}\text{Se})$ 258].

§ Compound **3**: Blue powder (ether); mp $166.0\text{--}172.0^\circ\text{C}$ (decomp.); [Found: C, 26.78; H, 3.29. $\text{C}_{12}\text{H}_{16}\text{F}_6\text{PSe}_3$ requires C, 26.59; H,

2.97%]; δ_{P} (162 MHz, $[\text{H}_8]\text{THF}$, relative to H_3PO_4) -144.0 [spt., $J(^{31}\text{P}^{19}\text{F})$ 706]

References

- 1 P. D. Boyle, S. Parsons, J. Passmore and D. J. Wood, *J. Chem. Soc., Chem. Commun.*, 1993, 199; T. S. Cameron, R. C. Haddon, S. M. Mattar, S. Parsons, J. Passmore and A. P. Ramirez, *Inorg. Chem.*, 1992, **31**, 2274; T. S. Cameron, R. C. Haddon, S. M. Mattar, S. Parsons, J. Passmore and A. P. Ramirez, *J. Chem. Soc., Chem. Commun.*, 1991, 358; E. G. Awere, J. Passmore, P. S. White and T. Klapötke, *J. Chem. Soc., Chem. Commun.*, 1989, 1415; E. G. Awere, J. Passmore, K. F. Preston and L. H. Sutcliffe, *Can. J. Chem.* 1988, **66**, 1776; E. G. Awere, N. Burford, C. Mailer, J. Passmore, M. J. Schriver, P. S. White, A. J. Banister, H. Oberhammer and L. H. Sutcliffe, *J. Chem. Soc., Chem. Commun.*, 1987, 66.
- 2 Recently, benzotriselenolium trifluoromethanesulfonate has been reported; G. Wolmershäuser and G. Heckmann, *Angew. Chem., Int. Ed. Engl.*, 1992, **31**, 779.
- 3 M. J. Earle, K. R. Griffiths and A. G. Massey, *Polyhedron*, 1992, **11**, 395; R. E. Humphries and A. G. Massey, *Phosphorus, Sulfur*, 1988, **36**, 135; N. Tokitoh, H. Ishizuka and W. Ando, *Chem. Lett.*, 1988, 657.
- 4 S. Ogawa, N. Yomoji, S. Chida and R. Sato, *Chem. Lett.*, 1994, 507; N. Yomoji, S. Takahashi, S. Chida, S. Ogawa and R. Sato, *J. Chem. Soc., Perkin Trans. 1*, 1993, 1995; and references cited therein.
- 5 B. L. Patel, C. B. Ziegler, N. A. Cortese, J. E. Plevyak, T. C. Zebrovitz, M. Terpkio and R. F. Heck, *J. Org. Chem.*, 1977, **42**, 3903.
- 6 D. G. Foster, *Org. Synth.*, Coll. Vol. 3, 771.
- 7 M. Suzuki, T. Kawagishi and R. Noyori, *Tetrahedron Lett.*, 1981, **22**, 1809; P. Dowd and P. Kennedy, *Synth. Commun.*, 1981, 935; N. Miyoshi, H. Ishii, K. Kondo, S. Murai and N. Sonoda, *Synthesis*, 1979, 300.
- 8 G. D. Figuly, C. K. Loop and J. C. Martin, *J. Am. Chem. Soc.*, 1989, **111**, 654; E. Block, V. Eswarakrishnan, M. Gernon, G. Ofori-Okai, C. Saha, K. Tang and J. Zubieta, *J. Am. Chem. Soc.*, 1989, **111**, 658; K. Smith, C. M. Lindsay and G. J. Pritchard, *J. Am. Chem. Soc.*, 1989, **111**, 665.
- 9 K. Grätz, F. Huber, A. Silvestri, G. Alonzo and R. Barbieri, *J. Organomet. Chem.*, 1985, **290**, 41; R. C. Poller and J. A. Spillman, *J. Chem. Soc. A*, 1966, 958.
- 10 P. Girard, J. L. Namy and H. B. Kagan, *J. Am. Chem. Soc.*, 1980, **102**, 2693; J. A. Soderquist, *Aldrichimica Acta*, 1991, **24**, 15.
- 11 Interconvertible two-electron redox reactions of selenium containing heterocycles have been reported; H. Fujihara, H. Mima, T. Erata and N. Furukawa, *J. Am. Chem. Soc.*, 1992, **114**, 3117.