

Novel Valence Expansion Reactions Using KC₈: A New Route to Hexavalent Organotellurium Compounds from Divalent Tellurium

Masataka Miyasato, Mao Minoura,[†] Yohsuke Yamamoto, and Kin-ya Akiba^{*††}
 Department of Chemistry, Graduate School of Science, Hiroshima University,
 1-3-1 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-8526

[†]Department of Chemistry, School of Science, Kitasato University, 1-15-1 Kitasato, Sagami-hara 228-8555


^{††}Advanced Research Center for Science and Engineering, Waseda University, 3-4-1 Ohkubo, Tokyo 169-8555

(Received December 18, 2001; CL-011272)

Hexavalent tellurium compounds, Ar₅(CH₃)Te, Ar₄(CH₃)₂Te, and Ar₂(CH₃)₄Te (Ar = 4-CF₃C₆H₄, Ph, 4-CH₃C₆H₄), were synthesized by the reaction of Ar_{2-n}(CH₃)_nTe (n = 0–1) or Ar_{3-m}(CH₃)_mTe⁺X[−] (m = 0–2) with KC₈ followed by the treatment with CH₃I.

Valence of main group element compounds are generally exalted higher by oxidizing reagents such as halogens and valence shell expansion takes place in certain cases. For example, Ph₄F₂Te was prepared by treatment of Ph₄Te with XeF₂ as an oxidant.¹ However, reactions of R₂Te or R₄Te with alkyl halides did not yield hypervalent tellurium compounds but the corresponding oniums R₃Te⁺.² Here we report one-pot novel valence expansion reactions from Ar_{2-n}(CH₃)_nTe(II) (n = 0–1) or Ar_{3-m}(CH₃)_mTe⁺X[−](IV) (m = 0–2) to Ar₅(CH₃)Te(VI), Ar₄(CH₃)₂Te(VI), and Ar₂(CH₃)₄Te(VI) (Ar = 4-CF₃C₆H₄, Ph, 4-CH₃C₆H₄) by use of KC₈ followed by treatment with CH₃I.

The reaction of (4-CF₃C₆H₄)₂Te(II) with potassium graphite (KC₈)³ (ca. 15 equiv) was carried out at −78 °C in THF for 5 min and the reaction mixture was treated with CH₃I (ca. 30 equiv) to afford several hexavalent tellurium compounds, that is, (4-CF₃C₆H₄)₅(CH₃)Te (**1**) (4%), (4-CF₃C₆H₄)₄(CH₃)₂Te (*trans*-**2**) (9%), and (4-CF₃C₆H₄)₂(CH₃)₄Te (*trans*-**3**) (10%) (Scheme 1).^{4,6} These hexavalent tellurium compounds are stable to atmospheric moisture and could be separated by recycling HPLC. They were characterized by spectroscopies and elemental analyses. X-ray analysis of *trans*-**2** and *trans*-**3** confirmed that these were octahedral and the two methyl groups in *trans*-**2** and the two 4-CF₃C₆H₄ groups in *trans*-**3** were located *trans* to each other. The ORTEP drawing of *trans*-**3** is shown in Figure 1.⁷



a: 1) KC₈, THF, −78 °C, 2) CH₃I, −78 °C–rt.

Scheme 1. Synthesis of hexaorganotellurium compounds.

The reaction of (4-CF₃C₆H₄)₃Te⁺Cl[−] with KC₈ followed by treatment with CH₃I afforded a similar mixture of the hexavalent tellurium compounds (**1**: 21%, *trans*-**2**: 7%, and *trans*-**3**: 7%) (Table 1). When (4-CF₃C₆H₄)(CH₃)Te, (4-CF₃C₆H₄)₂-(CH₃)Te⁺OTf[−], and (4-CF₃C₆H₄)(CH₃)₂Te⁺I[−] were used instead of homoleptic materials as mentioned above, i.e., (4-




Figure 1. X-ray structure of *trans*-**3** (30% thermal ellipsoid). The structure is almost perfectly octahedral; bond angles for each *cis* group is in the range of 90 ± 0.7° and for each *trans* group is in the range of 180 ± 1.4°. Selected bond lengths (Å): Te–C(1) 2.25(1), Te–C(8) 2.19(1), Te–C(9) 2.18(1), Te–C(10) 2.18(1).

Table 1. Yields of the reaction of organotellurium compounds with KC₈ (CH₃K, PhCH₂K) after treatment with CH₃I

Starting Material	Reagent	Yields (% based on Te)		
		1	<i>trans</i> - 2	<i>trans</i> - 3
(4-CF ₃ C ₆ H ₄) ₂ Te	A	4	9	10
	B	no	11	no
	C	no	13	no
(4-CF ₃ C ₆ H ₄)(CH ₃)Te	A	no	no	4
(4-CF ₃ C ₆ H ₄) ₃ Te ⁺ Cl [−]	A	21	7	7
	B	16	6	no
(4-CF ₃ C ₆ H ₄) ₂ (CH ₃)Te ⁺ CF ₃ SO ₃ [−]	A	no	31	4
(4-CF ₃ C ₆ H ₄)(CH ₃) ₂ Te ⁺ I [−]	A	no	no	16
		4 <i>trans</i> - 5		
Ph ₃ Te ⁺ Br [−] ^a	A	19	12	
		6 <i>trans</i> - 7		
(4-CH ₃ C ₆ H ₄) ₃ Te ⁺ Cl [−] ^b	A	36	7	

A; KC₈ (15 equiv), B; CH₃K (2 equiv), C; PhCH₂K (1 equiv). CH₃I (30 equiv) was used. no; not obtained.

^aC₆H₅I (4 equiv) was added. ^b4-CH₃C₆H₄I (4 equiv) was added.

CF₃C₆H₄)₂Te and (4-CF₃C₆H₄)₃Te⁺Cl[−], monomethylated **1** was not obtained at all but di- and tetramethylated compounds (*trans*-**2** and *trans*-**3**) were obtained with increased yields. For example, *trans*-**2** (31%) and *trans*-**3** (4%) were obtained from (4-CF₃C₆H₄)₂(CH₃)Te⁺OTf[−] and only *trans*-**3** was obtained from

