

Article



Subscriber access provided by University of Sunderland

Use of Cyclopropane as C1 Synthetic Unit by Directed Retro-Cyclopropanation with Ethylene Release

Sobi Asako, Takaaki Kobashi, and Kazuhiko Takai

J. Am. Chem. Soc., Just Accepted Manuscript • DOI: 10.1021/jacs.8b09297 • Publication Date (Web): 22 Oct 2018

Downloaded from http://pubs.acs.org on October 22, 2018

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Use of Cyclopropane as C1 Synthetic Unit by Directed Retro-Cyclopropanation with Ethylene Release

Sobi Asako*, Takaaki Kobashi, and Kazuhiko Takai*

Division of Applied Chemistry, Graduate School of Natural Science and Technology, Okayama University, 3-1-1 Tsushimanaka, Kita-ku, Okayama 700-8530

ABSTRACT: Cyclopropanation of alkenes is a well-established textbook reaction for the synthesis of cyclopropanes, where "high-energy" carbene species is exploited for driving the reaction forward. However, little attention has been paid to molecular transformations involving the reverse reaction, retro-cyclopropanation (RC). This is because of difficulties associated with both cleaving the two geminal C–C single bonds and exploiting the generated carbenes for further transformations in an efficient and selective manner. Here we report that a molybdenum-based catalytic system overcomes the above challenges and effects the RC of cyclopropanes bearing a pyridyl group with the release of ethylene (alkene) and the subsequent intramolecular cyclization leading to pyrido[2,1-a]isoindoles. The reaction allows for the uncommon use of cyclopropanes as C1 synthetic units in contrast to most conventional reactions in which cyclopropanes are used as C3 synthetic units. We anticipate that this new strategy will pave the way for C1 cyclopropane chemistry.

INTRODUCTION

Cyclopropanes are three-membered carbocycles that display unique properties and reactivities because of their unusual bonding and ring strain. Ubiquitous in naturally occurring compounds and rationally designed drugs, cyclopropane-containing molecules themselves are important, ^{1,2,3} ^{4,5,6,7,8} but they also serve as versatile synthetic in modern organic intermediates synthesis.8,9,10,11,12,13,14,15 Certain transition metals are capable of cleaving the C–C bond via oxidative addition to form metallacyclobutane intermediates with strain release (ca. 27 kcal/mol) as a driving force (Fig. 1a, top).^{16,17,18,19} Synthetic chemists have successfully utilized the key intermediates, mostly as C3 fragments for various transformations such as addition. cycloaddition, and cycloisomerization reactions, wherein activated cyclopropanes such as donor-acceptor cyclopropanes.²⁰ vinylcyclopropanes,²¹ and alkylidenecyclopropanes²² are often employed to facilitate the C–C bond cleavage step.

If the metallacyclobutane intermediate further undergoes the second cleavage of a C–C bond, a metal carbene species is formed with the release of ethylene, a common step involved in the olefin metathesis reaction (Fig. 1a, bottom). The overall sequence is retrocyclopropanation (RC), i.e., fragmentation of a cyclopropane into an ethylene and a carbene species. This is a highly uphill and difficult process unless appropriate stabilization of the generated metal carbene is exerted. Indeed, the generation of metal carbene largely relies on the cleavage of the weak bonds in "high-energy" compounds such as diazo compounds, frequently used but potentially explosive carbene precursors, and the metal carbene generation by breaking the two C-C bonds of stable cyclopropanes remains challenging.²³ Because of the lack of efficient and selective catalytic systems that can achieve both RC and subsequent transformation of metal carbenes, such reactions that use cyclopropanes as C1 synthetic units have scarcely been reported. In this context, the biosynthesis of ethylene, a natural plant hormone, by oxidative fragmentation of 1-aminocyclopropane-1carboxylic acid with concomitant release of HCN (C1 unit) and CO₂ may be relevant but lacks the intermediacy of a carbene species.^{24,25}

Cationic gold(I) can promote the retro-Buchner reaction, a special case of RC, where a substituted methylene moiety is removed from 1,3,5-cycloheptatriene, which is in equilibrium with norcaradiene, with the release of benzene (Fig. 1b).²⁶

R

The intramolecular RC of bicyclo[1.1.0]butane is proposed to be involved under transition-metal catalysis, such as Ni²⁷ and Rh²⁸ (Fig. 1c). These reactions take advantage of aromatization and the release of extremely high ring strain (63-67 kcal/mol) 18,19 as driving force. However, the substrates are then limited to highly strained cyclopropanes. The intermolecular RC was observed in the Rh(I)-catalyzed [2+2+1] cycloaddition reaction of alkylidenecyclopropanes²⁹ and the Rh(I)catalyzed [5+2-2] cycloisomerization-type reaction of allenylcyclopropanes.³⁰ However, the former reaction still relies on higher ring the strain of alkylidenecyclopropanes (38–41 kcal/mol)18,19 and requires slow addition of divnes to suppress side reactions following non-RC paths in some cases, and the latter reaction suffers from low selectivity between the cycloisomerizations with and without ethylene release (Fig. 1d and 1e). The PhWCl₃/RAlCl₂ binary system was found to catalyze RC of simple cyclopropanes³¹ and cyclopropane-olefin metathesis,32 albeit again with low efficiency (Fig. 1f).33

1

2

3

4

5

6

7

8

9 10

11

12

13

14

15

16

17

18

19

20

21

22

23

24 25

26 27

35

36 37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52 53

54

55

56

57 58 59

60

a Cyclopropane in synthetic chemistry: release of ring strain (27 kcal/mol) C3 unit via one-fold C-C bond cleavage (conventional)

C1 unit via two-fold gem C-C bond cleavage (uncommon)

$$\begin{array}{cccc} R & & Metal & \stackrel{H}{\longrightarrow} & M_{-} \\ R' & & & & \\ R' & & & & \\ \end{array} \xrightarrow{H} & & & \\ R' & & & \\ \end{array} \xrightarrow{H} & & & \\ \end{array} \xrightarrow{H} & \\ \xrightarrow{H} & \\ \end{array} \xrightarrow{H} & \\ \xrightarrow{H} & \\ \xrightarrow{H} & \\ \end{array} \xrightarrow{H} & \\ \xrightarrow{H} & \xrightarrow{H} & \\ \xrightarrow{H} & \\ \xrightarrow{H} & \xrightarrow{H} & \\ \xrightarrow{H} & \xrightarrow{H} & \\ \xrightarrow{H} & \xrightarrow{H} & \xrightarrow{H} & \\ \xrightarrow{H} & \xrightarrow{H}$$

b Retro-Buchner reaction of cycloheptatriene: aromatization

c Intramolecular RC of bicyclo[1,1,0]butane: release of very high ring strain (63-67 kcal/mol)

$$\bigcirc \quad \stackrel{\mathsf{M}}{\longrightarrow} \quad \bigvee_{=\mathsf{M}} \quad$$

d Intermolecular RC of alkylidenecyclopropane: release of high ring strain (38-41 kcal/mol) followed by cycloaddition: slow addition of divne



e Intermolecular RC of allenylcyclopropane: followed by intramolecular cyclization: low selectivity and yield



f Intermolecular RC of simple cyclopropane: followed by cyclopropanation (cyclopropane-olefin metathesis): low yield









Figure 1. Organic transformations involving retrocyclopropanation with metal catalysts.

Thus, to date, few efficient catalytic systems are available for molecular transformations featuring RC with synthetic utility. Here we describe a molybdenumcatalyzed directed RC/intramolecular cyclization sequence of cyclopropanes possessing a pyridine moiety that plays multiple roles (Fig. 1g). With the prevalence of the chelation assistance in the activation of chemically inert bonds, such as C-H and C-C bonds, together with our own experience in this field,^{34,35} we envisioned that the use of a pyridyl group at an appropriate position for RC would be beneficial for the following reasons: it would 1) facilitate the first cleavage of a proximal C-C bond in a regioselective manner, 2) prohibit the undesired decomposition of the metallacyclobutane intermediate I through a hydride elimination process,³⁶ 3) adequately stabilize the carbene species II formed after the elimination of ethylene from I, and 4) efficiently trap the carbene by intramolecular cyclization to incorporate itself into the

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46 47

48

49

50

51

52

53

54

55

60

value-added product. The directing group approaches for the regioselective transformations of cyclopropanes have been reported.³⁷ However, the cyclopropanes are still used as C3 synthetic units in these cases, because the C– C bond cleavage takes place only once.

RESULTS AND DISCUSSION

To test this hypothesis, we commenced our investigation using 2-[2-(1phenylcyclopropyl)phenyl]pyridine (1a) as a model substrate (Fig. 2). Close similarity between the proposed mechanism in our strategy (Fig. 1g) and that of the olefin metathesis, significant advances of which have been made with molybdenum catalysts^{38,39} besides our recent interest in molybdenum catalysis^{40,41} prompted us to use this metal as a catalyst of choice; hence we first various combinations of molybdenum explored complexes and ligands. After extensive experimentation. we found that a catalytic amount of Mo(benzene)₂ combined with 9,10-phenanthrenequinone (L1)effectively catalyzed the RC/intramolecular cyclization reactions to afford 6-phenyl[2,1-*a*]pyridoisoindole (2a) in quantitative yield. The reaction reached completion with 20 mol % catalyst after 24 h at 160 °C in mesitylene, whereas at lower temperatures the reaction rate was significantly decreased (Table S1), partly because the thermodynamic driving force of releasing ethylene decreases as the temperature decreases ($\Delta G = -$ 3.7, -2.3, and +1.1 kcal/mol, at 160, 120, and 25 °C, respectively). The reaction still proceeded with 10 mol % catalyst loading, but with slower reaction rates (Table S1).

Screening of various ligands revealed that *o*-quinones were uniquely effective in the reaction; L1 and L2 performed best (Fig. 2a). While catechol showed similar catalytic activity (L3), common bidentate nitrogen- or phosphine-based ligands completely shut off the reaction (L4–L7). Mo(benzene)₂ alone could effect the reaction to some extent, but the reaction in the absence of molybdenum did not proceed at all. Other Mo(0) complexes such as Mo(CO)₆ and Mo(CO)₃(MeCN)₃ also worked, but less efficiently (Table S2).

The scope of the directed RC reaction is illustrated in Fig. 2b. Cyclopropanes possessing an electron-rich and electron-deficient aryl group at the 1-position of the cyclopropane both reacted smoothly (2b-2g). Cyclopropanes with a bulky 2-tolyl or alkyl substituents were less reactive but they still underwent the RC reactions at higher temperature and after longer reaction time (2h-2k). Due to the moderate instability of 6alkylpyrido[2,1-*a*]isoindoles (2i-2k) under atmospheric conditions, these products were oxidized and then isolated as the corresponding alkyl 2-(2-pyridyl)phenyl ketones. 2-(2-Cyclopropylphenyl)pyridine remained unchanged, indicating the necessary presence of a substituent at the 1-position of the cyclopropane. Both an electron donating and withdrawing substituent on the 2-phenylpyridine moiety slowed down the reaction (**2m** and **2n**). In addition to a pyridyl group, a pyrimidyl group also worked as a multifunctional unit and the corresponding pyrimido[2,1-*a*]isoindole (**2o**) was obtained in good yield, whereas an imino group failed to effect the reaction. All the 6-arylpyrido[2,1-*a*]isoindole derivatives synthesized by this method are previously unknown compounds, and exhibit strong fluorescence.⁴²



Figure 2. Molybdenum-catalyzed synthesis of pyrido[2,1-*a*]isoindoles via retro-cyclopropanation. a, Gibbs free energies and enthalpies of the reaction calculated at the M06/6-31G(d,p)_{mesitylene(SMD)} level of theory and the effects of ligands. ¹H NMR yields are shown. b, Scope of the molybdenum-catalyzed RC. The reactions were performed on a 0.2 mmol scale with Mo(benezene)₂/L1 (20 mol%) in mesitylene (0.25 M). Isolated yields are shown. ^{*a*}160 °C, 24 h. ^{*b*}160 °C, 18 h.

^c180 °C, 24 h. ^d180 °C, 72 h. ^eIsolated as ketone after oxidation. Ph, phenyl; [']Bu, *tert*-butyl; Me, methyl.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21 22

23

24

25

26 27

28 29

30

31

32

33

34

35 36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60

The importance of a pyridyl directing group for the molybdenum-catalyzed RC was illustrated by the reaction of 2-(1-phenylcyclopropyl)-1,1'-biphenyl. It did not afford any products with ethylene loss or ringopened products both in the presence and absence of added pyridine (Fig. 3a). Monitoring the reaction in pxylene- d_{10} in an NMR tube confirmed the evolution of ethylene, observed in 26% ¹H NMR yield after 24 h at 160 °C and in 31% yield after leaving the resulting mixture at ambient temperature for another day (Fig. 3b). Results of the RC with 2-substituted cyclopropanes also confirmed the liberation of alkene. Thus, a cyclopropane possessing a cis-decyl substituent at the 2position (3_{cis}) was cleanly converted to 2a in 87% yield with dodecenes detected in 87% total yield (Fig. 3c). Similarly, a cyclopropane with a *trans*-decyl substituent (3_{trans}) afforded 2a in 57% yield with dodecenes detected in 54% total yield.

The proposed mechanism commences with coordination of a pyridyl group to the active molybdenum species generated in situ from Mo(0) and o-quinone (Fig. 3d). The regioselective cleavage of the proximal C-C bond of the cyclopropane ring affords the molybdenacyclobutane intermediate **(I)**, which undergoes a second cleavage of the C-C bond with release of ethylene to form the molybdenum carbene species (II). The C-N bond forming intramolecular cyclization finally yields the pyridoisoindole product and regenerates the active species.



Figure 3. Molybdenum-catalyzed retrocyclopropanation with release of alkene and the proposed mechanism. *Yield of ethylene determined after leaving the resulting reaction mixture in an NMR tube at ambient temperature for another day. Ph, phenyl.

The proposed mechanism was further corroborated by density functional theory (DFT) calculations using $Mo/L1/C_2H_4$ and 2-[2-(1methylcyclopropyl)phenyl]pyridine as a model metal complex and substrate, respectively (Fig. 4). Calculations with other possible conformations and model systems are provided in the Supporting Information. Gibbs free energies and enthalpies calculated at the UM06/SDD:6-31G(d,p) level of theory in mesitylene at 160 °C are shown here. The proximal selective cleavage of the C-C single bond proceeds from complex A with triplet ground state to afford molybdenacyclobutane **B** in its singlet state, during which spin crossover takes place. After this point, the singlet potential energy surface of the productive reaction path mostly lies lower than the triplet energy surface. The [2+2] cycloreversion from **B** affords molybdenum carbene C with coordinated ethylene. Dissociation of a pyridyl group from the molybdenum center (C to $\mathbf{D}_{singlet}$) and its intramolecular nucleophilic attack to the carbene affords the cyclized product

2

3

4

5

6 7 8

9

10

11

12

13 14

15

16

17

18

19

20

21

22 23

24

25 26

27

28

29

30

31

32

33

34

35

36

37

 $(\mathbf{D}_{singlet} \text{ to } \mathbf{E})$.⁴³ The transition state Gibbs energies of each step relative to **A** were calculated to be 14.7, 5.9, and 14.3 kcal/mol, respectively.



Figure 4. DFT-examined paths for the retrocyclopropanation. Gibbs free energies and enthalpies (italicized) at 160 °C are given in kcal/mol, calculated at the UM06/SDD:6-31G(d,p)mesitylene(SMD) level of theory. *Ref. 43.

CONCLUSION

We have demonstrated that cyclopropanes can be used as uncommon C1 carbene units by releasing ethylene with a simple molybdenum/quinone catalyst. The strategic use of a pyridyl group with the catalyst enabled the efficient RC of cyclopropanes without extra strain, and the convenient synthesis of pyridoisoindole derivatives, compounds of interest for materials science. Although diazo compounds have played a leading role as sources of carbene species in organic synthesis, their inherent explosive and toxic nature continues to demand stable and safe surrogates for the generation of carbene species. The C1-fragment cyclopropane chemistry described here adds to the growing repertoire of nondiazo approaches to carbene species.23 Investigations to expand the scope of cyclopropanes for the RC reactions by fine tuning of quinone ligands, are ongoing to further explore the applications of cyclopropanes as diazo surrogates.

ASSOCIATED CONTENT

Supporting Information. Experimental procedures, DFT studies, physical properties of the compounds, and NMR spectra. This material is available free of charge via the Internet at http://pubs.acs.org.

AUTHOR INFORMATION

Corresponding Authors

*asako@cc.okayama-u.ac.jp *ktakai@cc.okayama-u.ac.jp

ORCID

Sobi Asako: 0000-0003-4525-5317 Kazuhiko Takai: 0000-0002-2572-0851

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank MEXT for financial support (KAKENHI Grantin-Aid for Scientific Research (A) No. 26248030 to K.T. and Challenging Research (Exploratory) No. 17K19122 to S.A.). The generous amount of computational time from the Research Center for Computational Science, Okazaki, is gratefully acknowledged.

REFERENCES

- ¹ Faust, R. Fascinating natural and artificial cyclopropane architectures. Angew. Chem. Int. Ed. 2001, 40, 2251–2253.
- ² Wessjohann, L. A.; Brandt, W. Biosynthesis and metabolism of cyclopropane rings in natural compounds. Chem. Rev. 2003, 103, 1625–1647.
- ³ Pietruszka, J. Synthesis and properties of oligocyclopropyl-containing natural products and model compounds. Chem. Rev. 2003, 103, 1051-1070.
- ⁴ de Meijere, A.; Kozhushkov, S. I.; Schill, H. Three-membered-ring-based molecular architectures. Chem. Rev. 2006, 106, 4926–4996.
- ⁵ Chen, D. Y.-K.; Pouwer, R. H.; Richard, J.-A. Recent advances in the total synthesis of cyclopropane-containing natural products. *Chem. Soc.* Rev. 2012, 41, 4631-4642.
- ⁶ Taylor, R. D.; MacCoss, M.; Lawson, A. D. G. Rings in drugs. J. Med. Chem. 2014, 57, 5845-5859. 15
 - ⁷ Talele, T. T. The "cyclopropyl fragment" is a versatile player that frequently appears in preclinical/clinical drug molecules. J. Med. Chem. 2016, 59.8712-8756.
 - ⁸ Ebner, C.; Carreira, E. M. Cyclopropanation strategies in recent total syntheses. Chem. Rev. 2017, 117, 11651–11679.
 - ⁹ Rubin, M.; Rubina, M.; Gevorgyan, V. Transition metal chemistry of cyclopropenes and cyclopropanes. *Chem. Rev.* 2007, 107, 3117–3179.
- ¹⁰ Carson, C. A.; Kerr, M. A. Heterocycles from cyclopropanes; applications in natural product synthesis. *Chem. Soc. Rev.* **2009**, *38*, 3051–3060. 19
 - ¹¹ Tang, P.; Qin, Y. Recent applications of cyclopropane-based strategies to natural product synthesis. Synthesis 2012, 44, 2969–2984.
 - ¹² Mack, D. J.; Njardarson, J. T. Recent advances in the metal-catalyzed ring expansions of three- and four-membered rings. ACS Catal. 2013, 3, 272-286.
- 22 ¹³ Murakami, M.; Chatani, N. Eds., Cleavage of Carbon–Carbon Single Bonds by Transition Metals (Wiley-VCH, Weinheim, 2015). 23
 - ¹⁴ Kulinkovich, O. G. Ed., Cyclopropanes in Organic Synthesis (Wiley-VCH, Hoboken, 2015).
- 24 ¹⁵ Fumagalli, G.; Stanton, S.; Bower, J. F. Recent methodologies that exploit C-C single-bond cleavage of strained ring systems by transition metal complexes. Chem. Rev. 2017, 117, 9404-9432. 25
- ¹⁶ Souillart, L.; Cramer, N. Catalytic C-C bond activations via oxidative addition to transition metals. Chem. Rev. 2015, 115, 9410–9464. 26
 - ¹⁷ de Meijere, A. Bonding properties of cyclopropane and their chemical consequences. Angew. Chem. Int. Ed. Engl. **1979**, 18, 809–826.
- 27 ¹⁸ Wiberg, K. B. The concept of strain in organic chemistry. Angew. Chem. Int. Ed. Engl. 1986, 25, 312–322.
- 28 ¹⁹ Khoury, P. R.; Goddard, J. D.; Tam, W. Ring strain energies: substituted rings, norbornanes, norbornenes and norbornadienes. *Tetrahedron* 29 2004, 60, 8103-8112. 30
- ²⁰ Schneider, T. F.; Kaschel, J.; Werz, D. B. A new golden age for donor-acceptor cyclopropanes. *Angew. Chem. Int. Ed.* 2014, *53*, 5504–5523. 31
 - ²¹ Khusnutdinov, R. I.; Dzhemilev, U. M. Transition metal complexes in the chemistry of vinylcyclopropanes. J. Organomet. Chem. 1994, 471, 1-18.
- 32 ²² Brandi, A.; Cicchi, Cordero, S. F. M.; Goti, A. Progress in the synthesis and transformations of alkylidenecyclopropanes and 33 alkylidenecyclobutanes. Chem. Rev. 2014, 114, 7317-7420. 34
- ²³ Jia, M.; Ma, Shengming. New approaches to the synthesis of metal carbenes. Angew. Chem. Int. Ed. 2016, 55, 9134–9166. 35
 - ²⁴ Yang, S. F.; Hoffman, N. E. Ethylene biosynthesis and its regulation in higher plants. Annu. Rev. Plant Physiol. 1984, 35, 155–189.
 - ²⁵ Pirrung, M. C. Ethylene biosynthesis from 1-aminocyclopropanecarboxylic acid. Acc. Chem. Res. 1999, 32, 711–718.
- ²⁶ (a) Solorio-Alvarado, C. R.; Wang, Y.; Echavarren, A. M. Cyclopropanation with gold(I) carbenes by retro-Buchner reaction from 37 cycloheptatrienes. J. Am. Chem. Soc. 2011, 133, 11952-11955. (b) Wang, Y.; McGonigal, P. R.; Herlé, B.; Besora, M.; Echavarren, A. M. Gold(I) 38 carbenes by retro-Buchner reaction: generation and fate. J. Am. Chem. Soc. 2014, 136, 801-809. (c) Wang, Y.; Muratore, M. E.; Rong, Z.; 39 Echavarren, A. M. Formal (4+1) cycloaddition of methylenecyclopropanes with 7-aryl-1,3,5-cycloheptatrienes by triple gold(I) catalysis. Angew. 40 Chem. Int. Ed. 2014, 53, 14022-14026. (d) Yin, X.; Mato, M.; Echavarren, A. M. Gold(I)-catalyzed synthesis of indenes and cyclopentadienes:
- 41 access to ()-Laurokamurene B and the skeletons of the cycloaurenones and dysiherbols. Angew. Chem. Int. Ed. 2017, 56, 14591-14595. (e) Herlé,
- B.; Holstein, P. M.; Echavarren, A. M. Stereoselective cis-vinylcyclopropanation via a gold(I)-catalyzed retro-Buchner reaction under mild 42 Conditions. ACS Catal. 2017, 7, 3668-3675. (f) Mato, M.; Herlé, B.; Echavarren, A. M. Cyclopropanation by gold- or zinc-catalyzed retro-43 Buchner reaction at room temperature. Org. Lett. 2018, 20, 4341-4345.
- 44 ²⁷ Takaya, H.; Suzuki, T.; Kumagai, Y.; Hosoya, M.; Kawauchi, H.; Noyori, R. Nickel(0)-catalyzed reactions of bicyclo[1.1.0]butanes. Geminal 45 two-bond cleavage reaction and the stereospecific olefin trapping of the carbenoid intermediate. J. Org. Chem. 1981, 46, 2854-2861.
- 46 ²⁸ Walczak, M. A. A.; Wipf, P. Rhodium(I)-catalyzed cycloisomerizations of bicyclobutanes. J. Am. Chem. Soc. 2008, 130, 6924–6925.
- ²⁹ Yoshizaki, S.; Shibata, Y.; Tanaka, K. Fulvene synthesis by rhodium(I)-catalyzed [2+2+1] cycloaddition: synthesis and catalytic activity of 47 tunable cyclopentadienyl rhodium(III) complexes with pendant amides. Angew. Chem. Int. Ed. 2017, 56, 3590-3593. 48
- ³⁰ Kawamura, T.; Kawaguchi, Y.; Sugikubo, K.; Inagaki, F.; Mukai, C. Rhodium(I)-catalyzed cycloisomerization of allene–allenylcyclopropanes. 49 Eur. J. Org. Chem. 2015, 719-722.
- 50 ³¹ Gassman, P. G.; Johnson, T. H. Retrocarbene additions. Dissection of alkyl-substituted cyclopropanes under metathesis conditions. J. Am. Chem. 51 Soc. 1976, 98, 6057-6058.
- ³² Gassman, P. G.; Johnson, T. H. Cyclopropane–olefin cross metathesis. J. Am. Chem. Soc. 1976, 98, 6058–6059. 52
- ³³ For RC of alkoxycyclopropanes in gas phase, see: Fedorov, A.; Bach, L. A.; Birney, D. M.; Chen, P. Potential energy surface for (retro-53)cyclopropanation: metathesis with a cationic gold complex. J. Am. Chem. Soc. 2011, 133, 12162-12171. 54
- ³⁴ Shang, R.; Ilies, L.; Nakamura, E. Iron-catalyzed C-H bond activation. Chem. Rev. 2017, 117, 9086–9139. 55
 - ³⁵ Kuninobu, Y.; Takai, K. Organic reactions catalyzed by rhenium carbonyl complexes. Chem. Rev. 2011, 111, 1938–1953.
- 56 ³⁶ Jennings, P. W.; Johnson, L. L. Metallacyclobutane complexes of the group eight transition metals: synthesis, characterizations, and chemistry, 57

9

10

11

12

13

14

16

17

18

20

21

36

- Chem. Rev. 1994, 94, 2241-2290.
- ³⁷ (a) Bart, S. C.; Chirik, P. J. Selective, catalytic carbon-carbon bond activation and functionalization promoted by late transition metal catalysts, J. Am. Chem. Soc. 2003, 125, 886-887. (b) Shaw, M. H.; Melikhova, E. Y.; Kloer, D. P.; Whittingham, W. G.; Bower, J. F. Directing group enhanced carbonylative ring expansions of amino-substituted cyclopropanes: rhodium-catalyzed multicomponent synthesis of N-heterobicyclic enones, J. Am. Chem. Soc. 2013, 135, 4992-4995. (c) Kondo, H.; Itami, K.; Yamaguchi, J. Rh-catalyzed regiodivergent hydrosilylation of acylaminocyclopropanes controlled by monophosphine ligands, Chem. Sci. 2017, 8, 3799-3803. (d) Murai, M.; Nishiyama, A.; Nishinaka, N.; Morita, H.; Takai, K. Iridium-catalysed hydrosilylation of cyclopropanes via regioselective carbon-carbon bond cleavage, Chem. Commun. 2017, 53, 9281-9284.
- ³⁸ Grubbs, R. H.; Wenzel, A. G.; O'Leary, D. J.; Khosravi, E. Eds., Handbook of Metathesis (Wiley, Weinheim, 2014).
 - ³⁹ Hoveyda, A. H.; Zhugralin, A. R. The remarkable metal-catalysed olefin metathesis reaction. *Nature* **2007**, *450*, 243–251.
- ⁴⁰ Asako, S.; Ishikawa, S.; Takai, K. Synthesis of linear allylsilanes via molybdenum-catalyzed regioselective hydrosilylation of allenes. ACS. Catal. 2016, 6, 3387-3395.
- ⁴¹ Asako, S.; Sakae, T.; Murai, M.; Takai, K. Molybdenum-catalyzed stereospecific deoxygenation of epoxides to alkenes. Adv. Synth. Catal. 2016, 358, 3966-3970.
- 42 (a) Mitsumori, T.; Bendikov, M.; Dautel, O.; Wudl, F.; Shioya, T.; Sato, H.; Sato, Y. Synthesis and properties of highly fluorescent indolizino[3,4,5-ab]isoindoles. J. Am. Chem. Soc. 2004, 126, 16793-16803. (b) Yang, D.-T.; Radtke, J.; Mellerup, S. K.; Yuan, K.; Wang, X.; Wagner, M.; Wang, S. One-pot synthesis of brightly fluorescent Mes; B-functionalized indolizine derivatives via cycloaddition reactions. Org. Lett. 2015, 17, 2486–2489.
- ⁴³ Molybdenum carbene complex **D** without coordination of the pyridyl group could not be located at triplet spin state. Instead, molybdenacyclobutane complex D' without pyridyl coordination was found.

