

This article was downloaded by: [UQ Library]

On: 11 November 2014, At: 20:03

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Incn20>

STEREOSELECTIVE SYNTHESIS OF CONFORMATIONALLY RIGID APIO CARBANUCLEOSIDES AS POTENTIAL ANTIVIRAL AGENTS

Hyung Ryong Moon ^a, Kyung Ran Kim ^a, Bum Tae Kim ^b, Ki Jun Hwang ^b, Moon Woo Chun ^c & Lak Shin Jeong ^d

^a College of Pharmacy, Pusan National University, Pusan, Korea

^b Department of Chemistry, Chonbuk National University, Chonju, Korea

^c College of Pharmacy, Ewha Womans University, Seoul, Korea

^d College of Pharmacy, Seoul National University, Seoul, Korea

Published online: 15 Nov 2011.

To cite this article: Hyung Ryong Moon, Kyung Ran Kim, Bum Tae Kim, Ki Jun Hwang, Moon Woo Chun & Lak Shin Jeong (2005) STEREOSELECTIVE SYNTHESIS OF CONFORMATIONALLY RIGID APIO CARBANUCLEOSIDES AS POTENTIAL ANTIVIRAL AGENTS, *Nucleosides, Nucleotides and Nucleic Acids*, 24:5-7, 709-711, DOI: [10.1081/NCN-200060292](https://doi.org/10.1081/NCN-200060292)

To link to this article: <http://dx.doi.org/10.1081/NCN-200060292>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

STEREOSELECTIVE SYNTHESIS OF CONFORMATIONALLY RIGID APIO CARBANUCLEOSIDES AS POTENTIAL ANTIVIRAL AGENTS

Hyung Ryong Moon and Kyung Ran Kim □ *College of Pharmacy, Pusan National University, Pusan, Korea*

Bum Tae Kim and Ki Jun Hwang □ *Department of Chemistry, Chonbuk National University, Chonju, Korea*

Moon Woo Chun □ *College of Pharmacy, Ewha Womans University, Seoul, Korea*

Lak Shin Jeong □ *College of Pharmacy, Seoul National University, Seoul, Korea*


□ *Apio north-methanocarbo-cyclic nucleosides 1–3 with bicyclo[3.1.0]hexane template were first synthesized. Introduction of hydroxymethyl substituent was efficiently and stereoselectively accomplished by aldol and retro-aldol reaction and fixed conformation was achieved from a modified Simmons-Smith cyclopropanation on a cyclopentane ring.*

INTRODUCTION


Due to the lack of anomeric effect and *gauche* effect, carbocyclic nucleosides have a different conformation at the sugar moiety from that of conventional nucleosides and have seldom showed biological activities. Conventional nucleosides exist in the equilibrium between *south* and *north* conformations in solution. Since it has been suggested that conformation in sugar moiety of nucleosides was closely related with their biological activities, a number of nucleosides with a fixed conformation have been synthesized and evaluated for antiviral activity to identify relationships between conformation and biological activity. Among them, the representative is the nucleoside built on bicyclo[3.1.0]hexane template,^[1] which exists in one conformation, *north* or *south*, and the nucleoside analogues showed significant antiviral activity. As a part of efforts to search for antiviral agents with better biological profile, several groups including our laboratory have studied on

This research was supported by a grant from Korea Science and Engineering Research Foundation (KOSEF-R01-2002-000-00114-0).

Address correspondence to Hyung Ryong Moon, College of Pharmacy, Pusan National University, Pusan 609-735, Korea.



SCHEME 1 Reagents and conditions: (a) acetone, conc. H₂SO₄, rt, 2.5 h; (b) vinylmagnesium bromide, THF, −78°C to 0°C, 3 h; (c) NaIO₄, CH₂Cl₂, H₂O, 0°C to rt, 40 min; (d) K₂CO₃, 37% CH₂O, MeOH, 80°C, 36 h; (e) CH₃PPh₃Br, KO^tBu, THF, rt, overnight; (f) Grubbs catalyst 2nd generation, CH₂Cl₂, rt, 2 h; (g) TrCl, DMAP, pyridine, rt, 20 h; (h) Et₂Zn, CH₂I₂, CH₂Cl₂, rt, 9 h; (i) 6-chloropurine, PPh₃, DEAD, THF, 42°C, 4 h; (j) NH₃/MeOH, 80°C, 8 h, then 50% aq. CF₃CO₂H, THF, rt, 4 d for **1**; 40% methylamine, MeOH, 80°C, 5 h, then 50% aq. CF₃CO₂H, THF, rt, 4 d for **2**; 1N HCl, 1,4-dioxane, reflux, overnight for **3**.



SCHEME 2 The plausible mechanism of hydroxymethylation reaction.

the syntheses and biological activities of apio and isonucleosides. In addition to isonucleosides, apio nucleosides have been reported to show better stability in acidic conditions and potent biological activity.^[2] On the basis of the hypothesis that the fix of conformation in sugar moiety of carbocyclic nucleosides may elicit a biological activity lost and the fact that apio nucleosides have potent biological activity with an increased stability under acidic conditions, we designed and synthesized apio *north*-methanocarbocyclic nucleosides **1**, **2**, and **3**. Lactol **5** was obtained starting from D-ribose via stereoselective Grignard reaction. In order to synthesize apio-type sugar hydroxymethyl substituent was introduced at the C2-position of **5** in stereoselective manner,^[3] as shown in Scheme 1. The expected mechanism may be explained, as depicted in Scheme 2. Carbocyclic sugar **7** was synthesized by RCM reaction using second generation Grubbs catalyst and stereoselective fusion of cyclopropane on a cyclopentane ring to fix conformation was achieved by a modified Simmons-Smith cyclopropanation reaction using diethyl zinc and diiodomethane. The final apio carbanucleosides were synthesized via Mitsunobu reaction as a key step. The antiviral assay of the final nucleosides **1** – **3** is in progress and will be reported in due course.

REFERENCES

1. Altmann, K.-H.; Kesselring, R.; Francotte, E.; Rihs, G. 4',6'-Methano carbocyclic thymidine: a conformationally constrained building block for oligonucleotides. *Tetrahedron Lett.* **1994**, *35*(15), 2331–2334.
2. Terao, Y.; Akamatsu, M.; Achiwa, K. Synthesis of chiral 3-substituted gamma-lactones and 9-furanosyladenine

- from (R)-2-(2,2-diethoxyethyl)-1,3-propanediol monoacetate prepared by lipase-catalyzed reaction. *Chem. Pharm. Bull.* **1991**, *39*(3), 823–825.
3. Ho, P.-T. Branched-chain sugars. I. Reaction between furanoses and formaldehyde: a synthesis of D-hamamelose. *Tetrahedron Lett.* **1978**, *19*(19), 1623–1626.