Formation of a stable dicarbenoid and an unsaturated $C_2P_2S_2$ ring from two-electron oxidation of the $[C(PPh_2S)_2]^{2-}$ dianion[†]

Jari Konu and Tristram Chivers*

Received (in Berkeley, CA, USA) 25th June 2008, Accepted 19th July 2008 First published as an Advance Article on the web 12th September 2008 DOI: 10.1039/b810796c

Two-electron oxidation of the $[C(PPh_2S)_2]^{2-}$ dianion with iodine afforded an unexpected mixture of a dimeric Li–I carbenoid $[(Et_2O)(\mu-Li)][(\mu_4-Li){IC(PPh_2S)_2}_2]$ and a novel, unsaturated six-membered $C_2P_2S_2$ ring in $[(SPh_2P)_2C_2(PPh_2)_2S_2]$.

Our recent systematic studies of the oxidation of dichalcogenidoimidodiphosphinate monoanions $[(EPR_2)_2N]^-$ (1, E = S, Se, Te; $R = {}^{i}Pr$, ${}^{t}Bu$, Chart 1) have revealed new and unanticipated aspects of the chemistry of these well-studied anions.¹ We have shown that one-electron oxidation with iodine produces dimeric dichalcogenides EPR₂NR₂PE-EPR₂NPR₂E (2) with elongated central E–E bonds.^{2,3} In one case (E = Te, R = ${}^{t}Bu$) a structural isomer comprised of a contact ion pair in which a $[(TeP^tBu_2)_2N]^{-1}$ anion is Te, Te'-chelated to one Te atom of an incipient cyclic cation $[(TeP^tBu_2)_2N]^+$ was identified.³ The formally 6π -electron five-membered cations $[(EPR_2)_2N]^+$ (E = S, Se, Te; R = ^{*i*}Pr, ^tBu) are obtained as iodide salts by two-electron oxidation of the corresponding anions.³⁻⁵ By contrast, the oxidation of the phenyl-substituted derivatives $[(EPPh_2)_2N]^-$ (E = Se, Te) with one equivalent of iodine was found to be chalcogendependent; the five-membered ring $[(TePPh_2)_2N]^+$ was obtained for tellurium, whereas the unusual six-membered ring $[(\mu-Se)(SePPh_2)_2N]^+$ was the major product in the case of selenium.5

In the context of these intriguing and diverse results we turned our attention to the oxidative behaviour of the isoelectronic $[C(PPh_2S)_2]^{2-}$ dianion (3), which is prepared from bis(thiodiphenylphosphinoyl)methane upon treatment with two equivalents of methyl-lithium.⁶ Le Floch and co-workers have reported recently that mild oxidation of Li₂3 with hexachloroethane produces the remarkably stable carbenoid $[ClC(PPh_2S)_2]Li(OEt_2)_2$ (4).^{7,8} In this communication we give details of our complementary investigations of the oxidation of Li₂3 with iodine, which have led to the structural and spectroscopic characterisation of the stable dicarbenoid $[(Et_2O)-(\mu-Li)][(\mu_4-Li){IC(PPh_2S)_2}]$ (5) and the novel, unsaturated six-membered $C_2P_2S_2$ ring in $[(SPh_2P)_2C_2(PPh_2)_2S_2]$ (6).

The reaction between Li₂3 and one equivalent of I₂ was initially performed in a toluene–Et₂O mixture at -80 °C (Scheme 1).‡ The ³¹P NMR spectrum of the product in



d₈-THF revealed a major component (*ca.* 75%) that exhibits a singlet at δ 48.2 and a minor product (*ca.* 25%), which gives rise to two mutually coupled doublets at δ 46.4 and 42.1 [²J(³¹P, ³¹P) = 97 Hz]. Both of the new compounds were identified in the solid state by their X-ray crystal structures.†§

Single crystals of the major component were obtained from a toluene solution of the product mixture and identified as the dicarbenoid $[(Et_2O)(\mu-Li)][(\mu_4-Li){IC(PPh_2S)_2}_2]$ (5) (Scheme 1). In contrast to the monomeric structure of 4, the iodide derivative 5 adopts a dimeric arrangement of two molecules of the carbene $[:C(PPh_2S)_2]$ and two molecules of LiI. The two five-atom SPCPS fragments in the C_2 -symmetric dicarbenoid 5, each of which contains a C–I bond, are connected by a tetrahedral Li⁺ cation and a three-coordinate Li⁺ cation solvated by Et₂O (Fig. 1). The spirocyclic Li⁺ cation Li(1) has S–Li–S bond angles spanning a range of 99.8(1) to 119.5(3)°, while the 3-coordinate lithium is in a trigonal planar environment ($\sum \ Li(2)$ 359.9°). The Li(2) centre also engages in significant Li···I close contacts of 3.167(1) Å which form an almost linear



Scheme 1 Formation of 5 and 6 by two-electron oxidation of the dianion $[C(PPh_2S)_2]^{2-}$ (3) with iodine.

University of Calgary, Department of Chemistry, 2500 University Drive NW, Calgary, Alberta, Canada T2N 1N4. E-mail: chivers@ucalgary.ca; Fax: +1-403-289-9488; Tel: +1-403-220-5741 † Electronic supplementary information (ESI) available: Experimental and crystallographic data in pdf format. CCDC reference numbers 692935 (5) and 692936 (6). For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/b810796c



Fig. 1 Crystal structure of **5** with thermal ellipsoid at 50% probability level. Hydrogen atoms have been omitted for clarity. Relevant bond parameters (in Å and °): C(1)–I(1) 2.147(3), C(1)–P(1) 1.750(3), C(1)–P(2) 1.745(3), P(1)–S(1) 1.997(1), P(2)–S(2) 2.022(1), Li(1)–S(1) 2.466(4), Li(1)–S(2) 2.512(5), Li(2)–S(2) 2.461(5), Li(2)– $\cdot\cdot$ I(1) 3.167(1), P(1)–C(1)–P(2) 123.8(2), P(1)–C(1)–I(1) 111.8(2), P(2)–C(1)–I(1) 107.3(2). Symmetry operation (A): -x, y, 0.5 – z.

arrangement [\angle I(1)–Li(2)–I(1A) = 169.6(3)°]. Consequently, the carbon atom in the PCP unit in **5** is pyramidally distorted with the sum of bond angles being 342.9°, *cf.* 359.9° in **4**.⁷ Despite the disparity in the geometry around the carbon center in **4** and **5**, and the strength of the different carbon–halogen bonds, the calculated bond orders for the C–Cl and C–I bonds in **4** and **5**, respectively, are essentially identical (0.90 and 0.92, respectively).⁹ The endocyclic P–C and P–S bond distances in **5** are slightly elongated (by *ca.* 0.02 Å), compared to those in **4**.¹²

Notwithstanding the two inequivalent lithium and phosphorus environments in 5, only a single resonance was observed in both the ⁷Li (δ 0.73) and ³¹P NMR spectra at 23 °C indicating fluctionality in solution. Indeed, when a CD₂Cl₂ solution of the product is cooled to -90 °C, two singlets are observed in the ⁷Li NMR at δ 2.20 and 1.16. The broadness of the latter singlet, however, suggests that unresolved fluctionality persists even at low temperature affording a broad singlet for the 3-coordinate lithium. Consistently, the ³¹P NMR spectrum of 5 exhibits two broad, overlapping singlets rather than the expected two mutually coupled doublets at -90 °C. The ¹³C{¹H} NMR spectrum of 5 in d₈-THF shows a broad triplet at δ 2.4 ppm [¹J(¹³C, ³¹P) = ca. 60 Hz] for the PCP-carbon at 23 °C. The triplet of the PCP-carbon in 5 is shifted to higher field by ca. 36 ppm compared to that in 4 (38.5 ppm).⁷ This upfield shift can be attributed primarily to the greater shielding effect of iodine compared to chlorine,¹³ although the change of hybridization (towards sp³) in 5 compared to that in 4 (carbon lone pair in a pure p orbital)⁷ may also be a contributing factor.

Single crystals of the minor product formed in the reaction depicted in Scheme 1 were obtained from a diethyl ether solution. This compound was identified by X-ray crystallography as $[(SPh_2P)_2C_2(PPh_2)_2S_2]$ (6), which contains a novel, unsaturated six-membered $C_2P_2S_2$ ring in a chair conformation (Fig. 2).†§ This heterocycle can be viewed to result from the union of two $[:C(PPh_2S)_2]$ carbenes without the incorporation of LiI. The crystal structure of **6**·Et₂O contains exocyclic $Ph_2P=S$ units attached to each carbon, one of which is inclined towards the centre of the ring while the second unit



Fig. 2 Molecular structure of $6 \cdot Et_2O$ with thermal ellipsoids at 50% probability level. Hydrogen atoms and Et_2O solvate have been omitted for clarity. Pertinent bond lengths (in Å): C(1)–P(1) 1.762(3), C(1)–P(2) 1.692(3), C(1)–S(2) 1.775(3), C(2)–P(3) 1.766(3), C(2)–P(4) 1.692(3), C(2)–S(2) 1.769(3), P(1)–S(1) 1.975(1), P(2)–S(4) 2.133(1), P(3)–S(3) 1.969(1), P(4)–S(4) 2.137(1).

is bent away from the ring. The structure exhibits both endoand exo-cyclic P–C and P–S bonds with inequivalent bond lengths; the exocyclic P–C bonds are *ca*. 0.07 Å longer and the exocyclic P–S bonds are *ca*. 0.16 Å shorter than the corresponding endocyclic bond lengths, indicating significant double bond character in the endocyclic P—C and exocyclic P—S bonds, respectively, as illustrated in Scheme 1. The exocyclic P—S distances are similar to those reported for the related compounds $[Ph_2C=C(PPh_2S)_2]^{14}$ and $[H(Cl)C(PPh_2S)_2]^{.7}$ The C(1) and C(2) atoms in **6** are only slightly distorted from planarity with the sum of bond angles being 356.2 and 358.5°, respectively.

The formation of the $C_2P_2S_2$ ring in 6 Et₂O from the twoelectron oxidation of **3** involves a sulfur transfer process to generate a CSC unit. A tentative explanation of this observation is shown in Scheme 2. The first step (i) invokes an *inter*molecular nucleophilic attack of a sulfur atom of the initially formed carbene [:C(PPh₂S)₂] on the electron-deficient carbon centre of another carbene to create a C–S bond, accompanied by a



Scheme 2 Suggested reaction mechanism for the formation of the six-membered $C_2P_2S_2$ ring in 6.

rearrangement to afford a five-membered CSPSP ring 7. Step (ii) involves an *intra*molecular attack of a sulfur atom of the five-membered ring on the exocyclic carbene center in 7 with subsequent ring expansion to give the final product **6**.

Attempts were made to improve the yield of **6** by carrying out the oxidation of Li₂**3** with a mixture of iodine and 12-crown-4 (to remove LiI) at -80 °C.† Although this procedure prevented the formation of **5** and increased the yield of **6** to *ca*. 50% (³¹P NMR) it was not possible to separate pure samples of **6** from the other products H₂C(PPh₂S)₂ and H(I)C(PPh₂S)₂. When the reaction of Li₂**3** with iodine was carried out at 23 °C the dicarbenoid **5** was obtained in *ca*. 90% yield (³¹P NMR), together with H₂C(PPh₂S)₂ from which it is difficult to separate; the six-membered ring **6** is not formed under these conditions.†

In summary, participation of the LiI by-product in the twoelectron oxidation of Li₂3 results in the formation of the dimeric carbenoid 5, which exhibits distorted pyramidal carbon centers and a thermal stability comparable to that found in the monomeric Li–Cl carbenoid 4.¹⁵ In the absence of LiI incorporation, this oxidation process produces the novel unsaturated six-membered $C_2P_2S_2$ ring in 6.

Notes and references

‡ Formation of **5** and **6** from Li₂**3** and I₂: Li₂[C(PPh₂S)₂] was synthesized according to the literature.^{6,7} A solution of H₂C(PPh₂S)₂ (0.287 g, 0.64 mmol) in 30 mL of toluene was cooled to -80 °C and 0.80 mL of MeLi (1.6 M in Et₂O, 1.28 mmol) was added *via* syringe. The reaction mixture was stirred for 15 min at -80 °C and $2\frac{1}{2}$ h at room temperature.

The turbid solution of Li₂[C(PPh₂S)₂] was cooled to -80 °C and a solution of I₂ (0.162 g, 0.64 mmol) in 30 mL of Et₂O was added *via* cannula. The reaction mixture was stirred for $\frac{1}{2}$ h at -80 °C and 3 h at room temperature giving an orange-yellow solution. The solvents were evaporated under vacuum and the resulting tarry product was dissolved to 50 mL of Et₂O. White LiI powder was filtered with a PTFE-disk and the solvent was evaporated *in vacuo* to give a yellow, amorphous powder (0.335 g, 91% calculated as a 75:25 mixture of **5** and **6** based on ³¹P NMR data).

and **6** based on P INMR data). NMR data of **5**: ¹H NMR (THF-d₈, 23 °C): δ 7.17–8.02 [m, 40H, C₆H₅]. ¹³C{¹H} NMR; δ 139.5 [(dd) AXX', $J(^{13}C,^{31}P) = 101$ Hz; C_{ipso} of C₆H₅], 133.8 [t, ² $J(^{13}C,^{31}P) = 5.0$ Hz; C_{ortho} of C₆H₅], 130.2 [s; C_{para} of C₆H₅], 127.8 [t, ³ $J(^{13}C,^{31}P) = 6.1$ Hz; C_{meta} of C₆H₅], 2.4 [br, t, ¹ $J(^{13}C,^{31}P) = ca.$ 60 Hz; PCP-carbon]. ³¹P NMR: δ 48.2 ppm. ⁷Li NMR: δ 0.73 ppm. ³¹P NMR (CD₂Cl₂, -90 °C): δ 5.00 and 49.9 ppm (br, overlapping). ⁷Li NMR (CD₂Cl₂, -90 °C): δ 2.20 and 1.16 (br) ppm.

NMR data of **6**: ³¹P NMR (THF-d₈, 23 °C): δ 46.4 [d, ²J(³¹P, ³¹P) = 97 Hz] and 42.1 [d, ²J(³¹P, ³¹P) = 97 Hz] ppm. ¹H and ¹³C{¹H} NMR data for **6** could not be obtained due to the small quantity of the compound in the product mixture.

§ Crystal data of **5**: C₅₄H₅₀I₂Li₂OP₄S₄, $M_r = 1234.74$, monoclinic, space group C2/c, a = 26.997(5), b = 10.517(2), c = 20.050(4) Å, $\beta = 109.28(3)^{\circ}$, V = 5373(2) Å³, Z = 4, $\rho_{calcd} = 1.526$ g cm⁻³, $\mu = 1.483$ mm⁻¹, T = 173(2) K, 8595 reflections collected (θ range = 2.99–25.03°), 4707 unique ($R_{int} = 0.0284$), $R_1 = 0.0312$ [for 3 694

reflections with $I > 2\sigma(I)$] and $wR_2 = 0.0629$ (for all data). Crystal data of **6**·Et₂O: C₅₄H₅₀OP₄S₄, $M_r = 967.06$, monoclinic, space group $P_{2_1/c}$, a = 18.496(4), b = 11.907(2), c = 23.533(5) Å, $\beta = 108.68(3)^{\circ}$, V = 4910(2) Å³, Z = 4, $\rho_{calcd} = 1.308$ g cm⁻³, $\mu = 0.363$ mm⁻¹, T = 173(2) K, 16 547 reflections collected (θ range = 2.46–25.03°), 8 634 unique ($R_{int} = 0.0432$), $R_1 = 0.0484$ [for 6 137 reflections with $I > 2\sigma(I)$] and $wR_2 = 0.1149$ (for all data). The structures were solved and refined by using SHELXS-97 and SHELXL-97.¹⁶

- For recent reviews, I. Haiduc, in *Comprehensive Coordination Chemistry II*, ed. J. A. McCleverty and T. J. Meyer, Elsevier Ltd., Amsterdam, 2003, pp. 323–347; C. Silvestru and J. E. Drake, *Coord. Chem. Rev.*, 2001, 223, 117.
- 2 T. Chivers, D. J. Eisler, J. S. Ritch and H. M. Tuononen, *Angew. Chem.*, *Int. Ed.*, 2005, **44**, 4953.
- 3 T. Chivers, D. J. Eisler, J. S. Ritch and H. M. Tuononen, *Chem.-Eur. J.*, 2007, **13**, 4643; T. Chivers, J. Konu, J. S. Ritch, M. C. Copsey, D. J. Eisler and H. M. Tuononen, *J. Organomet. Chem.*, 2007, **692**, 2658.
- 4 J. Konu, T. Chivers and H. M. Tuononen, *Chem. Commun.*, 2006, 1634.
- 5 J. Konu, T. Chivers and H. M. Tuononen, *Inorg. Chem.*, 2006, 45, 10678.
- 6 T. Cantat, N. Mézailles, L. Ricard, Y. Jean and P. Le Floch, *Angew. Chem., Int. Ed.*, 2004, **43**, 6382; T. Cantat, L. Ricard, Y. Jean, P. Le Floch and N. Mézailles, *Organometallics*, 2006, **25**, 4965.
- 7 T. Cantat, X. Jacques, L. Ricard, X. F. Le Goff, N. Mézailles and P. Le Floch, *Angew. Chem., Int. Ed.*, 2007, **46**, 5947.
- 8 For a review on stable PCP-carbenes, see: T. Cantat, N. Mézailles, A. Auffrant and P. Le Floch, *Dalton Trans.*, 2008, 1957.
- 9 The C–Cl and C–I bond orders were calculated by the Pauling equation $N = 10^{(D R)/0.71}$ ¹⁰ where R is the observed bond length (Å). The single bond length D is estimated from the sums of appropriate covalent radii:^{10,11} C–Cl 1.75 and C–I 2.12 Å. A value of 0.73 Å was used for sp² hybridized carbons. If the covalent radii of 0.76 Å is used to indicate sp³ hybridized carbon, a bond order of 1.01 is obtained for **5**.
- 10 L. Pauling, *The Nature of the Chemical Bond*, Cornell University Press, Ithaca, NY, 3rd edn, 1960.
- 11 B. Cordero, V. Gomez, A. E. Platero-Prats, M. Reves, J. Echeverria, E. Cremades, F. Barragan and S. Alvarez, *Dalton Trans.*, 2008, 2832.
- 12 The crystal structure of **4** was determined at 150 K⁷ whereas the structure of **5** was measured at 173 K \dagger .
- 13 For example, the ¹³C NMR chemical shift of CH₃I appears *ca.* 46 ppm upfield from that of CH₃Cl and the shift for CH₃CH₂I is found *ca.* 40 ppm upfield from that of CH₃CH₂Cl: J. W. Emsley, J. Feeney and L. H. Sutcliffe, *High Resolution NMR Spectroscopy*, Pergamon Press Ltd., Oxford, 1966, vol. 2, pp. 990–997.
- 14 T. Cantat, F. Jaroschik, F. Nief, L. Ricard, N. Mezailles and P. Le Floch, *Chem. Commun.*, 2005, 5178.
- 15 The monomeric carbenoid **4** was reported to be stable in solution and in the solid state up to 60 °C.⁷ The dimeric carbenoid **5** showed no decomposition when the solid material was kept at 95 °C under vacuum (*ca.* 10^{-3} Torr) for several hours.
- 16 G. M. Sheldrick, SHELXS-97, Program for solution of crystal structures, University of Göttingen, Germany, 1997; G. M. Sheldrick, SHELXL-97, Program for refinement of crystal structures, University of Göttingen, Germany, 1997.