



## Asymmetric total synthesis of reported structure of eudistomidin B, an indole alkaloid isolated from a marine tunicate

Takeshi Ito, Mariko Kitajima, Hiromitsu Takayama \*

Graduate School of Pharmaceutical Sciences, Chiba University, 1-33 Yayoi-cho, Inage-ku, Chiba 263-8522, Japan

### ARTICLE INFO

#### Article history:

Received 21 April 2009

Revised 18 May 2009

Accepted 21 May 2009

Available online 25 May 2009

#### Keywords:

Indole

Alkaloid

Tetrahydro- $\beta$ -carboline

Spectroscopy

Asymmetric total synthesis

### ABSTRACT


Eudistomidin B, isolated from a marine tunicate, was originally assigned a tetrahydro- $\beta$ -carboline structure with a 2-*p*-tolylethanamine residue. Asymmetric total synthesis of the reported structure of natural product, which features an intramolecular diastereoselective Pictet-Spengler cyclization, suggests that the reported structure of the natural product needs to be revised.

© 2009 Elsevier Ltd. All rights reserved.

In connection with our chemical and pharmacological studies on the  $\beta$ -carboline class of indole alkaloids,<sup>1</sup> we are interested in the marine indole alkaloid eudistomidin B (**1**),<sup>2</sup> which was isolated from the Okinawan tunicate *Eudistoma glaucus* and has unique biological activities, including potent cytotoxicity to murine leukemia L1210 and L5178Y cells and the ability to activate rabbit heart muscle actomyosin ATPase. This natural product has a novel structure composed of a brominated tetrahydro- $\beta$ -carboline and a unique 2-*p*-tolylethanamine residue, which are joined together at the C-1 position of the former skeleton and the  $\alpha$  position of the tolylethanamine residue, generating consecutive chiral centers at C-1 and C-10 positions. The intriguing chemical structures and interesting biological activities prompted us to embark on the asymmetric total synthesis of eudistomidin B (**1**).

Our retrosynthesis, which is shown in Scheme 1, was based on the intramolecular Pictet-Spengler cyclization strategy that was successfully used in our synthesis of subincanadines.<sup>3</sup> Stereoselective construction of the tetrahydro- $\beta$ -carboline skeleton in **2** was expected to be accomplished by the intramolecular Pictet-Spengler reaction using aldehyde **3** that tethered 5-bromotryptamine (**4**) and optically active aldehyde prepared from diol **5**. Transformation of the secondary hydroxyl group at C-10 in **2** into the amino function, accompanied by inversion of the stereochemistry, would furnish target molecule **1**.

The synthesis began with the preparation of optically active diol **5** (Scheme 2) from *p*-tolualdehyde via a seven-step operation that




Scheme 1. Retrosynthetic analysis.

included the proline-catalyzed asymmetric aminooxylation<sup>4</sup> of aldehyde **6**. Reductive cleavage of the N–O bond in **7** using  $\text{PtO}_2$  under  $\text{H}_2$  atmosphere gave diol **5** (mp 65–67 °C,  $[\alpha]_D^{23} +17.4$  (c 1.0,  $\text{CHCl}_3$ )) in 82% yield, which showed 99% ee as determined by chiral HPLC column chromatography. After selective protection of the

\* Corresponding author. Tel./fax: +81 43 290 2901.


E-mail address: [htakayam@p.chiba-u.ac.jp](mailto:htakayam@p.chiba-u.ac.jp) (H. Takayama).



Scheme 2.

primary alcohol in diol **5** with a TBS group, the remaining hydroxyl group was condensed with 1,1'-carbonyldiimidazole (CDI) to give carbamate **8**, which was then treated with 5-bromotryptamine (**4**)<sup>5</sup> in DCM in the presence of DMAP, followed by PTS in MeOH, to afford carbamate **9** in 75% overall yield from **5**.

Next, the primary alcohol in **9** was oxidized with IBX to give aldehyde **3** in 87% yield (Scheme 3). Compound **3** was found to exist in hemiaminal form **3'** in solution, as demonstrated by the chemical shift appearing at  $\delta$  82 (C-1) in the <sup>13</sup>C NMR spectrum. Treatment of **3** with TFA in DCM at 0 °C gave the cyclization products in 98% yield with a diastereomeric ratio of 4:1, which could be separated by column chromatography. The stereochemistries of major product **2**<sup>6</sup> as well as minor product **2'** were determined by NOE experiments (Fig. 1): irradiation of the proton at C-1 caused a significant signal enhancement of the proton on the methylene group (C-11) of the side chain in **2** and of the proton at C-10 in min-



Scheme 3.





Figure 1.

or product **2'**, respectively, demonstrating the trans relationship of the two protons at C-1 and C-10 in the major product **2**. Diastereoselectivity of the intramolecular Pictet-Spengler reaction of **3**



Scheme 4.

**Table 1**<sup>1</sup>H and <sup>13</sup>C NMR data for synthetic and natural<sup>2</sup> eudistomidin B in CDCl<sub>3</sub> + CF<sub>3</sub>CO<sub>2</sub>D (10:1)

Position	$\delta_{\text{H}}$ (500 MHz)		$\delta_{\text{C}}$ (125 MHz)	
	Synthetic	Natural	Synthetic	Natural
1	5.28 (s)	5.19 (d, 8.9)	62.77	63.59
3	3.88 (m)	3.28 (dd, 12.2, 5.3)	50.36	45.80
4	3.77 (br dd, 14.0, 4.0)	2.86 (m)	15.29	15.00
	3.21 (2H, overlapped)	2.82 (m)		
4a		2.38 (dd, 15.6, 4.2)	107.27	105.53
4b			126.74	122.40
5	7.71 (d, 1.7)	7.52 (d, 1.6)	121.40	121.05
6			114.51	113.61
7	7.46 (dd, 8.7, 1.7)	7.39 (dd, 8.7, 1.6)	127.96	126.91
8	7.34 (d, 8.7)	7.42 (d, 8.7)	113.67	113.87
8a			136.05	135.49
9		11.08 (s)		
9a			121.24	137.57
10	4.56 (br d, 9.8)	4.18 (m)	56.75	63.73
11	3.21 (overlapped)	3.23 (dd, 14.7, 3.1)	34.32	33.27
12	3.09 (m)	3.03 (dd, 14.7, 9.7)	128.69	134.10
13, 17	6.97 (2H, d, 7.9)	7.05 (2H, d, 7.5)	128.58	128.87
14, 16	7.12 (2H, d, 7.9)	6.59 (2H, d, 7.5)	130.49	128.52
15			139.47	134.10
18	2.29 (3H, s)	2.88 (3H, s)	20.72	32.42
19	3.15 (3H, s)	2.92 (3H, s)	42.54	39.75

could be improved to 16:1 by using TMSCl as acid. The high diastereoselectivity of the cyclization could be due to the plausible acylium intermediate, in which the indole nucleus would attack from the less hindered side (anti from the side chain), as depicted in Figure 1.

To accomplish the total synthesis, we next investigated the introduction of an amino function to the C-10 position, which is accompanied by the inversion of stereochemistry. The carbonyl group in **2** was removed by alkaline hydrolysis and the resulting secondary amine in **10** was protected as carbamate or sulfonamide (Scheme 4). All attempts to conduct the nucleophilic substitution of the free alcohol in **11a** or **11b** under Mitsunobu conditions or of the C-10-O-mesyl derivatives prepared from **11a** or **11b** with sodium azide gave unsuccessful results. Then, we modified the cyclic sulfamate strategy developed by White and Garst<sup>7</sup> and applied it to the synthesis. Treatment of amino alcohol **10** with 1.2 equiv of sulfonyl chloride in DCM at –78 °C in the presence of Et<sub>3</sub>N for 2 h gave unstable cyclic sulfamate **12**. To this mixture was added 10 equiv of sodium azide in DMF. The reaction mixture was heated at 60 °C to facilitate nucleophilic ring opening with the azide group at C-10. After cooling to room temperature, the reaction mixture was treated with 10% aqueous sulfuric acid to hydrolyze resulting sulfamidic acid intermediate **13** to give azidoamine derivative **14** in 26% overall yield. By careful treatment and column chromatography, unstable cyclic sulfamate **12**<sup>8</sup> could be isolated in low yield (9–15%) and then transformed to **14** with the same procedure as above-mentioned in 64% yield. Finally, reductive methylation of the secondary amine in **14** and successive reduction of the azide function with Red-Al furnished the target molecule in 60% yield (2 steps).

The structure of synthetic compound **1**<sup>9</sup> ( $[\alpha]_{\text{D}}^{25}$  free base: +42.0 (c 0.57, CHCl<sub>3</sub>), TFA salt: +63.4 (c 0.78, CHCl<sub>3</sub>)) was confirmed by detailed spectroscopic analyses, including 2D-NMR (COSY, HMQC,

and HMBC) spectroscopy. However, significant differences were noted between our NMR data (see Table 1) and those reported for the natural product ( $[\alpha]_{\text{D}}^{25}$  –76.4 (c 0.3, CHCl<sub>3</sub>)). In particular, the chemical shifts at H-10, 13, 14, and 18 in the <sup>1</sup>H NMR spectra and at C-10 and 18 in the <sup>13</sup>C NMR spectra showed marked differences, as depicted in Scheme 4. Hence, we concluded that natural eudistomidin B does not have structure **1**, and inconsistencies in the data reported for the natural product hindered us from proposing a credible alternative structure.

## Acknowledgment

This work was supported by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science.

## References and notes

- (a) Takayama, H.; Kitajima, M.; Kogure, N. *Curr. Org. Chem.* **2005**, *9*, 1445–1464; (b) Takayama, H.; Misawa, K.; Okada, N.; Ishikawa, H.; Kitajima, M.; Hatori, Y.; Murayama, T.; Wongseripipatana, S.; Tashima, K.; Matsumoto, K.; Horie, S. *Org. Lett.* **2006**, *8*, 5705–5708; (c) Kitajima, M. *J. Nat. Med.* **2007**, *61*, 14–23; (d) Nakayama, A.; Kogure, N.; Kitajima, M.; Takayama, H. *Heterocycles* **2008**, *76*, 861–865.
- Kobayashi, J.; Cheng, J.-F.; Ohta, T.; Nozoe, S.; Ohizumi, Y.; Sasaki, T. *J. Org. Chem.* **1990**, *55*, 3666–3670.
- Suzuki, K.; Takayama, H. *Org. Lett.* **2006**, *8*, 4605–4608.
- Zhong, G. *Angew. Chem., Int. Ed.* **2003**, *42*, 4247–4250.
- 5-Bromotryptamine was prepared from 5-bromoindole via a three-step operation: (i) POCl<sub>3</sub>, DMF, 40 °C, y. 94%; (ii) NH<sub>4</sub>OAc, MeNO<sub>2</sub>, reflux, y. 74%; (iii) NaBH<sub>4</sub>, BF<sub>3</sub>–Et<sub>2</sub>O, THF, reflux, y. 77%.
- Compound **2**: mp 219–221 °C (from EtOAc);  $[\alpha]_{\text{D}}^{25}$  +1.7 (c 0.87, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.53 (1H, d, *J* = 1.9 Hz, H-5), 7.38 (2H, d, *J* = 8.0 Hz, H-14, H-16), 7.31 (2H, d, *J* = 8.0 Hz, H-13, H-17), 7.17 (1H, dd, *J* = 8.8, 1.9 Hz, H-7), 6.70 (1H, d, *J* = 8.8 Hz, H-8), 5.52 (1H, br s, N<sub>9</sub>-H), 4.80 (1H, d, *J* = 8.1, 1.8, 1.8 Hz, H-1), 4.46 (1H, ddd, *J* = 11.0, 8.1, 4.5 Hz, H-10), 4.26 (1H, dd, *J* = 13.5, 5.2 Hz, H-3), 3.61 (1H, dd, *J* = 13.0, 4.5 Hz, H-11), 3.09 (1H, ddd, *J* = 13.5, 11.5, 5.0 Hz, H-3), 2.98 (1H, dd, *J* = 13.0, 11.0 Hz, H-11), 2.78 (1H, dddd, *J* = 15.4, 11.5, 6.1, 2.2 Hz, H-4), 2.69 (1H, br dd, *J* = 15.4, 5.0 Hz, H-4), 2.53 (3H, s, H<sub>3</sub>-18); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  156.1 (C-19), 138.1 (C-15), 134.5 (C-8a), 131.6 (C-12), 130.5 (C-9a), 130.3 (C-14, C-16), 130.1 (C-13, C-17), 128.0 (C-4b), 125.2 (C-7), 121.2 (C-5), 112.9 (C-6), 112.1 (C-8), 108.5 (C-4a), 80.8 (C-10), 56.6 (C-1), 40.0 (C-11), 38.7 (C-3), 21.1 (C-18), 20.7 (C-4); UV (MeOH)  $\lambda_{\text{max}}$  nm 300 (sh), 291, 229; IR (ATR)  $\nu_{\text{max}}$  cm<sup>–1</sup> 3230, 2363, 1715, 1417, 1233, 789; EIMS *m/z* (%) 412 (100, M<sup>+</sup>+2), 410 (99, M<sup>+</sup>), 306 (98), 304 (92); HREIMS *m/z* 410.0629 (M<sup>+</sup>, calcd for C<sub>21</sub>H<sub>19</sub>N<sub>2</sub>O<sub>2</sub><sup>79</sup>Br, 410.0629).
- White, G. J.; Garst, M. E. *J. Org. Chem.* **1991**, *56*, 3177–3178.
- Compound **12**:  $[\alpha]_{\text{D}}^{25}$  +35.2 (c 0.18, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.54 (1H, d, *J* = 1.8 Hz, H-5), 7.39 (2H, d, *J* = 8.0 Hz), 7.31 (2H, d, *J* = 8.0 Hz), 7.18 (1H, dd, *J* = 8.6, 1.8 Hz, H-7), 6.63 (1H, d, *J* = 8.6 Hz, H-8), 5.45 (1H, br s, N<sub>9</sub>-H), 4.91 (1H, d, *J* = 7.8 Hz, H-1), 4.74 (1H, ddd, *J* = 10.9, 7.8, 4.6 Hz), 3.64 (2H, overlapped), 3.48 (1H, ddd, *J* = 12.8, 7.9, 4.7 Hz), 3.16 (1H, dd, *J* = 13.1, 10.9 Hz), 2.97 (1H, m), 2.74 (1H, m), 2.55 (3H, s, H<sub>3</sub>-18); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  136.6, 134.7, 134.3, 133.9, 129.41, 129.36, 128.7, 124.6, 120.8, 112.5, 112.3, 110.0, 67.2, 54.3, 41.6, 37.1, 22.5, 21.1 (C-18); UV (MeOH)  $\lambda_{\text{max}}$  nm 299 (sh), 291 (sh), 282, 229; IR (ATR)  $\nu_{\text{max}}$  cm<sup>–1</sup> 3384, 2920, 2850, 1715, 1518, 1441, 1317, 1312, 1176; EIMS *m/z* (%) 448 (32, M<sup>+</sup>+2), 446 (30, M<sup>+</sup>), 368 (9), 366 (10), 250 (96), 248 (100); HREIMS *m/z* 446.0300 (M<sup>+</sup>, calcd for C<sub>20</sub>H<sub>19</sub>N<sub>2</sub>O<sub>3</sub><sup>79</sup>Br, 446.0299).
- Synthetic **1**: as free base  $[\alpha]_{\text{D}}^{25}$  +42.0 (c 0.57, CHCl<sub>3</sub>); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  9.42 (1H, br s, N<sub>9</sub>-H), 7.62 (1H, d, *J* = 1.7 Hz, H-5), 7.19 (1H, dd, *J* = 8.5, 1.7 Hz, H-7), 7.18 (1H, d, *J* = 8.5 Hz, H-8), 7.13 (2H, d, *J* = 8.1 Hz, H-14, H-16), 7.12 (2H, d, *J* = 8.1 Hz, H-13, H-17), 3.41 (1H, dd, *J* = 13.7, 2.7 Hz, H-11), 3.38 (1H, br d, *J* = 8.9 Hz, H-1), 3.22 (1H, ddd, *J* = 13.7, 8.5, 5.1 Hz, H-3), 3.10 (2H, overlapped, H-3, H-10), 2.83 (1H, m, H-4), 2.60 (1H, br d, *J* = 15.3 Hz, H-4), 2.49 (1H, dd, *J* = 13.7, 9.5 Hz, H-11), 2.47 (3H, s, H<sub>3</sub>-19), 2.34 (3H, s, H<sub>3</sub>-18); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  135.93, 135.80, 135.27, 134.20, 129.37, 129.31, 128.92, 123.87 (C-7), 120.63 (C-5), 112.31 (C-8), 112.16, 107.44, 63.46 (C-1), 55.98 (C-10), 47.88 (br, C-3), 41.51 (C-11), 39.81 (br, C-19), 21.01 (C-18), 16.36 (C-4); UV (MeOH)  $\lambda_{\text{max}}$  nm 300 (sh), 291, 230; IR (ATR)  $\nu_{\text{max}}$  cm<sup>–1</sup> 2920, 2844, 1576, 1440, 1305, 1020; EIMS *m/z* (%) 399 (1, M<sup>+</sup>+2), 397 (1, M<sup>+</sup>), 265 (97), 263 (100); HREIMS *m/z* 397.1153 (M<sup>+</sup>, calcd for C<sub>21</sub>H<sub>24</sub>N<sub>3</sub><sup>79</sup>Br, 397.1153). As TFA salt  $[\alpha]_{\text{D}}^{25}$  +63.4 (c 0.78, CHCl<sub>3</sub>); <sup>1</sup>H and <sup>13</sup>C NMR data, see Table 1; UV (MeOH)  $\lambda_{\text{max}}$  nm 301 (sh), 292, 229; IR (ATR)  $\nu_{\text{max}}$  cm<sup>–1</sup> 2925, 2855, 1668, 1444, 1139, 797.