### Macrocyclic scaffolds derived from *p*-aminobenzoic acid<sup>†</sup>

Fred Campbell,<sup>*a*</sup> Jeffrey Plante,<sup>*a*</sup> Christopher Carruthers,<sup>*a*</sup> Michaele J. Hardie,<sup>*a*</sup> Timothy J. Prior<sup>*b*</sup> and Andrew J. Wilson<sup>\**a*</sup>

Received (in Cambridge, UK) 22nd March 2007, Accepted 24th April 2007 First published as an Advance Article on the web 8th May 2007 DOI: 10.1039/b704366j

### The regiospecific synthesis of $C_3$ macrocyclic scaffolds possessing multiple different functional groups is described.

The efficient synthesis of macrocyclic scaffolds such as cyclodextrins, calixarenes and porphyrins is the foundation on which modern supramolecular receptors are built. However, with the exception of cyclic peptides,<sup>1</sup> their syntheses are not amenable to construction of scaffolds possessing different peripheral functional groups in a defined sequence. This is a requirement for modern applications such as protein surface recognition.<sup>2</sup> Although statistical approaches<sup>3</sup> often prove useful, there is a need for synthetic methods that lead to scaffolds with these features. Cyclisation of *foldamers*<sup>4,5</sup>—of which aromatic oligoamides<sup>6–8</sup> are one class-represents a means of achieving this goal. Indeed, hydrogen-bonding has been used to promote cyclisation of the constituent building blocks of aromatic oligoamide foldamers9-12 and more recently to synthesise tetramers with a defined sequence of functional groups at the periphery.<sup>13</sup> Herein we report a different strategy that exploits the conformational bias of N-alkylated aromatic oligobenzamides<sup>14-17</sup> to achieve this goal via an iterative synthesis of linear trimers and subsequent cyclisation to yield macrocycles. Although cyclic trimers of this nature<sup>18,19</sup> can be made with identical substituents our approach leads to regiospecific incorporation of functional groups.

Our synthesis of the cyclic trimers 1a and 1b is shown in Scheme 1. The elegance of our approach derives from the fact that after each amide bond formation is carried out, the latent amine can be unmasked and then monoalkylated utilising reductive amination conditions. We first applied this synthesis to macrocycle 1a with three identical propyl side chains and then to an example in which three different substituents are incorporated- macrocycle 1b. Of a panel of different methods, we found in situ imine formation and reduction using borane-pyridine or borane-picoline complex to be ideal with unoptimised yields ranging from 79-95%. Other reducing agents such as sodium cyanoborohydride gave progressively poorer yields on chain elongation whilst isolation of the imine and reduction using sodium borohydride gave poor yields. As has previously been observed, we found that in situ formation of the acid chloride using dichlorotriphenylphosphorane<sup>20</sup> was a convenient method of amide bond formation. Many other peptide bond forming reagents including PyBOP, EDCI and

<sup>b</sup>CCLRC Daresbury Laboratory, Warrington, UK WA4 4AD

TFFH failed to give any reaction. Again unoptimised yields for this step ranged from 73–83%. It is also noteworthy that direct addition of 4-nitrobenzoyl chloride to a refluxing solution of the secondary aniline and triethylamine also afforded decent yields of product. Hydrolysis of the C-terminal esters 2a and 2b afforded free acids 3a and 3b. These both underwent ring closure using dichlorotriphenylphosphorane to afford the macrocycles 1a and 1b in yields of 89 and 93% yield respectively. In our hands, direct formation of macrocycle 1a from the *N*-propyl-4-aminobenzoic acid using dichlorotriphenylphosphorane resulted in poor yields and difficult purification.



Scheme 1 Synthesis of cyclic trimers 1a and 1b.

The cyclisation yields are remarkable, given that no special high dilution conditions or templates are used. The reaction is so efficient because the linear trimers adopt a folded conformation enforced by the known preference of N-alkylphenylbenzamides for the *cis* conformation.<sup>14–17</sup> This is confirmed by single crystal

<sup>&</sup>lt;sup>a</sup>School of Chemistry, University of Leeds, Woodhouse Lane, Leeds, UK LS2 9JT. E-mail: A.J.Wilson@leeds.ac.uk; Fax: +44(0)113-3436565; Tel: +44(0)113-3431409

<sup>†</sup> Electronic supplementary information (ESI) available: Synthetic procedures, additional NMR data and X-ray diffraction studies. See DOI: 10.1039/b704366j

studies of compound 1a and 2a and <sup>1</sup>H NMR and 2D NOESY experiments for 1b and 2b.



Fig. 1 X-Ray crystal structures of the acyclic protected trimer 2a and its corresponding cyclic trimer 1a (above CPK representation, below stick representation). Carbon is shown in grey, hydrogen in white, nitrogen in blue and oxygen in red.<sup>‡</sup>

Crystals suitable for X-ray diffraction were obtained by cooling a wet solution of methanol and ether of the propyl-appended linear trimer 2a.<sup>‡</sup> The structure (Fig. 1) clearly demonstrates the mutual proximity of the amine and carboxylate groups with both amides adopting a cis conformation. Crystals of macrocycle 1a were obtained by slow evaporation of a toluene-dichloromethane solution.† The asymmetric unit contains 4 molecules of trimer and 4 molecules of toluene which pack into columns of alternate trimer and toluene units held together by edge to face  $\pi$ - $\pi$  interactions (ESI Fig. S1<sup>†</sup>). Only one of the macrocycles 1a in the structure is shown in Fig. 1. Although slight differences are observed for each of the four trimers, the main structural features are the same. Thus, the structure is similar in nature to the previously disclosed N,N',N''-trimethyl derivative<sup>18</sup> and closely resembles the structure of the linear ester 2a from which it derives. For 1a all three amide groups necessarily adopt the cis conformation and the amides twist out of conjugation with the aromatic ring-clearly indicating why the structures can form so easily. From the crystal structure, the cavity volume is estimated to be  $\sim 15$  Å<sup>3</sup>. This is too small to be useful for host-guest chemistry and indicates the macrocycles will function exclusively as scaffolds onto which guest binding domains can be added.

The solid state conformational behaviour is maintained in solution. Fig. 2 shows the <sup>1</sup>H NMR spectra of compound **2b** and **1b**—NOE cross peaks (ESI Fig. S3) are highlighted for interacting protons.† The <sup>1</sup>H NMR spectrum of the acyclic trimer **2b** has only one set of resonances indicative of a strong bias towards one conformation. Observation of through-space correlation between protons situated on aromatic rings at opposing ends of the acyclic trimer indicate that the folded conformation is adopted. These



Fig. 2 <sup>1</sup>H NMR spectra (500 MHz, CDCl<sub>3</sub>) of acyclic trimer 2b and cyclic trimer 1b (arrows denote NOE cross peaks).

cross peaks are still present in the final macrocycle **1b** (not shown). In the <sup>1</sup>H NMR spectrum of **1b**, the amine resonance is lost whilst the aromatic resonances move closer together, although interestingly they remain well resolved despite only subtle differences in alkyl substitution. This is not a conformational issue as little change is observed upon heating a sample in  $C_2D_2Cl_4$  to 100 °C. Finally, the <sup>1</sup>H NMR and NOESY spectrum of compound **2a** is similarly complicated to that for **2b** but upon cyclisation simplifies to just five resonances consistent with a  $C_3$  symmetric cyclic compound (ESI Fig. S2 and S4).<sup>†</sup> This again suggests only one conformation is present in solution for these macrocycles.

In summary, we have described a robust method for regiospecific synthesis of differentially substituted  $C_3$  cyclic supramolecular scaffolds. Future studies will focus on incorporation of protected aldehyde fragments that contain useful functional groups for host–guest chemistry. We will report on these results in due course.

We would like to thank EPSRC (EP/D077842/1) for a financial support of this research and for an EPSRC DTA award (FC). We would also like to thank CCLRC for access to Station 16.2SMX, Daresbury Laboratory.

#### Notes and references

‡ CCDC 641457 and 641458. For crystallographic data in CIF or other electronic format see DOI: 10.1039/b704366j

- 1 D. T. Bong, T. D. Clark, J. R. Granja and M. R. Ghadiri, Angew. Chem., Int. Ed., 2001, 40, 988–1011.
- 2 A. J. Wilson, J. Hong and A. D. Hamilton, Org. Biomol. Chem., 2007, 5, 276–285.
- 3 L. Baldini, A. J. Wilson, J. Hong and A. D. Hamilton, J. Am. Chem. Soc., 2004, 126, 5656–5657.
- 4 S. H. Gellman, Acc. Chem. Res., 1998, 31, 178-180.
- 5 D. J. Hill, M. J. Mio, R. B. Prince, T. S. Hughes and J. S. Moore, *Chem. Rev.*, 2001, **101**, 3893–4012.
- 6 B. Gong, Chem.-Eur. J., 2001, 7, 4336-4342.
- 7 I. Huc, Eur. J. Org. Chem., 2004, 17-29.
- 8 Z.-T. Li, J.-L. Hou and H.-P. Yi, *Chem.–Asian J.*, 2006, 1, 766–778.

- 9 L. Y. Xing, U. Ziener, T. C. Sutherland and L. A. Cuccia, *Chem. Commun.*, 2005, 5751–5753.
- 10 H. Jiang, J. M. Leger, P. Guionneau and I. Huc, Org. Lett., 2004, 6, 2985–2988.
- 11 L. Yuan, W. Feng, K. Yamato, A. R. Sandford, D. Xu, H. Guo and B. Gong, J. Am. Chem. Soc., 2004, 126, 11120–11121.
- 12 L. He, Y. An, L. H. Yuan, W. Feng, M. F. Li, D. C. Zhang, K. Yamato, C. Zheng, X. C. Zeng and B. Gong, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 10850–10855.
- 13 S.-W. Kang, C. M. Gothard, S. Maitra, A.-T. Wahab and J. S. Nowick, J. Am. Chem. Soc., 2007, 129, 1486–1487.
- 14 K. Yamaguchi, G. Matsumura, H. Kagechika, I. Azumaya, Y. Ito, A. Itai and K. Shudo, J. Am. Chem. Soc., 1991, 113, 5474–5475.

- 15 I. Azumaya, H. Kagechika, K. Yamaguchi and K. Shudo, *Tetrahedron*, 1995, **51**, 5277–5290.
- 16 A. Tanatani, A. Yokoyama, I. Azumaya, Y. Takahura, C. Mitsui, M. Shiro, M. Uchiyama, A. Muranaka, N. Kobayashi and T. Yokozawa, J. Am. Chem. Soc., 2005, 127, 8553.
- H. M. Konig, R. Abbel, D. Schollmeyer and A. F. M. Kilbinger, *Org. Lett.*, 2006, 8, 1819–1822.
   I. A. K. M. Kilbinger, *Cont. Phys. Lett.*, 2001, 17
- I. Azumaya, T. Okamoto and H. Takayanagi, *Anal. Sci.*, 2001, **17**, 1363–1364.
   I. Azumaya, T. Okamoto, F. Imabeppu and H. Takayanagi,
- 17 I. AZUIIIAYA, I. OKAMOIO, F. IMADEPPU and H. Takayanagi, *Heterocycles*, 2003, **60**, 1419–1424.
  20 I. Azumaya, T. Okamoto, F. Imaharay, and H. Takayanagi, *Computer Science*, 2013, 100 (2014).
- 20 I. Azumaya, T. Okamoto, F. Imabeppu and H. Takayanai, *Tetrahedron*, 2003, **59**, 2325–2331.

# **Find a SOLUTION**

## ... with books from the RSC

### Choose from exciting textbooks, research level books or reference books in a wide range of subject areas, including:

- Biological science
- Food and nutrition
- Materials and nanoscience
- Analytical and environmental sciences
- Organic, inorganic and physical chemistry

### Look out for 3 new series coming soon ...

- RSC Nanoscience & Nanotechnology Series
- Issues in Toxicology
- RSC Biomolecular Sciences Series

RSC Advancing the Chemical Sciences



www.rsc.org/books