Tetrahedron Letters 50 (2009) 3719-3722

Contents lists available at ScienceDirect

Tetrahedron Letters

journal homepage: www.elsevier.com/locate/tetlet



Synthesis of polycyclic heterocycles via sequential Au-catalyzed cycloisomerization and Ru-catalyzed metathesis reactions

Chung-Meng Chao, Patrick Yves Toullec, Véronique Michelet*

Laboratoire de Synthèse Sélective Organique et Produits Naturels, ENSCP, UMR 7573, 11 rue P. et M. Curie, F-75231 Paris Cedex 05, France

ARTICLE INFO

Article history: Received 3 February 2009 Accepted 1 April 2009 Available online 5 April 2009

ABSTRACT

A sequential and efficient catalytic preparation of polycyclic heterocycles, including tetrahydro furo- or cyclopenta- pyran, hexahydropyrano pyrrolidine, and tetrahydroisochromene derivatives is described based on a gold-catalyzed cycloisomerization reaction followed by a Ru-catalyzed ring-closing metathesis. An isomerization of the alkene moiety leading diastereoselectively to cis tetrahydrocyclopenta[c]pyran and tetrahydro-1*H*-furo[3,4-c]pyran is observed in some cases. The catalytic hydrogenation reactions of these tetrahydropyrans led to the formation of polycyclic-functionalized saturated compounds.

© 2009 Elsevier Ltd. All rights reserved.

The transition metal-catalyzed reactions and specifically homogeneous catalysis have given a considerable impetus to the progress of the area of organometallic, organic, and industrial chemistries.¹ Among them, carbon-carbon and carbon-oxygen bond formations using gold² and ruthenium³ organometallic catalysts are fundamentally important because they provide successful solutions to environmental, economic concerns. Whereas gold catalysis was only recognized recently for the promotion of a myriad of transformations, ruthenium catalysis and more specifically ruthenium-catalyzed metathesis⁴ has been well established as an important tool for the organic chemist. As part of our ongoing program toward the development of catalytic cycloisomerization reactions,⁵ we have been interested in the synthesis of polycyclic derivatives and we have envisaged to test a sequential gold-catalyzed alkoxycyclization with a ruthenium-catalyzed ring-closing metathesis reaction. We and others have indeed described metalcatalyzed alkoxycyclization of enynes leading to cyclic ethers (Scheme 1, Eq 1).^{5,6} We anticipated that allylic derivatives might be engaged in Ru-catalyzed metathesis leading to bicyclic derivatives (Scheme 1, Eq 2). The resulting heterocycles would be valuable skeleton for the synthesis of natural and/or pharmaceutical product derivatives.⁷ Further hydrogenation reaction would then potentially allow the preparation of polycyclic heterocycles (Scheme 1, Eq. 3), controlling therefore three stereogenic centers by catalytic reactions. The sequential gold-catalyzed cycloisomerization/ring-closing metathesis has been scarcely described in the literature⁸ and only concerned the ring-closing metathesis reactions of a diene resulting from the addition of an allylsilane to 1,6-enynes^{8a} or a methoxycyclized bearing a geranyl and allyl side chains.^{8b} These strategies respond to environmental concerns and contribute to the concept of green chemistry.¹

As gold catalysts have been described for the alkoxycyclization step,^{5,6} we tested the allyloxycyclization reaction on various enynes. The catalyst combining Au(III)/PPh₃ and various silver salts was chosen for this study and the reaction was conducted in allylic alcohol which is used as a solvent at room temperature (Table 1). The addition of allylic alcohol was successfully realized on carboand heteroatom-linked 1,6-enynes (Table 1, entries 1–5). The allylic ethers **2a–e** were isolated in good to excellent yield. In the same manner as previously reported, one diastereomer was isolated for **2a**, **2b**, and **2e**. The lower reactivity of nitrogen-tethered enynes^{9,5c} prompted us to test the silver bis(trifluoromethanesulfonyl)imidate as an halide-abstracting agent (Table 1, entries 4 and 5). The corresponding ethers were thus prepared under mild



Scheme 1.

^{*} Corresponding author. Tel.: +33 1 44276742; fax: +33 1 44071062. *E-mail address:* veronique-michelet@enscp.fr (V. Michelet).

^{0040-4039/\$ -} see front matter @ 2009 Elsevier Ltd. All rights reserved. doi:10.1016/j.tetlet.2009.04.001

Table 1

Gold-catalyzed allyloxycyclization reaction of enynes 1c-j



Entry	Enyne 1		Conditions	<i>t</i> (h)	Product 2		Yield ^a (%)
1		1a	10 mol %AuCl ₃ , 10 mol %PPh ₃ , 30 mol %AgSbF ₆ 10 mol %AuCl ₃ , 10 mol %PPh ₃ , 30 mol %AgOTf	2 2.5		2a	79 66
2	MeO ₂ C Ph MeO ₂ C	1b	10 mol %AuCl ₃ , 10 mol %PPh ₃ , 30 mol %AgSbF ₆ 10 mol %AuCl ₃ , 10 mol %PPh3, 30 mol %AgOTf	1.5 2.5	MeO ₂ C MeO ₂ C	2b	90 89
3 ^b	MeO ₂ C	1c	10 mol %AuCl ₃ , 10 mol %PPh ₃ , 30 mol %AgNTf ₂	4.5	MeO ₂ C MeO ₂ C	2c	56
4	TsN	1d	10 mol %AuCl ₃ , 10 mol %PPh ₃ , 30 mol %AgNTf ₂	16	TsN	2d	70
5	TsN	1e	10 mol %AuCl ₃ , 10 mol %PPh ₃ , 30 mol %AgNTf ₂	2.5	TSN H	2e	53
6		1f	10 mol %AuCl ₃ , 10 mol %PPh ₃ , 30 mol %AgSbF ₆ 10 mol %AuCl ₃ , 10 mol %PPh ₃ , 30 mol %AgNTf ₂	72 72	SM SM		1
7	MeO ₂ C	1g	10 mol %AuCl ₃ , 10 mol %PPh ₃ , 30 mol %AgOTf	53	MeO ₂ C MeO ₂ C	2g	26 ^b
8	MeO ₂ C	1h	10 mol %mol %AuCl ₃ , 10 mol %PPh ₃ , 30 mol %AgOTf	14	MeO ₂ C MeO ₂ C	2h	42
9	MeO ₂ C	1i	10 mol %mol %AuCl ₃ , 10 mol %PPh ₃ , 30 mol %AgOTf	14	MeO ₂ C MeO ₂ C + MeO ₂ C MeO ₂ C 0 40:6		86
10		1j	10 mol %AuCl ₃ , 10 mol %PPh ₃ , 30 mol %AgSbF ₆	1	H(H Ph + 0 86 : 14	2j 2j	65

^a Isolated yield.

conditions at room temperature and isolated in 70% and 53% yields, respectively. No reactivity was observed for aryl-substituted enyne **1f** (Table 1, entry 6). The reaction of methyl-substituted prop-2-enyl substrates **1g**, **1h** (Table 1, entries 7 and 8) was more sluggish,

with several unidentified by-products being observed. The ether **2g** was isolated in 26% yield. In the case of crotyl-substituted derivative **1h**, as previously reported,^{5c} the reaction led selectively to the six-membered ether **2h**, which was obtained in 42% yield.

^b 60% conversion.



Scheme 2.

The allyloxycyclization reaction of 1,7-enyne **1i** afforded the desired ether in a modest 33% yield, the major product **3** resulting from the formal hydration of the alkyne moiety (Table 1, entry 9). We also prepared a chiral 1,6-enyne **1j** starting from commercially available (R)-but-1-yn-3-ol via a classical Williamson reaction. We were pleased to find that the tandem addition/ cyclization smoothly occurred and led to the formation of two diastereomers **2j** and **2j**' in 65% yield and in a 86/14 ratio. The stereochemistry of the major isomer was determined by NMR NOESY experiment.

Table 2

Ruthenium-catalyzed metathesis reaction of dienes ${\bf 2b-g}$





^a Isolated yield. Ar = $2,3-(CH_2OCH_2)-C_6H_3$



We also examined the possibility of introduction of propargylic alcohol, which would potentially allow an enyne metathesis process. The addition of propargylic alcohol to the enyne **1b** gave rise to the degradation of the starting material, either in the presence of allylic alcohol as the solvent or in a 6:1 mixture of dioxane/propargylic alcohol. The introduction of trimethylsilylpropargylic alcohol was successful and led to a mixture of **4** and **5** in 44% overall yield (Scheme 2). This mixture was directly engaged in a desilylation process under classical conditions, which afforded **4** in 43% yield. Other attempts to introduce trimethylsilylpropargylic alcohol on enynes **1a** or **1e** unfortunately did not give the desired products.

Our efforts next focused on the Ru-catalyzed metathesis reaction of the cyclopentanyldienes **2** in the presence of Grubbs II catalyst (2 or 4 mol %).⁴ The carbo- and heterocyclic ethers **2c** and **2d** reacted smoothly in the presence of 4 mol % of Grubbs II catalyst and the expected alkenes **6c** and **6d** were isolated in 64% yield (Table 2, entries 1 and 2). The reactivity of **2e** was particularly interesting as an isomerized derivative **7e** was isolated in 40% yield (Table 2, entry 3). The diastereoselectivity of this isomerization was further demonstrated after the hydrogenation reaction step on derivative **9**. Ru-catalyzed isomerization of olefins has been recently highlighted and was firstly described by Nishida, Cossy, and Alcaide groups in the presence of Grubbs II catalyst.¹⁰

The **6e**/**7e** ratio was showed to depend on the catalyst loading. The use of 2 mol % of catalyst led to a quasi equimolar quantity of alkenes **6e** and **7e** (Table 2, entry 3). The reactivity of enyne **1b** followed the same trend (Table 2, entries 4 and 5). The completely isomerized derivative **7b** was isolated in 99% yield in the presence of 4 mol %catalyst (Table 2, entry 4), whereas a 80:20 **6b**/**7b** mixture was obtained in the presence of 2 mol %of Grubbs II catalyst (Table 2, entry 5). The ring-closing metathesis of **2g** was also successful and afforded the tetrahydro-1*H*,3*H*-isochromene derivative **6g** in 80% yield (Table 2, entry 6). The enyne metathesis was attempted on substrate **5**, no conversion was observed after 102 h.

We then envisaged synthesizing completely saturated **5/6** bicyclic derivatives via the catalytic hydrogenation of the alkene moiety (Scheme 3). The classical hydrogenation of **7b** and **7e**, conducted in the presence of 5 mol %Pd/C in methanol at room temperature under 1 atm pressure H_2 afforded the corresponding bicyclic adducts **8** and **9** in 60% and 97% yields, respectively. The diastereoselectivity of the tandem gold-catalyzed reaction and the Ru-catalyzed ring-closing metathesis/isomerization process was fully demonstrated at this stage by NMR NOESY experiment on compound **9**.

In conclusion, we have developed a sequential catalytic preparation of polycyclic heterocycles based on a gold-catalyzed cycloisomerization reaction followed by a Ru-catalyzed ring-closing metathesis, leading efficiently to tetrahydro furo- or cyclopentapyran, hexahydropyrano pyrrolidine and tetrahydroisochromene skeletons. The second catalytic step was accompanied in some cases with an isomerization of the alkene moiety leading diastereoselectively to cis tetrahydrocyclopenta[c]pyran and tetrahydro-1*H*-furo[3,4-*c*]pyran. The catalytic hydrogenation reactions of these tetrahydropyrans led to the formation of polycyclic-functionalized saturated compounds.

Acknowledgments

This project was supported by the Centre National de Recherche Scientifique (CNRS) and the Ministère de l'Education et de la Recherche. C.-M.C. thanks the Ministère de l'Education et de la Recherche for a grant (2007–2009). The authors thank Dr. M. N. Rager for NOESY NMR analysis of compounds **9**, **2***j*, and **2***j*'.

Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.tetlet.2009.04.001.

References and notes

- (a) Cornils, B.; Herrmann, W. A. Applied Homogeneous Catalysis with Organometallic Compound; VCH: New-York, 1996; (b) Trost, B. M. Acc. Chem. Res. 2002, 35, 695; (c) Trost, B. M. Angew.Chem., Int. Ed. Engl. 1995, 34, 259; (d) Trost, B. M. Science 1991, 254, 1471; (e) Anastas, P.; Warner, J. C. Green Chemistry, Theory and Practice; Oxford University Press, 1998; (f) Anastas, P. T.; Kirchhoff, M. M. Acc. Chem. Res. 2002, 35, 686; (g) Anastas, P. T.; Zimmerman, J. B. Environ. Sci. Technol. 2003, 37, 94A; (h) Poliakoff, M.; Fitzpatrick, J. M.; Farren, T. R.; Anastas, P. T. Science 2002, 297, 807; (i) Trost, B. M.; Toste, D. F.; Pinkerton, A. B. Chem. Rev. 2001, 101, 2067.
- (a) Arcadi, A. Chem. Rev. 2008, 108, 3266; (b) Patil, N. T.; Yamamoto, Y. Chem. Rev. 2008, 108, 3395; (c) Li, Z.; Brouwer, C.; He, C. Chem. Rev. 2008, 108, 3239; (d) Shen, H. C. Tetrahedron 2008, 64, 3885; (e) Skouta, R.; Li, C.-J. Tetrahedron 2008, 64, 4917; (f) Hashmi, A. S. K.; Rudolph, M. Chem. Soc. Rev. 2008, 37, 1766; (g) Muzart, J. Tetrahedron 2008, 64, 5815; (h) Hashmi, A. S. K. Chem. Rev. 2007, 107, 3180; (i) Gorin, D. J.; Toste, D. Nature 2007, 446, 395; (j) Hashmi, A. S. K.; Hutchings, G. J. Angew Chem., Int. Ed. 2006, 45, 7896; (k) Jimenez-Nunez, E.; Echavarren, A. M. Chem. Commun. 2007, 333; (l) Ma, S.; Yu, S.; Gu, Z. Angew. Chem., Int. Ed. 2006, 45, 200; (m) Hashmi, A. S. K. Angew. Chem., Int. Ed. 2005, 44, 6090; (n) Höffmann-Röder, A.; Krause, N. Org. Biomol. Chem. 2005, 3, 387.
- (a)For selected reviews on Ru-catalyzed reactions: Hand Book of Metathesis; Grubbs, R. H., Ed.; Wiley-VCH: Weinheim, 2003; Vols. 1–3, (b) Kotha, S.; Mandal, K. Chem. Asian J. 2009. 10.1002/asia.200800244; (c) Grubbs, R. H. Tetrahedron 2004, 60, 7117; (d) McReynolds, M. D.; Dougherty, J. M.; Hanson, P. R. Chem. Rev. 2004, 104, 2239; (e) Nakamura, I.; Yamamoto, Y. Chem. Rev. 2004, 104, 2127–2198; (f) Deiters, A.; Martin, S. F. Chem. Rev. 2004, 104, 2199; (g) Diver, S. T.; Giessert, A. J. Chem. Rev. 2004, 104, 1317; (h) Villar, H.; Frings, M.; Bolm, C. Chem. Soc. Rev. 2007, 55; (i) Sémeril, D.; Bruneau, C.; Dixneuf, P. H. Adv. Synth. Catal. 2002, 344, 585; (j) Nicolaou, K. C.; Bulger, P. G.; Sarlah, D. Angew. Chem., Int. Ed. 2005, 44, 4490; (k) Kotha, S.; Lahiri, K. Synlett 2007, 2767; (l) Trost, B. M.; Frederiksen, M. U.; Rudd, M. T. Angew. Chem., Int. Ed. 2005, 44, 6630; (m) Naota, T.; Takaya, H.; Murahashi, S.-I. Chem. Rev. 1998, 98, 2599.

- (a) Schwab, P.; Grubbs, R. H.; Ziller, J. W. J. Am. Chem. Soc. 1996, 118, 100;
 (b) Scholl, M.; Ding, S.; Lee, C. W.; Grubbs, R. H. Org. Lett. 1999, 1, 953;
 (c) Garber, S. B.; Kingsbury, J. S.; Gray, B. L.; Hoveyda, A. H. J. Am. Chem. Soc. 2000, 122, 8168.
- (a) Nevado, C.; Charruault, L.; Michelet, V.; Nieto-Oberhuber, C.; Muñoz, M. P.; Méndez, M.; Rager, M.-N.; Genêt, J.-P.; Echavarren, A. M. Eur. J. Org. Chem. 2003, 706; (b) Charruault, L.; Michelet, V.; Taras, R.; Gladiali, S.; Genêt, J.-P. Chem. Comm. 2004, 850; (c) Genin, E.; Leseurre, L.; Toullec, P. Y.; Genêt, J.-P.; Michelet, V. Synlett 2007, 1780; (d) Leseurre, L.; Toullec, P. Y.; Genêt, J.-P.; Michelet, V. Org. Lett. 2007, 9, 4049; (e) Genin, E.; Toullec, P. Y.; Antoniotti, S.; Brancour, C.; Genêt, J.-P.; Michelet, V. J. Am. Chem. Soc. 2006, 128, 3112; (f) Genin, E.; Antoniotti, S.; Michelet, V.; Genêt, J.-P. Angew. Chem., Int. Ed. 2005, 44, 4949; (g) Antoniotti, S.; Genin, E.; Michelet, V.; Genêt, J.-P. J. Am. Chem. Soc. 2005, 127, 9976; (h) Genin, E.; Toullec, P. Y.; Marie, P.; Antoniotti, S.; Brancour, C.; Genêt, J.-P.; Michelet, V. Arkivoc 2007, v, 67; (i) Neatu, F.; Li, Z.; Richards, R.; Toullec, P. Y.; Genêt, J.-P.; Dumbuya, K.; Gottfried, J. M.; Steinrück, H.-P.; Pârvulescu, V. I.; Michelet, V. Chem. Eur. J. 2008, 14, 9412; (j) Chao, C.-M.; Genin, E.; Toullec, P. Y.; Genêt, J.-P.; Michelet, V. J. Organomet. Chem. 2008, 694, 538; (k) Toullec, P. Y.; Chao, C.-M.; Chen, Q.; Genêt, J.-P.; Michelet, V. Adv. Synth. Catal. 2008, 250, 2401; (1) Toullec, P. Y.; Genin, E.; Leseurre, L.; Genêt, J.-P.; Michelet, V. Angew. Chem., Int. Ed. 2006, 45, 7427; (m) Chao, C.-M.; Vitale, M. R.; Toullec, P. Y.; Genêt, J.-P.; Michelet, V. Chem. Eur. J. 2009, 15, 1319; (n) Leseurre, L.; Chao, C. M.; Seki, T.; Genin, E.; Toullec, P. Y.; Genêt, J.-P.; Michelet, V. Tetrahedron 2009, doi:10.1016/j.tet.2008.11.105.
- (a) Nevado, C.; Cárdenas, D. J.; Echavarren, A. M. Chem. Eur. J. 2003, 9, 2627; (b) Martín-Matute, B.; Nevado, C.; Cárdenas, D. J.; Echavarren, A. M. J. Am. Chem. Soc. 2003, 125, 5757; (c) Méndez, M.; Muñoz, M. P.; Nevado, C.; Cárdenas, D. J.; Echavarren, A. M. J. Am. Chem. Soc. 2001, 123, 10511; (d) Méndez, M.; Muñoz, M. P.; Echavarren, A. M. J. Am. Chem. Soc. 2000, 122, 11549; (e) Nishizawa, M.; Yadav, V. K.; Skwarczynski, M.; Takao, H.; Imagawa, H.; Sugihara, T. Org. Lett. 2003, 5, 1609; (f) Nishizawa, M.; Imagawa, H. J. Synth. Org. Chem. Japan 2006, 64, 744; Faller, J. W.; Fontaine, P. P. J. Organomet. Chem. 2006, 691, 1912; (g) Muñoz, M. P.; Adrio, J.; Carretero, J. C.; Echavarren, A. M. Organometallics 2005, 24, 1293; (h) Nieto-Oberhuber, C.; Lopez, S.; Echavarren, A. M. J. Am. Chem. Soc. 2005, 127, 6178; (i) Nieto-Oberhuber, C.; Muñoz, M. P.; Nevado, C.; Cárdenas, D. J.; Echavarren, A. M. Angew. Chem., Int. Ed. 2004, 43, 2402; (j) Mézailles, N.; Ricard, L.; Gagosz, F. Org. Lett. 2005, 7, 4133; (k) Nieto-Oberhuber, C.; Muñoz, M. P.; Lopez, S.; Jimenez-Nuñez, E.; Nevado, C.; Herrero-Gomez, E.; Raducan, M.; Echavarren, A. M. Chem. Eur. J. 2006, 12, 1677; (1) Buzas, A. K.; Istrate, F. M.; Gagosz, F. Angew. Chem., Int. Ed. **2007**, 46, 1141; (m) Zhang, L.; Kozmin, S. A. J. Am. Chem. Soc. 2005, 127, 6962-6963; (n) Cabello, N.; Rodriguez, C.; Echavarren, A. M. Synlett 2007, 1753.
- (a) Durst, G. L.; Norman, B. H.; Pfeifer, L. A.; Richardson, T. I. PCT Int. Appl. 2004, WO 2004094400.; (b) Norman, B. H.; Richardson, T. I. PCT Int. Appl. 2006, WO 2006044176.; (c) Ratnayake, R.; Covell, D.; Ransom, T. T.; Gustafson, K. R.; Beutler, J. A. Org. Lett. 2008, 10, 57; (d) Jimenez-Alonso, S.; Perez-Lomas, A.; Estevez-Braun, A.; Munoz Martinez, F.; Chavez Orellana, H.; Ravelo, A. G.; Gamarro, F.; Castanys, S.; Lopez, M. J. Med. Chem. 2008, 51, 7132; (e) Wang, H.-X.; Liu, C.-M.; Liu, Q.; Gao, K. Phytochemistry 2008, 69, 2088; (f) Korthals, K. A.; Wulff, W. D. J. Am. Chem. Soc. 2008, 130, 2898; (g) Richardson, T. I.; Dodge, J. A.; Wang, Y.; Durbin, J. D.; Krishnan, V.; Norman, B. H. Bioorg. Med. Chem. Lett. 2007, 17, 5563. and references cited therein; (h) Barlow, A. J.; Compton, B. J.; Hertewich, U.; Lorimer, S. D.; Weavers, R. T. J. Nat. Prod. 2005, 68, 825; (i) Zheng, Y.; Shen, Y. Org. Lett. 2009, 11, 109.
- (a) Amijs, C. H. M.; Lopez-Carrillo, V.; Raducan, M.; Perez-Galan, P.; Ferrer, C.; Echavarren, A. M. J. Org. Chem. 2008, 73, 7721; (b) 61
- 9. Dr. E. Genin, Dissertation from the University Pierre et Marie Curie, 2006.
- For leading references, see: (a) Donohoe, T. J.; O'Riordan, T. J. C.; Rosa, C. P. Angew. Chem. Int. Ed. 2009, 48, 1014; (b) Alcaide, B.; Almendros, P.; Alonso, J. M. Org. Lett. 2001, 3, 3781; (c) Arisawa, M.; Terada, Y.; Nakagawa, M.; Nishida, A. Angew. Chem., Int. Ed. 2002, 41, 4732; (d) Cadot, C.; Dalko, P. I.; Cossy, J. Tetrahedron Lett. 2002, 43, 1839.