

This article was downloaded by: [University of Cambridge]

On: 20 October 2014, At: 01:19

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

Rhodium-Catalyzed Anti-Markovnikov-Type Hydrophosphination of Terminal Alkynes with Diphosphines and Hydrosilanes in the Presence of Oxygen

Shin-ichi Kawaguchi^a, Mao Kotani^a, Takashi Ohe^a, Shoko Nagata^a, Akihiro Nomoto^a, Motohiro Sonoda^a & Akiya Ogawa^a

^a Department of Applied Chemistry, Graduate School of Engineering, Osaka Prefecture University, Osaka, Japan
Published online: 27 May 2010.

To cite this article: Shin-ichi Kawaguchi, Mao Kotani, Takashi Ohe, Shoko Nagata, Akihiro Nomoto, Motohiro Sonoda & Akiya Ogawa (2010) Rhodium-Catalyzed Anti-Markovnikov-Type Hydrophosphination of Terminal Alkynes with Diphosphines and Hydrosilanes in the Presence of Oxygen, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 185:5-6, 1090-1097, DOI: [10.1080/10426501003773423](https://doi.org/10.1080/10426501003773423)

To link to this article: <http://dx.doi.org/10.1080/10426501003773423>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

RHODIUM-CATALYZED ANTI-MARKOVNIKOV-TYPE HYDROPHOSPHINATION OF TERMINAL ALKYNES WITH DIPHOSPHINES AND HYDROSILANES IN THE PRESENCE OF OXYGEN

Shin-ichi Kawaguchi, Mao Kotani, Takashi Ohe, Shoko Nagata, Akihiro Nomoto, Motohiro Sonoda, and Akiya Ogawa

Department of Applied Chemistry, Graduate School of Engineering, Osaka Prefecture University, Osaka, Japan

A novel rhodium(I)-catalyzed hydrophosphination of terminal alkynes with diphosphines and hydrosilanes takes place regioselectively in the presence of small amounts of oxygen. After air-oxidation during workups, the corresponding anti-Markovnikov-type vinylic phosphine oxides were obtained in good yields.

Keywords Alkyne; anti-Markovnikov; diphosphine; hydrophosphination; hydrosilane; rhodium catalyst

INTRODUCTION

Numerous studies of the additions of heteroatom compounds bearing a heteroatom–heteroatom linkage such as B–B,¹ Si–Si,² Ge–Ge,³ Sn–Sn,⁴ S–S,⁵ and Se–Se⁵ to alkynes catalyzed by transition metal catalysts have been reported.⁶ These addition reactions provide useful tools to vicinally bifunctionalized vinylic compounds. Recently, we reported the addition of tetraphenyldiphosphine as heteroatom compounds bearing a P–P linkage, to terminal alkynes in the presence of palladium catalyst.⁷ Surprisingly, this reaction afforded not the corresponding bisphosphination⁸ product but a Markovnikov-type hydrophosphination^{9–12} product (**3**), regioselectively, in the co-presence of oxygen (Scheme 1).


Based on the result of the hydrophosphination using deuterized alkyne, it has been apparent that the substrate alkynes also act as a hydrogen source in this hydrophosphination.⁷ Furthermore, we found that this hydrophosphination of alkynes proceeded efficiently by the addition of hydrosilane as a hydrogen source in the presence of small amounts of

Received 3 December 2008; accepted 14 January 2009.

Dedicated to Professor Naomichi Furukawa on the occasion of his 70th birthday.


This research was supported by a Grant-in-Aid for Scientific Research on Priority Areas (Area 444, No. 19020061) and Scientific Research (B, 19350095), from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

Address correspondence to Akiya Ogawa, Department of Applied Chemistry, Graduate School of Engineering, Osaka Prefecture University, 1-1 Gakuen-cho, Nakaku, Sakai, Osaka 599-8531, Japan. E-mail: ogawa@chem.osakafu-u.ac.jp




Scheme 1

oxygen (Scheme 2).¹³ Oxygen seems to oxidize appropriate amounts of tetraphenyldiphosphine to tetraphenyldiphosphine oxide, which inhibits the deactivation of the catalyst in this hydrophosphination reaction.



Scheme 2

By switching the catalyst simply from palladium(0) to rhodium(I), *anti*-Markovnikov-type hydrophosphination of terminal alkynes has been found to take place regioselectively (Scheme 3). In this article, we report a novel rhodium-catalyzed *anti*-Markovnikov-type hydrophosphination of terminal alkynes using a diphosphine–hydrosilane binary system in the presence of oxygen.




Scheme 3

RESULTS AND DISCUSSION

In a sealed NMR tube under N₂ atmosphere, 1-octyne (**1a**) (0.3 mmol), tetraphenyldiphosphine (**2**) (0.1 mmol), diethylmethylsilane (**5**) (0.3 mmol), tris(triphenylphosphine)rhodium(I) chloride (0.005 mmol), and C₆D₆ (600 μl) were placed, and then a quarter equivalent of oxygen (0.025 mmol) was introduced into the tube. Heating the resulting mixture at 80 °C for 18 h, and the subsequent air-oxidation during workups, afforded the corresponding *anti*-Markovnikov-type hydrophosphination product, (*E*)-1-diphenylphosphinyl-1-octene (**7a**), in 90% yield (Scheme 3).

In addition, the formation of silylphosphine oxide as a byproduct was observed by ³¹P NMR (Et₂MeSiP(O)Ph₂: ³¹P NMR (C₆D₆) δ 21.8 ppm).¹³

Since tetraphenyldiphosphine (**2**) may be oxidized by oxygen to the corresponding oxide **8**, we next examined this hydrophosphination of 1-octyne (**1a**) by adding tetraphenyldiphosphine oxide (**8**) in the absence of oxygen. When a mixture of 1-octyne (**1a**) (0.6 mmol), tetraphenyldiphosphine (**2**) (0.1 mmol), tetraphenyldiphosphine oxide (**8**) (0.1 mmol), diethylmethylsilane (**5**) (0.6 mmol), and C₆D₆ (600 μl) in the presence of tris(triphenylphosphine)rhodium(I) chloride (0.025 mmol) was heated under the same conditions, the corresponding vinylphosphine as the hydrophosphination product and its oxide **7a** were formed. After air-oxidation during workups, vinylphosphine oxide **7a** was obtained in 83% yield (Scheme 4). (The yield was based on the sum of the mole numbers of tetraphenyldiphosphine and tetraphenyldiphosphine oxide.)



Scheme 4

We next examined the hydrophosphination of several alkynes by using the methods shown in Schemes 3 and 4 (Table I). Aromatic alkynes such as phenylacetylene underwent the Rh(I)-catalyzed regioselective hydrophosphination, providing the corresponding (*E*)-vinylic phosphine oxide (**7b**) in good yield (Table I, entry 3). Chloro and silyl substituents are tolerant of the reaction (Table I, entries 4, 5, and 7). In the case of a conjugate enyne, the corresponding phosphinyl diene (**7d**) was obtained in good yield (Table I, entry 6).

Although additional detailed experiments are required in order to clarify the reaction mechanism of this rhodium(I)-catalyzed hydrophosphination, possible pathways¹³ are shown in Scheme 5: (i) at first, oxidative addition of diphosphine oxide to rhodium(I) to generate active species (**9**); (ii) regioselective phosphinorhodation of alkyne, to form vinylrhodium intermediate (**10**); two possible pathways (path A and B) are considered: path A: (iii) the reaction of vinylrhodium **10** with the hydrosilane to form vinylphosphine (**6**) and

Table 1 Rhodium(I)-catalyzed hydrophosphination of alkynes using diphosphine-hydrosilane binary systems

Entry	Alkyne	Condition	Product	Yield% ^b
1	n Hex-≡	A	n Hex-CH=CH-P(=O)(Ph) ₂ 7a	90%
2		B		83%
3		A		82% ^d
4		A		80%
5		B		63%
6		A		90% ^d
7 ^c	TMS-≡	A	TMS-CH=CH-P(=O)(Ph) ₂ 7e	71%

^aReaction condition A: alkyne (0.3 mmol), (Ph₂P)₂ (0.1 mmol), Et₂MeSiH (0.3 mmol), oxygen (0.025 mmol), RhCl(PPh₃)₃ (0.005 mmol). Reaction condition B: alkyne (0.6 mmol), (Ph₂P)₂ (0.1 mmol), Ph₂PP(O)Ph₂ (0.1 mmol), Et₂MeSiH (0.6 mmol), RhCl(PPh₃)₃ (0.01 mmol).

^bDetermined by ¹H NMR.


^cRhCl(PPh₃)₃ (10 mol%, 0.02 mmol) was used.

^dThe yield of this Rh-catalyzed reaction was higher compared with that of the Pd(PPh₃)₄-catalyzed hydrophosphination,¹³ in which some minor byproducts were formed concomitantly.

silylphosphinyrhodium species (**11**) and (iv) exchange reaction of the silyl group of **11** with the phosphino group of tetraphenyldiphosphine to regenerate Rh species **9** with the concomitant formation of silylphosphine as a byproduct; path B: (iii) exchange of phosphinyl group with hydride group to form vinylrhodium hydride species (**12**) and silylphosphine oxide and (iv) reductive elimination to give vinylphosphine **6** and addition of diphosphine oxide to regenerate **9**.

An alternative pathway may involve the formation of phosphinyrhodium hydride (**13**) (Scheme 6): (i) the reaction of rhodium species **9** with the hydrosilane to generate phosphinyrhodium hydride **13**¹⁴; (ii) hydrorhodation of alkyne to generate vinylrhodium hydride (**14**); (iii) the reaction of **14** with the diphosphine to give vinylphosphine **6** along with the regeneration of **9**.

In summary, rhodium-catalyzed highly regioselective hydrophosphination of terminal alkynes has been developed by using diphosphine-hydrosilane binary systems, which affords the *anti*-Markovnikov-type addition products. This reaction is regiocomplementary with the palladium-catalyzed Markovnikov-type hydrophosphination reported previously.



Scheme 5


EXPERIMENTAL

General Procedure for Rhodium-Catalyzed Hydrophosphination of Alkynes Using Diphosphine-Hydrosilane Binary Systems

Condition A. $(\text{Ph}_2\text{P})_2$ (37.0 mg, 0.1 mmol), $\text{Ph}_2\text{P(=O)Ph}_2$ (38.6 mg, 0.1 mmol), $\text{RhCl(PPh}_3)_3$ (9.3 mg, 0.01 mmol), alkyne (0.6 mmol), Et_2MeSiH (87 μl , 0.6 mmol), and C_6D_6 (0.6 mL) were placed in a sealed Pyrex glass NMR tube under a nitrogen atmosphere. The mixture was stirred for 30 sec, and the tube was covered with aluminum foil. The mixture was heated under reflux for 18 h, and then was left under air overnight. Purification of the crude materials was performed by preparative TLC.

Condition B. $(\text{Ph}_2\text{P})_2$ (37.0 mg, 0.1 mmol), $\text{RhCl(PPh}_3)_3$ (4.6 mg, 0.005 mmol), alkyne (0.3 mmol), Et_2MeSiH (43 μl , 0.3 mmol), and C_6D_6 (0.6 mL) were placed in a sealed Pyrex glass NMR tube. Next, oxygen (0.6 mL, 0.025 mmol) was introduced into the tube. The mixture was stirred for 30 sec, and the tube was covered with aluminum foil. The mixture was heated under reflux for 18 h, and was left under air overnight. Purification of the crude products was performed by preparative TLC.

(E)-1-(Diphenylphosphinyl)-1-octene (7a). Colorless oil; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.18–7.72 (m, 10H), 6.73 (ddt, $J = 6.5, 16.9, J_{\text{HP}} = 19.4$ Hz, 1H), 6.22 (dd, $J = 17.0, J_{\text{HP}} = 24.7$ Hz, 1H), 2.30 (q, 2H), 1.28–1.50 (m, 8H), 0.86 (t, 3H); $^{13}\text{C NMR}$ (100



Scheme 6

MHz, CDCl_3) δ 153.0, 133.2 ($J_{\text{CP}} = 104.5$ Hz), 131.6 ($J_{\text{CP}} = 2.5$ Hz), 131.3 ($J_{\text{CP}} = 9.9$ Hz), 128.5 ($J_{\text{CP}} = 12.3$ Hz), 121.5 ($J_{\text{CP}} = 102.8$ Hz), 34.5 ($J_{\text{CP}} = 16.5$ Hz), 31.5, 28.8, 27.8, 22.5, 14.0; ^{31}P NMR (200 MHz, CDCl_3) δ 24.2.

REFERENCES

- (a) T. Ishiyama, N. Matsuda, N. Miyaura, and A. Suzuki, *J. Am. Chem. Soc.*, **115**, 11018 (1993); (b) T. Ishiyama, N. Matsuda, M. Murata, F. Ozawa, A. Suzuki, and N. Miyaura, *Organometallics*, **15**, 713 (1996).
- (a) H. Sakurai, Y. Kamiyama, and Y. Nakadaira, *J. Am. Chem. Soc.*, **97**, 931 (1975); (b) H. Sakurai, Y. Kamiyama, and Y. Nakadaira, *Chem. Lett.*, **4**, 887 (1975); (c) H. Okinoshima, K. Yamamoto, and M. Kumada, *J. Organomet. Chem.*, **86**, C27 (1975); (d) H. Watanabe, M. Kobayashi, K. Higuchi, and Y. Nagai, *J. Organomet. Chem.*, **186**, 51 (1980); (e) H. Yamashita, N. P. Reddy, and M. Tanaka, *Organometallics*, **16**, 5223 (1997).
- (a) E. Piers and R. T. Skerlj, *J. Chem. Soc., Chem. Commun.*, 1025 (1987); (b) T. Tsumuraya and W. Ando, *Organometallics*, **8**, 2286 (1989); (c) T. Hayashi, H. Yamashita, T. Sakakura, Y. Uchamaru, and M. Tanaka, *Chem. Lett.*, **20**, 245 (1991).
- (a) T. N. Mitchell, A. Amamria, H. Killing, and D. Rutschow, *J. Organomet. Chem.*, **241**, C45 (1983); (b) E. Piers and R. T. Skerlj, *Can. J. Chem.*, **72**, 2468 (1994).
- (a) For the metal-catalyzed addition of E–E (E = group 16) to carbon–carbon unsaturated bonds, see: (a) H. Kuniyasu, A. Ogawa, S. Miyazaki, I. Ryu, N. Kambe, and N. Sonoda, *J. Am. Chem. Soc.*, **113**, 9796 (1991); (b) T. Kondo, S. Uenoyama, K. Fujita, and T. Mitsudo, *J. Am. Chem. Soc.*, **121**, 482 (1999); (c) M. Arisawa and M. Yamaguchi, *Org. Lett.*, **3**, 763 (2001); (d) H. Kuniyasu, In *Catalytic Heterofunctionalization*, A. Togni and H. Grützmacher, Eds. (Wiley-VCH,

- Weinheim, Germany, 2001) Chapter 7; (e) A. Ogawa, In *Handbook of Organopalladium Chemistry for Organic Synthesis*, E. Negishi, Ed. (Wiley-Interscience, New York, 2002), Chapter VII; (f) V. P. Ananikov, M. A. Kabeshov, I. P. Beletskaya, G. G. Aleksandrov, and I. L. Eremenko, *J. Organomet. Chem.*, **687**, 451 (2003); (g) M. Arisawa, Y. Kozuki, and M. Yamaguchi, *J. Org. Chem.*, **68**, 8964 (2003); (h) S. Usugi, H. Yorimitsu, H. Shinokubo, and K. Oshima, *Org. Lett.*, **6**, 601 (2004); (i) V. P. Ananikov and I. P. Beletskaya, *Org. Biomol. Chem.*, **2**, 284 (2004); (j) A. Ogawa, In *Main Group Metals in Organic Synthesis*, H. Yamamoto and K. Oshima, Eds. (Wiley-VCH, Weinheim, Germany, 2004), vol. 2, Chapter 15; (k) A. V. Moro, C. W. Nogueira, N. B. V. Barbosa, P. H. Menezes, J. B. T. daRocha, and G. Zeni, *J. Org. Chem.*, **70**, 5257 (2005); (l) V. P. Ananikov, M. A. Kabeshov, and I. P. Beletskaya, *Synlett*, **6**, 1012 (2005).
- (a) I. P. Beletskaya and C. Moberg, *Chem. Rev.*, **106**, 2320 (2006); (b) K. A. Horn, *Chem. Rev.*, **95**, 1317 (1995); (c) H. K. Sharma and K. H. Pannell, *Chem. Rev.*, **95**, 1351 (1995); (d) I. Beletskaya and C. Moberg, *Chem. Rev.*, **99**, 3435 (1999); (e) L.-B. Han and M. Tanaka, *Chem. Commun.*, 395 (1999); (f) M. Suginome and Y. Ito, *Chem. Rev.*, **100**, 3221 (2000); (g) A. Ogawa, *J. Organomet. Chem.*, **611**, 463 (2000).
 - S. Nagata, S.-i. Kawaguchi, M. Matsumoto, I. Kamiya, A. Nomoto, M. Sonoda, and A. Ogawa, *Tetrahedron Lett.*, **48**, 6637 (2007).
 - (a) V. A. Tzschach and S. Baensch, *J. Prakt. Chem.*, **313**, 254 (1971); (b) A. Sato, H. Yorimitsu, and K. Oshima, *Angew. Chem. Int. Ed.*, **44**, 1694 (2005); (c) S.-i. Kawaguchi, S. Nagata, K. Tsuchii, A. Nomoto, and A. Ogawa, *Tetrahedron Lett.*, **47**, 3919 (2006); (d) D. L. Dodds, M. F. Haddow, A. G. Orpen, P. G. Pringle, and G. Woodward, *Organometallics*, **25**, 5937 (2006); (e) A. Kondoh, H. Yorimitsu, and K. Oshima, *J. Am. Chem. Soc.*, **129**, 4099 (2007).
 - For reviews concerning transition-metal-catalyzed hydrophosphination and hydrophosphinylation to carbon-carbon unsaturated bonds, see: D. K. Wicht and D. S. Glueck, In *Catalytic Heterofunctionalization*; A. Togni and H. Grützmaier, Eds. (Wiley-VCH, Weinheim, Germany, 2001), Chapter 5.
 - For the metal-catalyzed hydrophosphination to carbon-carbon unsaturated bonds, see: (a) P. G. Pringle, and M. B. Smith, *J. Chem. Soc., Chem. Commun.*, 1701 (1990); (b) M. A. Kazankova, I. V. Efimova, A. N. Kochetkov, V. V. Afanas'ev, I. P. Beletskaya, and P. H. Dixneuf, *Synlett*, **4**, 497 (2001); (c) M. O. Shulyupin, M. A. Kazankova, and I. P. Beletskaya, *Org. Lett.*, **4**, 761 (2002); (d) K. Takaki, G. Koshiji, K. Komeyama, M. Takeda, T. Shishido, A. Kitani, and K. Takehira, *J. Org. Chem.*, **68**, 6554 (2003); (e) K. Komeyama, T. Kawabata, K. Takehira, and K. Takaki, *J. Org. Chem.*, **70**, 7260 (2005); (f) H. Ohmiya, H. Yorimitsu, and K. Oshima, *Angew. Chem. Int. Ed.*, **44**, 2368 (2005); (g) C. Scriban, I. Kovacic, and D. S. Glueck, *Organometallics*, **24**, 4871 (2005); (h) K. Takaki, K. Komeyama, D. Kobayashi, T. Kawabata, and K. Takehira, *J. Alloys Compounds*, **408**, 432 (2006); (i) M. Hayashi, Y. Matsuura, and Y. Watanabe, *J. Org. Chem.*, **71**, 9248 (2006); (j) K. Komeyama, D. Kobayashi, Y. Yamamoto, K. Takehira, and K. Takaki, *Tetrahedron*, **62**, 2511 (2006); (k) B. Join, D. Mimeau, O. Delacroix, and A.-C. Gaumont, *Chem. Commun.*, 3249 (2006); (l) M. R. Crimmin, A. G. M. Barrett, M. S. Hill, P. B. Hitchcock, and P. A. Procopiou, *Organometallics*, **26**, 2953 (2007).
 - For the transition-metal-catalyzed hydrophosphinylation to carbon-carbon unsaturated bonds, see: (a) L.-B. Han and M. Tanaka, *J. Am. Chem. Soc.*, **118**, 1571 (1996); (b) L.-B. Han, N. Choi, and M. Tanaka, *Organometallics*, **15**, 3259 (1996); (c) L.-B. Han, R. Hua, and M. Tanaka, *Angew. Chem. Int. Ed.*, **37**, 4196 (1998); (d) C.-Q. Zhao, L.-B. Han, and M. Tanaka, *Organometallics*, **19**, 4196 (2000); (e) L.-B. Han, C.-Q. Zhao, and M. Tanaka, *J. Org. Chem.*, **66**, 5929 (2001); (f) S. Deprère and J.-L. Montchamp, *J. Am. Chem. Soc.*, **124**, 9386 (2002); (g) A. Allen Jr., L. Ma, and W. Lin, *Tetrahedron Lett.*, **43**, 3707 (2002); (h) M. D. Milton, G. Onodera, Y. Nishibayashi, and S. Uemura, *Org. Lett.*, **6**, 3993 (2004); (i) L.-B. Han, C. Zhang, H. Yazawa, and S. Shimada, *J. Am. Chem. Soc.*, **126**, 5080 (2004); (j) P. Ribière, K. Bravo-Altamirano, M. I. Antczak, J. D. Hawkins, and J.-L. Montchamp, *J. Org. Chem.*, **70**, 4064 (2005); (k) R. A. Stockland Jr., A. J. Lipman, J. A. Bawiec III, P. E. Morrison, I. A. Guzei, P. M. Findeis, and J. F. Tamblin,

- J. Organomet. Chem.*, **691**, 4042 (2006); (l) N. Dobashi, K. Fuse, T. Hoshino, J. Kanada, T. Kashiwabara, C. Kobata, S. K. Nune, and M. Tanaka, *Tetrahedron Lett.*, **48**, 4669 (2007).
12. For the rhodium-catalyzed synthesis of 1-alkynylphosphine oxides from 1-alkynes and tetraphenyldiphosphine, see: M. Arisawa, M. Onoda, C. Hori, and M. Yamaguchi, *Tetrahedron Lett.*, **47**, 5211 (2006).
 13. S-i. Kawaguchi, S. Nagata, A. Nomoto, M. Sonoda, and A. Ogawa, *J. Org. Chem.*, **73**, 7928 (2008).
 14. It is difficult to form phosphinylrhodium hydride from $\text{Rh}(\text{PPh}_3)_3\text{Cl}$ and Ph_2PH directly.