This article was downloaded by: [Monash University Library] On: 26 November 2014, At: 10:11 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/lncn20</u>

SYNTHESIS AND ANTIVIRAL EVALUATION OF ISATIN RIBONUCLEOSIDES

Mara R.P. deOliveira ^a , José C. Torres ^a , Simon J. Garden ^a , Carla Veronica B. dos Santos ^b , Thatyana Rocha Alves ^b , Angelo C. Pinto ^a , Helena de S. Pereira ^a , Luiz Roberto Leão Ferreira ^c , Nissin Moussatché ^a , Izabel Christina de P. P. Frugulhetti ^c , Vitor F. Ferreira ^b & Maria Cecília B. V. de Souza ^d

^a Departamento de Química Orgânica, Instituto de Química, Universidade Federal do Rio de Janeiro, Ilha do Fundão, Rio de Janeiro, CEP 21945-970, Brazil

^b Departamento de Química Orgânica, Outeiro de São João Batista s/no, Centro, Niterói, Instituto de Química, Rio de Janeiro, CEP 21945-970, Brazil

^c Instituto de Biologia , Outeiro de São João Batista s/no, Centro, Niterói , Universidade Federal Fluminense , Rio de Janeiro, CEP 21945-970, Brazil

^d Departamento de Química Orgânica, Outeiro de São João Batista s/no, Centro, Niterói, Instituto de Química, Rio de Janeiro, CEP 21945-970, Brazil Published online: 31 Aug 2006.

To cite this article: Mara R.P. deOliveira, José C. Torres, Simon J. Garden, Carla Veronica B. dos Santos, Thatyana Rocha Alves, Angelo C. Pinto, Helena de S. Pereira, Luiz Roberto Leão Ferreira, Nissin Moussatché, Izabel Christina de P. P. Frugulhetti, Vitor F. Ferreira & Maria Cecília B. V. de Souza (2002) SYNTHESIS AND ANTIVIRAL EVALUATION OF ISATIN RIBONUCLEOSIDES, Nucleosides, Nucleotides and Nucleic Acids, 21:11-12, 825-835, DOI: <u>10.1081/NCN-120016510</u>

To link to this article: <u>http://dx.doi.org/10.1081/NCN-120016510</u>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

©2002 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

NUCLEOSIDES, NUCLEOTIDES & NUCLEIC ACIDS Vol. 21, Nos. 11 & 12, pp. 825–835, 2002

SYNTHESIS AND ANTIVIRAL EVALUATION OF ISATIN RIBONUCLEOSIDES

Mara R. P. de Oliveira,¹ José C. Torres,¹ Simon J. Garden,¹ Carla Veronica B. dos Santos,² Thatyana Rocha Alves,² Angelo C. Pinto,¹ Helena de S. Pereira,⁴ Luiz Roberto Leão Ferreira,³ Nissin Moussatché,⁴ Izabel Christina de P. P. Frugulhetti,³ Vitor F. Ferreira,² and Maria Cecília B. V. de Souza^{2,*}

 ¹Universidade Federal do Rio de Janeiro, Instituto de Química, Departamento de Química Orgânica, Ilha do Fundão, CEP 21945-970, Rio de Janeiro, Brazil
²Instituto de Química, Departamento de Química Orgânica, and ³Instituto de Biologia, Universidade Federal Fluminense,

Outeiro de São João Batista s/no, Centro, Niterói,

CEP 24210-150, Rio de Janeiro, Brazil

⁴Universidade Federal do Rio de Janeiro, Instituto de Biofísica Carlos Chagas Filho, Ilha do Fundão, CEP 21941-590, Rio de Janeiro, Brazil

ABSTRACT

A series of novel substituted isatin ribonucleosides 3b-3f were synthesized in good yields by a TMSOTf catalysed coupling reaction between the silylated nitrogenated base (1b-1f) and 1-O-acetyl-2,3,5-tri-O-benzoyl- β -D-ribofuranose (2). Isatin nucleoside 3a previously reported was also prepared using this method giving high yield. From the compounds

825

DOI: 10.1081/NCN-120016510 Copyright © 2002 by Marcel Dekker, Inc. 1525-7770 (Print); 1532-2335 (Online) www.dekker.com

^{*}Corresponding author. Fax: 55 021 21 27193349; E-mail: gqocica@vm.uff.br

tested, ribonucleoside **3f** proved to be the most active one when assayed for antiviral activity on HSV-1 infected cells, leading to 66% of inhibition of virus yield. All the isatin derivatives tested did not inhibit HIV-1 Reverse Transcriptase (RT) activity.

The synthesis of new ribonucleosides continues to be important for the search of therapeutically useful agents.^[1] Isatin derivatives^[2] have been found to possess a wide variety of pharmacological properties as follows- monoamine oxidase inhibitor,^[3] antibacterial^[4] and antiviral activities.^[4b,4c,5] However, the literature reports few studies on the preparation of isatin nucleosides.^[6–9]

As part of an ongoing program^[10,11] for investigating the transformation of isatins into new derivatives with potential biological activity, this work describes the synthesis of five new isatin ribonucleosides: 5-methyl-2,3-dihydro-1-(2,3,5-tri-*O*-benzoyl- β -D-ribofuranosyl)indole-2,3-dione (**3b**), 5-fluoro-2,3-dihydro-1-(2,3,5-tri-*O*-benzoyl- β -D-ribofuranosyl)indole-2,3-dione (**3c**), 5-chloro-2,3-dihydro-1-(2,3,5-tri-*O*-benzoyl- β -D-ribofuranosyl)indole-2,3-dione (**3d**), 5-iodo-2,3-dihydro-1-(2,3,5-tri-*O*-benzoyl- β -D-ribofuranosyl) indole-2,3-dione (**3e**), 4,6-dibromo-2,3-dihydro-1-(2,3,5-tri-*O*-benzoyl- β -Dribofuranosyl)indole-2,3-dione (**3f**) as well as derivative 2,3-dihydro-1-(2,3,5tri-*O*-benzoyl- β -D-ribofuranosyl)indole-2,3-dione (**3a**)^[6a] previously reported. These isatin ribonucleosides were screened for their in vitro antiviral properties in HSV-1 infected cells. These compounds were also tested on HIV-1 RT activity. The reaction sequence for the preparation of the target compounds is outlined in Sch. 1.

RESULTS AND DISCUSSION

Isatin derivatives used in this work (**1a–1f**) were prepared according to published procedures.^[10,11] The method chosen for the synthesis of the corresponding ribonucleosides was the coupling between the previously silylated nitrogenated base and 1-*O*-acetyl-2,3,5-tri-O-benzoyl- β -D-ribofuranose (**2**) using Lewis acid catalysis.^[12] Heating **1** with bis(trimethylsilyl)trifluoro-acetamide (BSTFA)^[13] containing 1% of trimethylchlorosilane (TMSCI), at 60°C, under nitrogen atmosphere, for 16 h, followed by reaction with **2**, in the presence of trimethylsilyltrifluoroamethanesulfonate (TMSOTf)^[14] as catalyst, at room temperature, for 6 h, gave high yields of the corresponding ribonucleosides **3a–3f** (Sch. 1) (**3a**: 90%; **3b**: 80%; **3c**: 82%; **3d**: 78%; **3e**: 78%; **3f**: 80%). The use of a mixture of hexamethyldisilazane/trimethylchlorosilane as the silylating agent and toluene as solvent led to clear solutions which failed to give any nucleoside product.

Although the de-O-benzoylation of 3a is successfully reported in the literature,^[6a] all attempts to remove the O-benzoyl groups from 3a-3f using



Scheme 1. Synthetic route used for the preparation of isatin nucleosides 3a-3f.

different basic or acidic reagents such as methanol/sodium methoxide solution, methanol/sodium bicarbonate solution and Dowex H^+/H_2O , under different reaction conditions of time and temperature, led to nucleoside decomposition products, probably as a result of the initial fragmentation of N-glycosyl bond. This cleavage, in basic conditions, may be rationalized by the attack of the alcoxide ion on the most reactive benzoyl group (C-2), followed by a fast β -attack of the oxygen anion from the ortho-ester to the anomeric center, eliminating the isatin moiety, which is a good leaving group due to the vicinal carbonyl, as outlined in Sch. 2. Thus, it seems that this cleavage reverses the glycosyl bond by the same kind of mechanism of the regiocontrolled and stereospecific glycoylation of isatin derivatives with 1-Oacetyl-2,3,5-tri-O-benzoyl- β -D-ribofuranose (2). This cleavage was also observed in FAB mass spectra.

The structures of the nucleosides were confirmed by IR, UV, ¹H and ¹³C NMR experiments [¹H (Table 1), ¹³C (Table 2), DEPT, ¹H-¹H-COSY, HET-COR (¹J_{CH} and ⁿJ_{CH}, n = 2, 3)] and by FAB-high resolution mass spectra which revealed the molecular-ion peaks (M+H)⁺as well as the ion m/z 445 attributed to the riboside fragment resulting from C-N cleavage of glicosyl bond.

¹H and ¹³C spectra of 3a-3f showed very similar chemical shifts for the riboside moiety in each case (Tables 1 and 2).

The observation of nuclear Overhauser effect (n.O.e.) for the H4' signals while the anomeric proton was being irradiated (**3a**: 3.09%; **3b**: 2.46%; **3c**: 3.72%; **3d**: 3.04%; **3e**: 3.88%; **3f**: 2.97%) confirmed the β -configuration of all nucleosides.^[15] The point of attachment of the sugar to the bases assigned to the



R= H or Me

Scheme 2. Cleavage of the N-glycosyl bond of 3a-3f under basic conditions.

nitrogen atom from the isatin nucleus was based upon carbon-13 NMR chemical shifts for C2 and C3. These were compared with data for N-alkylated isatins in the literature.^[11] Additional proof for N1-ribosylation was also visible through the nuclear Overhauser effect in the spectra of the compounds where the irradiation of the anomeric hydrogen, H1', resulted in an *observed* n.O.e. on H7 (**3a**: 5.45%; **3b**: 3.28%; **3c**: 6.11%; **3d**: 6.89%; **3e**: 7.11%; **3f**: 5.25%).

BIOLOGICAL STUDIES

HSV-1 infected cells were incubated with the isatins and isatin ribonucleosides at a concentration of 50 μ M for 72 h. The antiviral activity was determined according to the computation of the 50 percent endpoint method described by Reed and Muench.^[5] Among the heterocycles, **1c** and **1f** were found to reduce virus yield significantly. Ribonucleosides **3a**, **3b** and **3f** proved to be more active than the corresponding isatin heterocycles, while the heterocycle **1c** was more active than the corresponding ribonucleoside. Thus, these five compounds were considered to be the most efficient derivatives, leading to statistically significant inhibition of virus yield (Table 3). The most efficient isatin derivatives were still less efficient than the potent anti HSV-1 compound, acyclovir, which reduced virus yield nearly 100% at the same concentration. Sincitium formation was exclusively observed in infected cells incubated with ribonucleoside **3c**. Compound **3d** was toxic to VERO cells and therefore was not tested.

Further investigation on the antiviral activities of the heterocycles 1a-1fand the ribonucleosides 3a-3f was assessed by measuring [3H]dTTP incorporation in a poly(rA).oligo(dT) template primer by the activity of the

ISATIN RIBONUCLEOSIDES

		<i>Table 1.</i> ¹ H (300.00	MHz) NMR Spectral D	ata for 3a–3f [(CDCl ₃ /'	TMS), J (Hz)]	
H	3a	3b	3c	3d	3e	3f
4 v	7.66–7.58 (m)	7.43–7.33 (m)	7.30 (dd, 6.6, 2.7)	7.65–7.48 (m)	7.88 (d, 1.8)	
9	7.13–7.06 (m)	- 6.90 (dd, 8.1, 1.2)	6.75 (dt, 8.8, 2.7)	7.01 (dd, 8.7, 2.1)	7.33 (dd, 8.4, 1.8)	(IIII) <i>CC.1</i> -0 + .1
7	7.24–7.21 (m)	7.10 (d, 8.1)	7.21 (dd, 3.6, 9.0)	7.18 (d, 8.7)	7.01 (d, 8.4)	7.48–7.35 (m)
1′	6.25 (d, 6.3)	6.23 (d, 6.0)	6.23 (d, 6.0)	6.19 (d, 6.3)	6.18 (d, 6.0)	6.09 (d, 6.0)
5	6.13 (t, 6.3)	6.12 (t, 6.0)	6.10 (t, 6.3)	6.10 (t, 6.1)	6.1.0 (t, 6.3)	6.20 (t, 6.0)
3,	6.05 (dd, 6.3, 4.5)	6.04 (dd, 6.6, 4.8)	6.04 (dd, 6.3, 4.5)	6.03 (dd, 6.3, 4.5)	6.02 (dd, 6.6, 4.5)	6.07–6.03 (m)
4	4.77–4.68 (m)	4.73–4.66 (m)	4.72–4.67 (m)	4.72–4.66 (m)	4.70–4.65 (m)	4.75–4.66 (m)
5'a	4.96–4.90 (m)	4.95–4.86 (m)	4.95 (dd, 13.2, 3.3)	4.94 (dd, 13.5, 3.9)	4.94 (dd, 13.5, 3.6)	4.94 (dd, 13.5, 3.9)
5'b	4.74–4.60 (m)	4.73–4.66 (m)	4.72–4.67 (m)	4.72–4.66 (m)	4.70–4.65 (m)	4.75–4.66 (m)
2''	8.13 (dd, 8.4, 1.5)	8.13 (dd, 8.4, 1.5)	8.12 (dd, 8.4, 1.5)	8.11 (dd, 8.4, 1.2)	8.10 (dd, 8.4, 1.2)	8.08 (dd, 8.4, 1.5)
	7.97 (dd, 8.4, 1.5)	7.97 (dd, 8.4, 1.5)	7.98 (dd, 8.4, 1.2)	7.98 (dd, 8.1, 1.2)	7.98 (dd, 8.4, 1.5)	7.99 (dd, 8.4, 1.2)
	7.93 (dd, 8.7, 1.5)	7.92 (dd, 8.4, 1.5)	7.92 (dd, 8.4, 1.5)	7.91 (dd, 8.4, 1.2)	7.91 (dd, 8.1, 1.2)	7.93 (dd, 8.4, 1.2)
3''	7.57–7.46 (m)	7.57–7.47 (m)	7.68–7.49 (m)	7.65–7.48 (m)	7.69–7.48 (m)	7.48–7.35 (m)
	7.41–7.33 (m)	7.43–7.33 (m)	7.43–7.30 (m)	7.30–7.33 (m)	7.43–7.31 (m)	
4"	7.66–7.58 (m)	7.66–7.58 (m)	7.68–7.49 (m)	7.65–7.48 (m)	7.69–7.48 (m)	7.62–7.53 (m)
	7.57–7.46(m)	7.57–7.47 (m)				
<u>C</u> H ₃	Ι	2.25 (s)	Ι	Ι	Ι	I

Downloaded by [Monash University Library] at 10:11 26 November 2014

829

		<i>Table</i> 2. ¹³ C (75.0 MH	(z) NMR Spectral Data fo	or 3a–3f [(CDCl ₃ /TMS	[(§	
¹³ C	3a	3b	3c	3d	3e	3f
5	157.6	157.8	157.6	157.0	156.6	156.4
3	181.5	181.8	180.93	180.6	180.2	179.9
3a	117.7	117.8	118.6 (d, 7.5)	118.7	119.3	115.4
4	125.6	125.8	112.3 (d, 24.0)	125.3	134.0	122.34
5	124.2	134.1	159.3 (d, 246.0)	130.1	86.9	131.5
6	138.2	138.7	124.6 (d, 23.5)	137.5	146.2	133.3
7	112.2	112.0	113.8 (d, 7)	113.6	114.3	114.4
7a	148.3	146.1	144.4	146.6	147.7	150.6
1′	84.3	84.2	84.4	84.5	84.7	84.8
2'	70.3 or 70.2	70.3 or 70.2	70.3 or 70.2	70.3	70.3	71.0
3'	70.3 or 70.2	70.3 or 70.2	70.3 or 70.2	70.3	70.3	70.6
4′	79.9	79.8	80.0	80.0	84.5	80.2
5'	63.3	63.4	63.2	63.2	63.1	63.4
1′′	129.2, 128.2	129.2, 128.4,	129.2, 128.5,	129.2, 128.1	129.2, 128.1	129.2, 128.5
		128.1	128.1			
2''	129.7,	129.7, 129.6,	129.7, 129.6,	129.7, 129.6,	129.7, 129.6,	129.73, 129.7,
	129.6, 129.5	129.5	129.5	129.5	129.5	129.6
3//	128.5,	128.5, 128.3	128.6, 128.4,	128.6, 128.5,	128.6, 128.4	128.6, 128.4
	128.4, 128.3		128.3	128.4		
4"	133.6,	133.6, 133.5,	133.7, 133.6,	133.7, 133.6,	133.7, 133.6,	133.7, 133.6,
	133.5, 133.3	133.3	133.5	133.5	133.5	133.4
\overline{CH}_3		20.4	I	Ι		
C=0	165.8,	165.8, 165.2,	165.8, 165.2,	165.8, 165.2,	165.8, 165.2,	166.0, 165.2,
	165.2, 165.1	165.1	165.1	165.1	165.1	165.1

830

DE OLIVEIRA ET AL.

Downloaded by [Monash University Library] at 10:11 26 November 2014

Table 3. Antiviral Activity of Isatins 1a-1f and Isatin Ribonucleosides 3a-3f (50 μ M) on HSV-1 Yield

Compounds	Herpes Simplex Type 1 Virus (% of Inhibition of Virus Yield)
1a	0
1b	- 16
1c	60*
1d	28
1e	28
lf	46*
3a	61*
3b	46*
3c	15
3d	N.D.
3e	37
3f	66*

N.D. = not determined.

Vero cells (10^6 cells/well) were infected with HSV-1 at 1PFU/cell for 2 h. At zero time post-infection, virus inocullum was replaced by culture medium containing or not isatin derivatives. After 72 h post-infection, virus yield of each sample was determined by 50% tissue culture infective dose (TCID₅₀/mL). The data are mean values from one of the representative experiments.

*P values < 0.05 for comparing drug-treated and non-treated cells.

recombinant HIV-1 reverse transcriptase (HIV-1 RT), as already described.^[17] Our results showed that none of the isatin derivatives reduced the RNA-dependent DNA polymerase activity of the recombinant HIV-1 RT.

In conclusion, the modest but significant anti-HSV-1 activities of the isatin derivatives described here suggests that these compounds may be selectively targeted to HSV-1 replication, since they were not cytotoxic to VERO cells at the concentration tested.

EXPERIMENTAL

General Procedures

Melting points were determined on a Fisher-Johns apparatus and are uncorrected. The IR spectra were recorded on a Perkin-Elmer 1420 spectrometer as potassium bromide pellets and frequencies are expressed in cm⁻¹. Ultraviolet (UV) spectra were obtained on a Schimadzu spectrophotometer; λ are in nm and ϵ in mol⁻¹ cm⁻¹. Mass spectra analysis

(FABHRMS) were recorded in a 3-nitrobenzylalcohol matrix in the positive ion mode on a VG ZAB-E mass spectrometer, by Department of Chemistry and Biochemistry, University of Oklahoma. NMR spectra were recorded on a Varian Unity Plus 300 spectrometer operating at 300.00 MHz (¹H) and 75.0 MHz (¹³C), in specified solvents. Chemical shifts are reported in ppm (δ) relative to tetramethylsilane. Proton and carbon spectra were typically obtained at room temperature. The two dimensional experiments were acquired using standard Varian Associates automated programs for data acquisition and processing.

General Procedure for the Preparation of 2,3-Dihydro-1-(2,3,5-tri-Obenzovl-B-D-ribofuranosvl)indole-2,3-dione 5-Methyl-2,3-dihydro-1-(**3**a), (2,3,5-tri-*O*-benzoyl-β-D-ribofuranosyl)indole-2,3-dione (3b), 5-Fluoro-2,3dihydro-1-(2,3,5-tri-O-benzoyl-β-D-ribofuranosyl)indole-2,3-dione (3c), 5-Chloro-2,3-dihydro-1-(2,3,5-tri-O-benzoyl-B-D-ribofuranosyl)indole-2,3-dione (3d), 5-Iodo-2,3-dihydro-1-(2,3,5-tri-*O*-benzoyl-β-D-ribofuranosyl)indole-2,3dione (3e) and 4,6-Dibromo-2,3-dihydro-1-(2,3,5-tri-O-benzoyl-β-D-ribofuranosyl)indole-2,3-dione (3f). A mixture of isatin 1a-1f (2.0 mmol) and bis(trimethylsilyl)trifluoroacetamide (BSTFA, 7.46 mmol) containing 1% of trimethylchlorosilane, in acetonitrile (4mL) was stirred at 60°C, under nitrogen atmosphere, for 16 h. The reaction mixture was allowed to cool to room temperature and a solution of 1-O-acetyl-2,3,5-tri-O-benzoyl-β-Dribofuranose (2.1 mmol) in 15 mL of acetonitrile was added. This procedure was followed by the dropwise addition of a solution of TMSOTf (2.1 mmol) in 10 mL of acetonitrile. After stirring for 6 h. the resulting mixture was poured into ice-cold water (20g). Afterwards, it was neutralized with a saturated aqueous solution of sodium bicarbonate and extracted with ethyl acetate $(3 \times 30 \text{ mL})$. The combined organic phases were dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel using hexane:methylenechloride gradient (up to 40%) as the eluant, to give:

3a: as orange crystals (1.08 g, 90%), m.p. 73°C; UV λ_{max} (CHCl₃) 411 (ϵ 426), 244 (ϵ 35,449); IR 1730 (C=O), 1260 (C-O); FABHRMS m/z calcd for C₃₄H₂₆NO₉ (M+H)⁺ 592.1619, found 592.1607; ¹H NMR (300.00 MHz, CDCl₃) see Table 1; ¹³C NMR (75.0 MHz, CDCl₃) see Table 2.

3b: as orange crystals (0.97 g, 80%), m.p. 78°C; UV λ_{max} (CHCl₃) 424 (ϵ 406), 244 (ϵ 17,939); IR 1730 (C=O), 1260 (C-O); FABHRMS m/z calcd for C₃₅H₂₈NO₉ (M+H)⁺ 606.1715, found 606.1171; ¹H NMR (300.00 MHz, CDCl₃) see Table 1; ¹³C NMR (75.0 MHz, CDCl₃) see Table 2.

3c: as orange crystals (1.00 g, 82%), m.p. 86°C; UV λ_{max} (CHCl₃) 411 (ϵ 633), 245 (ϵ 25,443); IR 1725 (C=O), 1260 (C-O); FABHRMS m/z calcd

ISATIN RIBONUCLEOSIDES

for $C_{34}H_{25}NO_9F$ (M+H)⁺ 610.1525, found 610.1513; ¹H NMR (300.00 MHz, CDCl₃) see Table 1; ¹³C NMR (75 MHz, CDCl₃) see Table 2.

3d: as orange crystals (0.98 g, 78%), m.p. 86°C; UV λ_{max} (CHCl₃) 419 (ϵ 609), 245 (ϵ 25,749); IR 1730 (C=O), 1260 (C-O); FABHRMS m/z calcd for C₃₄H₂₅NO₉³⁵Cl (M+H)⁺ 626.5229; found 626.1217; ¹H NMR (300.00 MHz, CDCl₃) see Table 1; ¹³C NMR (75.0 MHz, CDCl₃) see Table 2.

3e: as orange crystals (1.10 g, 78%), m.p. 83°C; UV λ_{max} (CHCl₃) 424 (ϵ 989), 243 (ϵ 22,198); IR 1730 (C=O), 1260 (C-O); FABHRMS m/z calcd for C₃₄H₂₅NO₉I (M+H)⁺ 718.0585, found 718.0574; ¹H-NMR (CDCl₃) see Table 1; ¹³C-NMR (CDCl₃) see Table 2.

3f: as yellow crystals (1.20 g, 80%), m.p. 223°C; UV λ_{max} (CHCl₃) 406 (ϵ 922), 243 (ϵ 19,324); IR 1730 (C=O), 1260 (C-O); FABHRMS m/z calcd for C₃₄H₂₄NO₉⁸¹Br₂ (M+H)⁺ 751.9790, found 751.9788; ¹H NMR (300.00 MHz, CDCl₃) see Table 1; ¹³C NMR (75.0 MHz, CDCl₃) see Table 2.

Cells. VERO epithelial cell line derived from monkey kidney cells obtained from the American tissue culture Collection (Rockville, MD) were grown in monolayer cultures in Dulbecco's modified Eagle's medium (Gibco laboratories) containing 2% heated-inactivated fetal bovine serum (FBS; Pigue Farm), 8% calf serum (CS; Pan-American Center of aftose fever), 2.25% sodium bicarbonate, 500 U/mL penicilin, 100 µg/mL streptomycin, 50 µg/mL gentamycin, 2.5 µg/mL fungison. Cell cultures were incubated at 37°C in humidified air containing 5% CO₂. For passaging, confluent monolayers (6.5×10^5 cells) were washed with phosphate-buffered saline solution (NaCl, 0.8 g%, KCl, 0.02 g%, Na₂HPO₄, 0.15 g%, NaH₂PO₄, 0.02 g%, EDTA, 1 mM PBS pH 7.2), and after a short trypsinisation (Trypsin at 0.25 g%) the cells were suspended in the culture medium.

Virus. HSV-1 were kindly provided by Dr. Marcia Wigg (Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brasil), propagated in Hep-2-cells and stored at -70° C until use.

Antiviral Activity. Antiviral activity on HSV-1 virus infection was assessed by virus yield assay. Briefly, sub-confluent VERO cells, grown in 96-well plates (10^6 cells/well), were infected with HSV-1 at 1 PFU (plaque forming unity)/cell for 120 min at 37° C in $100 \,\mu$ L of DMEM. After this period, virus inocullum was replaced by culture medium, containing isatin derivatives or not. We defined this point as the zero time post-infection. After 72 h post-infection, culture medium was harvested and virus titer of each sample was quantified in terms of the 50% tissue culture infective dose (TCID₅₀/mL) by endpoint dilution, according to Reed and Muench.^[16]

ACKNOWLEDGMENTS

Fellowship granted to M.R.P.O. from CNPq (Brazil) is gratefully acknowledged. A.C.P., S.J.G., M.C.B.V.S. and V.F.F. are grateful to CNPq for their individual research fellowships. This work was partially supported by CNPq (National Council of Research from Brazil) and FAPERJ.

REFERENCES

- Kolb, V.M. Novel and Unusual Nucleosides as Drugs. Prog. Drug Res. 1997, 48, 195–232.
- da Silva, J.F.M.; Garden, S.J.; Pinto, A.C. The Chemistry of Isatins: A Review from 1975 to 1999. J. Braz. Chem. Soc. 2001, 12 (3), 273–286.
- Medvedev, A.E.; Clow, A.; Sandler, M.; Glover, V. Isatin: A Link Between Natriuretic Peptides and Monoamines. Biochem.-Pharmacol. 1996, 52 (3), 385–391.
- (a) Atta-Ur-Rehman; Ijaz, A.S.; Choudhary, M.I.; Amtul, Z.; Atta-Ur-Rahman. New Isatin Derivatives and Their Antimicrobial Activity. J. Chem. Soc. Pakistan, 1997, 19 (3), 230–236; (b) Pandeya, S.N.; Sriram, D.; Nath, G.; de Clercq, E. Synthesis, Antibacterial; Antifungal and Anti HIV-Activities of Norfloxacin Mannich Bases. Eur. J. Med. Chem. 2000, 35 (2), 249–255; (c) Pandeya, S.N.; Sriram, D.; Nath, G.; de Clercq, E. Synthesis, Antibacterial, Antifungal and Anti HIV Activity of Schiff and Mannich Bases Derived from Isatin Derivatives and N-[4-(4'-chlorophenyl)thiazol-2-yl]thiosemicarbazide. Eur. J. Pharm. Sci. 1999, 9, 25–31.
- Teitz, Y.; Barko, N.; Abramoff, M.; Ronen, D. Relationships Between Structure and Antiretroviral Activity of Thiosemicarbazone Derivatives. Chemotherapy 1994, 40 (3), 195–200.
- (a) Yartseva, I.V.; Ektova, L.V.; Preobrazhenskaya, M.N. 1-Ribosylisatins. Bioorg. Khim. 1975, 1 (2), 189–194; (b) Yartseva, I.V.; Ektova, L.V.; Preobrazhenskaya, M.N.; Lesnaya, N.A.; Yavorskaya, N.P.; Platonova, G.N.; Sofyina, Z.P. 1-Glucosylisatins and their Biological Activities on Animals with Transplantable Tumors and In vitro. Bioorg. Khim. 1975, 1 (11), 1589–1592; (c) Yartseva, I.V.; Ektova, L.V.; Sakharova, V.I.; Dobrynin, Y.V.; Yavorskaya, N.P.; Nikolaeva, T.G.; Sofina, Z.P.; Preobrazhenskaya, M.N. Synthesis and Study of 1-Glycosylations. Zh. Org. Khim. 1977, 13 (8), 1743–1749.
- Dobrynin, Y.V.; Nikolaeva, T.G.; Shkrgova, A.D.; Lesnaya, N.A.; Peretolchina, N.M.; Sofyina, Z.P.; Éktova, L.V.; Tolkachev, V.N.; Preobrazhenskaya, M.N. Structure-Cyto-Toxic Activity Relationship in 1-Glycosylisatins 3-Derivatives. Khim.-Farm. Zh. 1984, 18 (12), 1440–1444.
- Éktova, L.V.; Tolkachev, V.N.; Yartseva, I.V.; Paramonova, T.D.; Lesnaya, N.A.; Sofyina, Z.P.; Marennikova, S.S; Chekunova, E.V.; Preobrazhenskaya, M.N. Synthesis and Study of the Derivatives of 5-Bromo-Glycosylisatins, 6-Nitro-1-Glycosylisatins and 5-Bromo-6-Nitro-1-Glycosylisatins. Khim.-Farm. Zh. 1984, 7, 776–785.

ISATIN RIBONUCLEOSIDES

- Preobrazhenskaya, M.N.; Yartseva, I.V.; Éktova, L.V. 1-Beta-D-Glucopyranosides of Isatin and Methylisatins. Dokl. Akad. Nauk SSSR 1974, 215 (4), 873–876.
- (a) Garden S.J.; Torres, J.C.; Ferreira, A.A.; Silva, R.B.; Pinto, A.C. A Modified Sandmeyer Methodology and the Synthesis of (±) Convolutamydine A. Tetrahedron Lett. **1997**, *38* (9), 1501–1504 and references cited therein; (b) Garden, S.J.; Torres, J.C.; Melo, S.C.D.; Lima, A.S.; Pinto, A.C.; Lima, E.L.S. Aromatic Iodination in Aqueous Solution. A New Base of Life for Aqueous Potassium Dichloroiodate. Tetrahedron Lett. **2001**, *42* (11), 2089–2092; (c) Torres, J.C.; Garden, S.J.; Pinto, A.C. J. An Investigation of a Palladium Catalysed Biaryl Synthesis of Pyrrolophenanthidine Derivatives: Extension of Heck Reaction. J. Braz. Chem. Soc. **2000**, *11* (5), 441–446.
- Garden S.J.; Torres, J.C.; Silva L.E.; Pinto, A.C. A Convenient Methodology for the N-Alkylkation of Isatin Compounds. Synth. Commun. **1998**, 28 (9), 1679–1689.
- (a) Niedballa, U.; Vorbrüeggen, H. A General Synthesis of N-Glycosides. I. Synthesis of Pyrimidine Nucleosides J. Org. Chem. **1974**, *39* (25), 3654–3660;
 (b) Niedballa, U.; Vorbrüeggen, H. On the Mechanism of the Stannic Chloride Catalysed Silyl Hilbert-Johnson Reaction. J. Org. Chem. **1976**, *41* (12), 2084–2086.
- (a) da Matta, A.D.; Bernardino, A.M.R.; Romeiro, G.A.; de Oliveira, M.R.P.; de Souza, M.C.B.V.; Ferreira, V.F. Nucleosides Having Quinolone Derivatives as Nitrogenated Bases: Regiospecific and Stereospecific Ribosylation of 3-Carbethoxy-1,4-dihydro-4-oxoquinolines. Nucleosides & Nucleotides 1996, 15 (4), 889–898; (b) da Matta, A.D.; dos Santos, C.V.B.; Pereira, H.S.; Frugulhetti, I.C.P.P.; de Oliveira, M.R.P.; de Souza M.C.B.V.; Moussatché, N.; Ferreira, V.F. Synthesis of Novel Nucleosides of 4-Oxoquinoline-3-carboxylic Acid Analogues. Heteroatom Chemistry 1999, 10 (3), 197–202.
- 14. Vorbrüeggen, H.; Krolikiewicz, K.; Bennua, B. Nucleoside Syntheses, XXII. Nucleoside Synthesis with Trimethylsilyl Triflate and Perchlorate as Catalysts. Chem. Ber. **1981**, *114* (4), 1234–1255.
- Rosemeyer, H.; Tóth, G.; Seela, F. Assignment of Anomeric Configuration of D-Ribonucleosides, Arabinonucleosides, 2'-Desoxyribonucleosides by NOE Difference Spectroscopy. Nucleosides & Nucleotides 1989, 8, 587–597.
- Reed L.J.; Muench, M.A. A Simple Method of Estimating Fifty Percent End Points. Am. J. Hyg. 1938, 27, 493–497.
- 17. Hizi, A.; Mc Gill, C.; Hughes, H. Expression of Soluble, Enzymatically Active Human Immunodeficiency Virus Reverse Transcriptase in Escherichia coli and Analysis o Mutants. Proc. Natl. Acad. Sci. USA **1988**, *85*, 1218–1222.

Received February 8, 2002 Accepted August 26, 2002