

Fused Tetrazoles as Azide Surrogates in Click Reaction: Efficient Synthesis of N-Heterocycle-Substituted 1,2,3-Triazoles


Buddhadeb Chattopadhyay, Claudia I. Rivera Vera, Stepan Chuprakov, and Vladimir Gevorgyan*

Department of Chemistry, University of Illinois at Chicago, 845 West Taylor Street, Chicago, Illinois 60607-7061

vlad@uic.edu

Received March 30, 2010

ABSTRACT





It has been shown that various pyrido-, quinolino-, pyrazino-, and quinoxalinotetrazoles can be used efficiently as azide components in Cu-catalyzed click reaction with alkynes. This method allows for efficient synthesis of a wide variety of N-heterocyclic derivatives of 1,2,3-triazoles.

1,2,3-Triazoles are biologically important units.¹ Pyridotriazoles and quinolinotriazoles are particularly interesting as they exhibit a wide range of biological properties, including control of arthropod pests,^{2a} substance-related disorders,^{2b} ATP-competitive inhibition of vascular endothelial growth factor receptors I and II,^{2c} and antibacterial^{2d} and antimicrobial activity.^{2e}

Unarguably, Cu-catalyzed click chemistry³ of azide with alkyne is the most efficient way to assemble the 1,2,3-triazole ring⁴ (eq 1). However, preparation of pyrido- and quinoli-

notriazoles is not straightforward because these azides exist in equilibrium between closed form (tetrazole **A**) and open form (azide **B**) (eq 2).⁵




(1) (a) Alvarez, R.; Valaquez, S.; San, F.; Aquaro, S.; Penro, C. D. C. F.; Karlsson, A.; Balzarini, J.; Camarasa, M. J. *J. Med. Chem.* **1994**, *37*, 4185. (b) Buckle, D. R.; Rockell, C. J. M.; Smith, H.; Spicer, B. A. *J. Med. Chem.* **1986**, *29*, 2262. (c) Genin, M. J.; Allwine, D. A.; Anderson, D. J.; Barbachyn, M. R.; Emmert, D. E.; Garmon, S. A.; Graber, D. R.; Grege, K. C.; Hester, J. B.; Hutchinson, D. K.; Morris, J.; Reischer, R. J.; Ford, C. W.; Zurenko, G. E.; Hamel, J. C.; Schaadt, R. D.; Stapert, D.; Yagi, B. H. *J. Med. Chem.* **2000**, *43*, 953.

(2) (a) Bretschneider, T.; Franken, E.-M.; Goergens, U.; Fuesslein, M.; Hense, A.; Kluth, J.; Schwarz, H.-G.; Koehler, A.; Malsam, O.; Voerste, A. PCT Int. Appl. 2010, WO 2010006713 A2 20100121. (b) Garzya, V.; Watson, S. P. PCT Int. Appl. 2009, WO 2009115486 A1 20090924. (c) Kiselyov, A. S.; Semenova, M.; Semenov, V. V. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 1344. (d) Gordeev, M. F.; Yuan, Z.; Liu, J. PCT Int. Appl. 2008, WO 2008108988 A1 20080912. (e) Japelj, B.; Rečnik, S.; Cebasek, P.; Stanovnik, B.; Svete, J. *J. Heterocycl. Chem.* **2005**, *42*, 1167.

Usually, the position of this equilibrium depends on several factors, such as the nature of substituents,⁵ solvent,⁶ and temperature.⁵ Thus, it has been reported^{7a} that a NO₂ group at the C-6 position of tetrazole favors the open form (azide **B**). On the contrary, tetrazoles with NO₂, COOH, and Cl groups at the C-8 position and unsubstituted tetrazole

(3) (a) Sharpless, K. B.; Finn, M. G. *Angew. Chem.* **2001**, *113*, 2056; *Angew. Chem., Int. Ed.* **2001**, *40*, 2004. (b) Rostovtsev, V. V.; Green, L. G.; Fokin, V. V.; Sharpless, K. B. *Angew. Chem.* **2002**, *114*, 2708; *Angew. Chem., Int. Ed.* **2002**, *41*, 2596. (c) Tornøe, C. W.; Christensen, C.; Meldal, M. *J. Org. Chem.* **2002**, *67*, 3057.


predominantly exist^{7,8} in closed form **A**. It should be mentioned that pyridotetrazole has been employed in the preparation of organometallic complexes of late transition metals.^{7a} Furthermore, there have been contradictory reports^{9,10} on the employment of tetrazoles in the click reaction. For instance, it has been shown that pyridotetrazoles, existing in closed form, are inert toward click reaction under standard conditions.⁹ However, there have been two reports^{10a,b} in which single examples of successful click reaction of in situ generated pyridotetrazoles with alkynes were demonstrated. Moreover, when this manuscript was in preparation, a paper describing successful click reaction of purinotetrazole, which mainly exists in an open form, has appeared.^{10c} Accordingly, motivated by the high biological importance of pyridyl- and quinoliny- containing triazoles² and intrigued by the contradictory results on the employment of triazoles in click reaction,^{9,10} we undertook an investigation aiming at the development of an efficient method using differently substituted tetrazoles in the synthesis of heterocyclic derivatives of 1,2,3-triazoles. Herein, we wish to report that various pyrido-, quinolino-, pyrazino-, and quinoxalinotetrazoles **1** can efficiently be employed in click reaction with alkynes to give the corresponding heterocyclic derivatives of 1,2,3-triazoles **2** (eq 3).



We first examined the reaction of tetrazole **1a** with phenyl acetylene employing the most popular^{3b} click chemistry

conditions (Table 1, entry 1). However, no formation of desired product **2a** was observed. Employment of other

Table 1. Optimization of Click Reaction of Tetrazoles



no.	catalyst, 10 mol %	solvent, 0.25 M	<i>t</i> [°C]	time [h]	yield [%] ^a
	CuSO ₄ ·5H ₂ O,				
1	Na-ascorbate	DCM/H ₂ O (1:1)	60	24	0
2	CuI	toluene	100	24	10
3	Cu(OTf) ₂	toluene	100	24	5
4	(CuOTf) ₂ ·C ₆ H ₆	toluene	100	2	52 ^b
5	(CuOTf) ₂ ·C ₆ H ₆	toluene	rt	7	81
6	(CuOTf) ₂ ·C ₆ H ₆	THF	60	12	76
7	(CuOTf) ₂ ·C ₆ H ₆	DCE	100	24	0
8	(CuOTf) ₂ ·C ₆ H ₆	1,4-dioxane	100	24	0

^a Isolated yields. ^b Some decomposition products were found.

copper salts was more effective. Thus, when the reaction was performed in the presence of 10 mol % CuI,^{4a} it afforded the product **2a** in 10% yield (entry 2). Use of Cu(OTf)₂⁴ⁱ gave 5% of product (entry 3). A substantial improvement of the yield (52%) has been achieved with (CuOTf)₂·C₆H₆^{4j} (entry 4). Gratifyingly, analogous reaction at room temperature gave 81% of **2a** (entry 5). THF was equally efficient as toluene in the reaction (entry 6). Switching to other solvents (entries 7 and 8) was not beneficial for this reaction.

With the optimized conditions in hand, we tested the generality of the click reaction of tetrazoles (Table 2). To our delight, these newly developed conditions appeared to be very general for a spectrum of N-fused tetrazoles, giving an easy access to 1,4-triazoles **2**. Thus, reaction of ester-containing pyridotetrazole (**1a**) with various alkynes proceeded smoothly at room temperature to produce differently substituted pyridyl-containing triazoles in good to excellent yields (entries 1–12). Reactions of unsubstituted (**1b**) and C-5 methyl-substituted (**1c**) tetrazoles were efficient at elevated temperatures (entries 13–21). It was also found that various N-fused heterocyclic tetrazoles, such as quinolinotetrazoles (**1d**, entries 22–28), pyrazinotetrazole (**1e**, entry 29 and 30), and quinoxalinotetrazole (**1f**, entries 31 and 32), successfully underwent click reaction to give the corresponding N-heterocycle-substituted 1,4-triazoles **2** in good yields. These reaction conditions appeared to be very general with respect to the alkyne component, as alkynes possessing various alkyl, aryl, alkenyl, benzyl, homobenzyl, ester, trimethylsilyl, alkyl chloride, secondary alcohol, acetal, thiophenyl, and even sugar groups provided good to high yields of triazoles **2**.

After developing the “tetrazole-clicking” approach for the synthesis of 1,4-triazoles, we next examined the

(4) For original [3 + 2] cycloaddition, see: Huisgen, R. *Angew. Chem.* **1963**, 75, 604. For selected recent reviews of click chemistry, see: (a) Meldal, M.; Tornøe, C. W. *Chem. Rev.* **2008**, 108, 2952. (b) Bock, V. D.; Hiemstra, H.; van Maarseveen, J. H. *Eur. J. Org. Chem.* **2006**, 51. (c) Nandivada, Jiang, H. X.; Lahann, J. *Adv. Mater.* **2007**, 19, 2197. (d) Tron, G. C.; Pirali, T.; Billington, R. A.; Canonico, P. L.; Sorba, G.; Genazzani, A. A. *Med. Res. Rev.* **2008**, 28, 278. (e) Becer, C. R.; Hoogenboom, R.; Schubert, U. S. *Angew. Chem., Int. Ed.* **2009**, 48, 4900. (f) Spiteri, C.; Moses, J. E. *Angew. Chem., Int. Ed.* **2010**, 49, 31. (g) Kolb, H. C.; Finn, M. G.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2001**, 40, 2004. (h) Lutz, J.-F. *Angew. Chem., Int. Ed.* **2007**, 46, 1018. (i) Fukuzawa, S.-i.; Shimizu, E.; Kikuchi, S. *Synlett* **2007**, 2436. (j) Aucagne, V.; Berna, J.; Crowley, J. D.; Goldup, S. M.; Hanni, K. D.; Leigh, D. A.; Lusby, P. J.; Ronaldson, V. E.; Slawin, A. M. Z.; Viterisi, A.; Walker, D. B. *J. Am. Chem. Soc.* **2007**, 129, 11950.

(5) (a) Boyer, J. H.; Miller, E. J. *J. Am. Chem. Soc.* **1959**, 81, 4671. (b) Lowe-Ma, Ch. K.; Nissan, R. A.; Wilson, W. S. *J. Org. Chem.* **1990**, 55, 3755. (c) Evans, R. A.; Wentrup, C. *J. Chem. Soc., Chem. Commun.* **1992**, 1062. (d) Sasaki, T.; Kanematsu, K.; Murata, M. *J. Org. Chem.* **1971**, 36, 446.

(6) Pochinok, V. V.; Avramenko, L. F.; Grigorenko, P. S.; Skopenko, V. N. *Russ. Chem. Rev.* **1975**, 44, 481, and references therein.

(7) (a) Pizzotti, M.; Cenini, S.; Porta, F. *J. Chem. Soc., Dalton Trans.* **1978**, 1155. (b) Messmer, A.; Juhasz-Riedl, H. Z.; Sohar, P. *J. Org. Chem.* **1988**, 53, 973. (c) Wentrup, C. *Tetrahedron* **1970**, 26, 4969. (d) *The Chemistry of Azido Group*; Patai, S., Ed.; Interscience: London, 1971. (e) Reimlinger, H. *Chem. Ber* **1970**, 103, 1900, and references therein.

(8) Kanyalkar, M.; Coutinho, E. C. *Tetrahedron* **2000**, 56, 8775.

(9) Colombano, G.; Travelli, C.; Galli, U.; Caldarelli, A.; Chini, M. G.; Canonico, P. L.; Sorba, G.; Bifulco, G.; Tron, G. C.; Genazzani, A. A. *J. Med. Chem.* **2010**, 53, 616.

(10) (a) Klein, M.; Diner, P.; Dorin-Semlat, D.; Doerig, C.; Grotli, M. *Org. Biomol. Chem.* **2009**, 7, 3421. (b) Saha, B.; Sharma, S.; Sawant, D.; Kundu, B. *Synlett* **2007**, 1591. (c) Lakshman, M. K.; Singh, M.; Parrish, D.; Balachandran, R.; Day, B. W. *J. Org. Chem.* **2010**, Published ASAP Mar 18, 2010; DOI: 10.1021/jo902342z.


Table 3. Toward the Synthesis of 1,5-Disubstituted Triazoles^a

no.	substrate	catalyst	4	5
1		RuCpCl(PPh ₃) ₂	-	-
2		RuCp*Cl(PPh ₃) ₂	-	-
3		(CuOTf) ₂ C ₆ H ₆	81%	-
4		RuCp*Cl(PPh ₃) ₂	23%	36% ^[b]
5		(CuOTf) ₂ C ₆ H ₆	91%	-
6		RuCp*Cl(PPh ₃) ₂	3%	56%
7		(CuOTf) ₂ C ₆ H ₆	87%	-

^a Isolated yield. Reaction conditions: 5 mol % catalyst, 1,4-dioxane 0.25 M, 110 °C (entries 1, 2, 4 and 6); 10 mol % catalyst, PhMe 0.25 M, 100 °C (entries 3, 5 and 7). ^b Inseparable mixture of **4** and **5** (1:1.5).

of 3-azido- and 4-azido-pyridines proceeded uneventfully providing 1,4-disubstituted triazoles in excellent yields

(entries 5 and 7). Thus, it became evident that under the Ru-catalysis tested, pyridotetrazoles could not be used as precursors for 1,5-disubstituted triazoles.



In summary, it has been shown that pyrido-, quinilino-, pyrazino-, and quinoxalinotetrazoles, which exist in open/close form equilibrium (between **1** and **1'**, eq 4) can be employed as azide surrogates in the Cu-catalyzed click reaction. This reaction is efficient with a wide variety of alkynes to produce N-heterocyclic derivatives of 1,4-disubstituted triazoles **2**. It has also been found that, probably because of deactivation of the Ru-catalyst, pyridotetrazoles cannot be used as azide precursors in the synthesis of 1,5-disubstituted triazoles.

Acknowledgment. The support of the NIH (GM-6444) is gratefully acknowledged.

Supporting Information Available: Experimental procedures and characterization of new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL100745D