A new combination of donor and acceptor: $bis(\eta^6$ -benzene)chromium and hexafluorobenzene form a charge-transfer stacked crystal

Catherine J. Aspley,^{*a*} Clive Boxwell,^{*a*} Maria L. Buil,^{*a*} Catherine L. Higgitt,^{*a*} Conor Long^{*b*} and Robin N. Perutz^{**a*}

^a Department of Chemistry, University of York, York, UK YO10 5DD. E-mail: rnp1@york.ac.uk ^b Department of Chemistry, Dublin City University, Dublin, Ireland

Received (in Oxford, UK) 3rd February 1999, Accepted 23rd April 1999

Bis(η^6 -benzene)chromium reacts with hexafluorobenzene to yield a red charge-transfer complex [Cr(η^6 -C₆H₆)₂•C₆F₆] which contains stacks of alternating donor and acceptor molecules with close inter- and intrastack contacts; in addition to the charge-transfer complex, formation of [Cr(η^6 -C₆H₆)₂]⁺ is demonstrated by EPR and IR spectroscopy.

Non-covalent interactions have a major role in determining structures of molecular assemblies. The formation of stacks of donors alternating with acceptors can lead to special electronic and magnetic properties as in $[Fe(\eta^5-C_5Me_5)_2][TCNE]$ or $[Fe(\eta^6-C_6Me_3H_3)_2][C_6(CN)_6]$.^{1,2} Stacking of aromatic rings (π stacks) represents another motif, this time driven by quadrupolar interactions, e.g. in co-crystals of benzene and hexafluorobenzene.³ Complex analogues of $C_6H_6 \cdot C_6F_6$ are attracting interest for their optoelectronic properties.4,5 Bis(η^6 -benzene)chromium 1 has potential for both these types of interaction: it has featured as a donor in $[Cr(\eta^{6} C_6H_6_2$ [S₂O₆•2SO₂] and has been employed for crystal engineering.^{6,7} Donor-acceptor (DA) complexes are also formed by 1 in $[Cr(\eta^6-C_6H_6)_2][TCNE]$ and $[Cr(\eta^6-C_6Me_3H_3)_2][TCNQ]$ but they are arranged in stacks of the type ... A₂DD... or in chains.8

Several authors have proposed that an electron-transfer process precedes C–F bond activation of perfluoroalkanes and perfluoroarenes by transition metal complexes,^{9–11} although experimental support for redox initiation is sparse. If this proposal is to be reconciled with the redox potentials, there must be either preassociation of the fluorocarbon and transition metal complex, or the products must be stabilised by rapid irreversible dissociation (*e.g.* $C_6F_6^-$ to $C_6F_5 + F^-$), or more likely both.¹² Crabtree *et al.* recently reported that $[Fe(\eta^5-C_5R_5)_2]$ (R = H or Me) associates with perfluoronaphthalene or perfluorophenanthrene in the solid state to form an irregular DA stack.¹³ However, there was no optical evidence for a charge-transfer (CT) interaction. We report that 1 reacts with hexafluorobenzene to form a red CT complex in solution which crystallises as a donor–acceptor stack, and additionally that 1 undergoes one-electron oxidation by C_6F_6 to a limited extent.

When a large excess of dry hexafluorobenzene is condensed onto a freshly sublimed sample of 1 and thawed under argon, a claret-coloured solution is obtained from which crystals grow over several weeks at room temperature. The crystals have two habits, red-pink blocks and thin yellow plates, both of which are stable only in the mother liquor. An X-ray crystal structure of a red-pink crystal mounted in its mother liquor in a capillary at room temperature revealed it to be $[Cr(\eta^6-C_6H_6)_2 \cdot C_6F_6]$.† The structure contains face-to-face stacks of $[Cr(\eta^6-C_6H_6)_2]$ moieties alternating with C_6F_6 units at regular intervals of *ca*. 3.5 Å, a value similar to that in $[Fe(\eta^5-C_5Me_5)_2][TCNE]^2$ The stacks lie approximately parallel to the body diagonal of the triclinic unit cell (Fig. 1) such that the benzene rings of the [Cr(η^{6} - C_6H_6] units eclipse the C_6F_6 rings in the same stack. The angle between the ring normals of the C_6F_6 and $[Cr(\eta^6-C_6H_6)_2]$ units is 5.9°. Each C_6F_6 lies approximately in line with one of the

 C_6H_6 rings in a neighbouring stack. The minimum C···C distance between C_6H_6 and C_6F_6 lies within a stack, while the minimum Cr···F distance lies between stacks (Fig. 2, Table 1). The regularly spaced DA stacks contrast with those of $[Fe(\eta^5-C_5Me_5)_2 \bullet C_{14}F_{10}]^{13}$ and [1][TCNE].⁸

We have also attempted to solve the structure of the thin yellow plates at -100 °C. Although the crystal did not diffract well, we found that the structure was little changed from that of the red–pink blocks. A compression along *b* and *c* led to a contraction of 1.9% in the minimum intrastack C…C distance and 3.7% in the minimum interstack Cr…F distance.



Fig. 1 Packing diagram for $[Cr(\eta^6-C_6H_6)_2 \bullet C_6F_6]$, viewed down the body diagonal of the triclinic cell (hydrogen atoms omitted).



Fig. 2 ORTEP diagram (ref. 15) showing the key intra- and inter-stack spacings within the crystal structure of $[Cr(\eta^6-C_6H_6)_2 \cdot C_6F_6]$, viewed down the *c* axis. The hydrogen atoms are omitted for clarity. The average C–C and Cr–C bond lengths within a $Cr(C_6H_6)_2$ unit are 1.397 and 2.134 Å. For the C_6F_6 molecule, the C–C and C–F bond lengths average 1.365 and 1.341 Å.

Table 1 Minimum contact distances in the structure of $[Cr(\eta^{6-}C_{6}H_{6})_{2}\bullet C_{6}F_{6}]$

Labela	Type of contact	Contact atoms	Distance/Å
A B C D	Intrastack C ₆ H ₆ C ₆ F ₆ Intrastack CrC ₆ F ₆ Intrastack CrC ₆ F ₆ Interstack CrC ₆ F ₆	$r(C \cdots C)$ $r(Cr \cdots C)$ $r(Cr \cdots F)$ $r(Cr \cdots F)$	3.480(4) 5.074(3) 5.312(3) 4.490(2)
^{<i>a</i>} labels A, B, C, D refer to Fig. 2.			

When the reaction of 1 with C_6F_6 was carried out with higher concentrations of 1, a fine yellow precipitate separated from the claret solution. A UV–VIS absorption spectrum of the solution, measured after filtering off the precipitate, showed an absorption at 503 nm and a shoulder at 390 nm.¹⁶⁺ On freezing in liquid nitrogen, the claret solution turned to a yellow glass, but the claret colour returned on melting. The corresponding absorption maxima lie to longer wavelength (542, 414 nm) on reaction of $[Cr(\eta^6-1,4-C_6H_4Me_2)_2]$ with C_6F_6 and to shorter wavelength (424 nm)§ on reaction of 1 with C_6F_5H . These absorption bands are assigned to charge-transfer transitions of the complexes $[Cr(\eta^6-arene)_2 \cdot C_6F_{6-n}H_n]$ (n = 0, 1) which must be present in solution. Redox potentials suggest that the ground state of the complex will be close to the A•D description and the excited state close to A+•D⁻.¶

The solutions from reaction of 1 with C_6F_6 (whether dilute or more concentrated) gave a broad EPR signal at g = 1.987consistent with formation of $[Cr(\eta^6-C_6H_6)_2]^+$ 1⁺.⁶ Solid-state EPR spectra of the yellow precipitate from reaction of 1 with C_6F_6 also revealed the presence of $1^+(g_{\parallel} = 2.002, g_{\perp} = 1.983)$, while IR spectra showed characteristic bands of both 1 and 1^{+.17} The presence of fluoride as a corresponding anion was revealed by its reaction with Me₃SiOTf yielding Me₃SiF. In order to ascertain the proportion of 1 which is oxidised, we investigated the effect of addition of C_6F_6 to toluene solutions of **1**. A control sample of 1 in toluene showed only traces of 1⁺. On addition of 2 and 5 equiv. of C_6F_6 , well-resolved resonances ($A_H = 3.4$ G) for 1^+ were observed with intensities *ca*. 20 fold and 48 fold greater than the control, respectively. Comparison with the resonance of a standard solution of TEMPO (10-4 mol dm-3 in toluene) provided lower-limiting estimates that the solutions of 1 were 0.5 and 3.3% oxidised, respectively.** The extent of oxidation is appreciably higher than expected to arise from impurities in the hexafluorobenzene. The formation of $[Cr(C_6H_6)_2]$ +F⁻ is reminiscent of the reaction of cobaltocene with perfluoroalkanes.10

These experiments lead to the following conclusions. (i) A donor-acceptor complex is formed between [$Cr(\eta^6-C_6H_6)_2$] **1** and C_6F_6 with a long-wavelength absorption in solution, assigned to a CT transition between **1** and C_6F_6 . (ii) The complex crystallises as [$Cr(\eta^6-C_6H_6)_2 \bullet C_6F_6$] with a \cdots DADA \cdots stacked structure with close intra- and inter-stack contacts. The claret colour is observed in the larger crystals. The structure is probably stabilised by charge-transfer and $\pi-\pi$ interactions. (iii) In addition to [$Cr(\eta^6-C_6H_6)_2 \bullet C_6F_6$], salts including [$Cr(\eta^6-C_6H_6)_2$]+F— are formed in low conversion although half-wave potentials suggest that **1** is incapable of reducing C_6F_6 .¶ (iv) The formation of the donor-acceptor complex and the oxidation of **1** provide support for related mechanisms for reactions of metal-hydride complexes with fluoroarenes.¹¹, ¹²

We thank Dr T. Braun and Professors N. Connelly and W. E. Geiger for helpful discussions, the referees for helpful criticism, S. Foxon and Dr A. C. Whitwood for experimental help and the EPSRC and the European Commission for support.

Notes and references

† *Crystal data* for [Cr(η⁶-C₆H₆)₂•C₆F₆]: C₁₈H₁₂CrF₆, M = 394.28, redpink blocks, triclinic, space group PI, a = 7.3254(8), b = 9.2627(9), c = 6.6102(8) Å, $\alpha = 100.026(9)$, $\beta = 112.327(8)$, $\gamma = 97.532(9)^\circ$, V = 398.87(8) Å³, Z = 1, T = 293(2) K, μ (Mo-K α) = 0.778 mm⁻¹, F(000) = 198, 1477 reflections measured, 1401 unique ($R_{int} = 0.040$), 116 parameters. The crystal was mounted in a capillary in its mother liquor. The structure was solved by direct methods (SHELX) (ref. 14) and refined by full-matrix least-squares on F^2 . Goodness of fit on F^2 1.083, final R1 [$I > 2\sigma(I)$] R1 = 0.0392, wR2 = 0.1027. CCDC 182/1238. See http://www.rsc.org/suppdata/cc/1999/1027/ for crystallographic files in .cif format.

 \ddagger [Cr(C₆H₆)₂] is yellow–green with a weak band at 640 nm and an intense band at 320 nm. The corresponding cation exhibits weak bands in the near-IR region and a more intense band at 340 nm (ref. 16). The CT band of Cr(η^{6} -C₆H₆)₂•SO₂ is observed at 540 nm (ref. 6).

§ The short-wavelength shoulder will overlap the bands of $[Cr(\eta^6\text{-}C_6H_6)_2]$ in $C_6F_5H.$

 \P The half-wave potentials of $C_6F_6/C_6F_6^-$ and $[Cr(\eta^6-C_6H_6)_2]^+/$ $[Cr(\eta^6-C_6H_6)_2]$ are -2.55 and -0.68 V respectively vs. SCE. (ref. 18).

|| Characterisation of the anion in the yellow precipitate was complicated by paramagnetic 1⁺. The precipitate was dissolved in CD₂Cl₂, Me₃SiOTf added and the volatiles condensed into an NMR tube. Me₃SiF was identified by the eight central lines of the decet at δ –157.96 (J_{HF} 7.6 Hz) in the ¹⁹F NMR spectrum (ref. 19). A control experiment with C₆F₆, CD₂Cl₂ and Me₃SiOTf generated only traces of Me₃SiF.

** Complex **1** was freshly sublimed before use. Solutions in toluene (0.047 mol dm⁻³) were made up in an argon-filled glove-box; C_6F_6 (99.9%), previously dried over molecular sieves and degassed by freeze-pump-thaw methods, was added in the box with a microsyringe. Typical impurities are C_6F_5Cl and $C_6F_5Cl_2$. Since the addition of C_6F_6 causes slight precipitation, the values of the percentage oxidation are lower limits.

- 1 J. S. Miller and A. J. Epstein, Angew. Chem., Int. Ed. Engl., 1994, 33, 385.
- 2 J. S. Miller, J. C. Calabrese, H. Rommelmann, S. R. Chittipeddi, J. H. Zhang, W. M. Reiff and A. J. Epstein, *J. Am. Chem. Soc.*, 1987, **109**, 769; M. D. Ward, *Organometallics*, 1987, **6**, 754; M. D. Ward and J. C. Calabrese, *Organometallics*, 1989, **8**, 593.
- 3 J. H. Williams, Acc. Chem. Res., 1993, 26, 593; A. P. Weat, S. Mecozzi and D. A. Dougherty, J. Phys. Org. Chem., 1997, 10, 347.
- 4 G. W. Coates, A. R. Dunn, L. M. Henling, J. W. Ziller, E. B. Lobkovsky and R. H. Grubbs, J. Am. Chem. Soc., 1998, 120, 3641.
- 5 P. Kirsch and K. Tarumi, Angew. Chem., Int. Ed., 1998, 37, 484.
- 6 C. Elschenbroich, R. Gondrum and W. Massa, Angew. Chem., Int. Ed. Engl., 1985, 24, 967.
- 7 D. Braga, A. L. Costa, F. Grepioni, L. Scaccianoce and E. Tagliavini, Organometallics, 1997, 16, 2070.
- 8 J. Š. Miller, D. M. O'Hare, A. Chakraborty and A. J. Epstein, J. Am. Chem. Soc., 1989, 111, 7853; D. O' Hare, M. D. Ward and J. S. Miller, Chem. Mater., 1990, 2, 758.
- 9 J. Burdeniuc and R. H. Crabtree, J. Am. Chem. Soc., 1996, 118, 2525; Organometallics, 1998, 17, 1582.
- 10 B. K. Bennett, R. G. Harrison and T. G. Richmond, J. Am. Chem. Soc., 1994, **116**, 11165.
- 11 M. K. Whittlesey, R. N. Perutz and M. H. Moore, *Chem. Commun.*, 1996, 787.
- 12 B. L. Edelbach and W. D. Jones, J. Am. Chem. Soc., 1997, 119, 7734.
- 13 C. M. Beck, J. Burdeniuc, R. H. Crabtree, A. L. Rheingold and G. P. A. Yap, *Inorg. Chim. Acta*, 1998, **270**, 559.
- 14 G. M. Sheldrick, SHELXL 93, Program for the Refinement of Crystal Structures, University of Göttingen, 1995.
- 15 C. K. Johnson, ORTEP, Report ORNL-5138, Oak Ridge National Laboratory, Oak Ridge, TN, 1976.
- 16 K. D. Warren, Struct. Bonding (Berlin), 1976, 27, 45.
- 17 H. P. Fritz, W. Lüttke, H. Stammreich and R. Forneris, Spectrochim. Acta, 1961, 17, 1068.
- 18 J. A. Marsella, A. G. Galicinski, A. M. Coughlin and G. P. Pez, J. Org. Chem., 1992, 57, 2856; D. Astruc, Electron Transfer and Radical Processes in Transition Metal Chemistry, VCH, Weinheim, 1995.
- 19 D. C. England, F. J. Weigart and J. C. Calabrese, J. Org. Chem., 1984, 49, 4816.

Communication 9/00919A