

Nitration of moderately deactivated arenes with nitrogen dioxide and molecular oxygen under neutral conditions. Zeolite-induced enhancement of regioselectivity and reversal of isomer ratios

Xinhua Peng,^a Naoyuki Fukui,^b Masayuki Mizuta^b and Hitomi Suzuki^{*b}

^a Department of Chemistry, School of Chemical Engineering, Nanjing University of Science and Technology, Xiao Ling Wei, Nanjing 210094, China. E-mail: xinhpeng@public1.ptt.js.cn

^b Department of Chemistry, School of Science and Technology, Kwansei Gakuin University, Gakuen 2-1, Sanda 669-1337, Japan. E-mail: hsuzuki@ksc.kwansei.ac.jp; Fax: +81-79-565-9077

Received 19th February 2003, Accepted 2nd May 2003

First published as an Advance Article on the web 29th May 2003

In the presence of zeolites, moderately deactivated arenes such as 1-nitronaphthalene, naphthonitriles, and methylated benzonitriles can be smoothly nitrated at room temperature by the combined action of nitrogen dioxide and molecular oxygen. The regioselectivity is considerably improved as compared with the conventional nitration methodology based on nitric and sulfuric acids. In some cases, the minor isomer became favoured to a significant extent, resulting in the reversal of ordinary isomer ratios of nitration products.

Introduction


In recent years, there has been a growing worldwide concern over the adverse effect of chemical wastes that arise from industrial activities. With regard to this, aromatic nitration is quite notorious for its unfriendly nature towards the environment.¹ The use of large quantities of nitric and sulfuric acids, corrosiveness, potential danger of explosion, low regioselectivity, oxidative degradation byproducts, and inefficient atomic economy all constitute the disadvantages of this time-honored reaction. Among a number of recent efforts to develop alternative methodologies, the combined use of nitrogen oxides (NO and NO₂) and oxygen in place of the classical nitric acid–sulfuric acid system appears to be the most attractive and promising. As everybody knows, commercial nitric acid is manufactured from lower nitrogen oxides *via* an energy-consuming multi-step process.

In the presence of molecular oxygen (dioxygen) and an appropriate catalyst, nitrogen dioxide is activated to react with a wide variety of nonactivated and moderately activated aromatic substrates to afford the corresponding nitro compounds in good yield.² However, the isomeric composition of the nitration products obtained resembles in most cases those of classical nitration based on nitric and sulfuric acids. This means the formation of a large proportion of unwanted products, *i.e.*, inevitable loss of materials. Smith and coworkers have recently reported that halobenzenes can be nitrated with nitrogen dioxide and dioxygen in the presence of zeolite H β catalyst to give products of nitration, in which the *para*-nitro isomer predominates over the *ortho*-isomer (*ortho* : *para* = 0.10–0.24).³ The *ortho* : *para*-isomer ratio of conventional nitration is usually in the range 0.6–0.8. We have also observed a significant improvement of the isomer product ratios in the zeolite-assisted double Kyodai-nitration of toluene and chlorobenzene.⁴ In the present paper, we wish to report a remarkable example of zeolite catalysis in the *nonacid* aromatic nitration, where the interaction between a polar substituent and the acidic sites on the inorganic solid surface would considerably improve the regioselectivity of nitration and in certain cases it can even lead to the reversal of isomer ratio of nitration products.

Results and discussion

Nitration of 1-nitronaphthalene

1,5-Dinitronaphthalene **2** and 1,8-dinitronaphthalene **3** are useful intermediates in the manufacture of dyestuffs, pigments, and high-energy materials. The former compound is also important as the precursor to high performance polyurethane resins and consequently of greater market demand than the latter. Industrial preparation of these polynitronaphthalenes still employs the traditional methodology involving a mixture of nitric and sulfuric acids, where the 1,5- : 1,8-dinitro isomer ratio of around 0.5–0.6 is always obtained either by direct nitration of naphthalene or by further nitration of 1-nitronaphthalene **1** (Table 1). The second nitro group preferentially occupies the position *peri* to the first nitro group irrespective of the nitrating conditions employed (Scheme 1).



The Kyodai-nitration which uses a combination of lower nitrogen oxides and ozone as the nitrating agent has demonstrated excellent conversion of a wide variety of aromatic compounds into the corresponding nitro derivatives.¹³ The non-acidic nature of this methodology is highly attractive and environmentally friendly. Unfortunately, when this nitrating system was applied to the nitration of compound **1**, the isomer proportion of the dinitration product was again similar to those of classical methodologies (Table 2, runs 1–3). No obvious improvement was observed in the presence of solid inorganic catalysts.

However, when compound **1** was subjected to the combined action of nitrogen dioxide and dioxygen in the presence of a

Table 1 Nitration of 1-nitronaphthalene with nitric acid alone or mixed acid

Entry	Nitrating agent	Reaction conditions	Yield (%)	Isomer proportion (%)			Ref.
				5-Nitro 2	8-Nitro 3	2 : 3 ratio	
1	HNO ₃ (<i>d</i> = 1.42)-H ₂ SO ₄ (<i>d</i> = 1.84)	No solvent, 0 °C	—	31	69	0.45	5
2	HNO ₃ (<i>d</i> = 1.42)-H ₂ SO ₄ (<i>d</i> = 1.84)	No solvent, 0 °C	98.1	33.5	66.5	0.50	6
3	HNO ₃ (<i>d</i> = 1.42)-H ₂ SO ₄ (<i>d</i> = 1.84)	(CH ₃ CO) ₂ O, −40 °C	—	32.4	67.6	0.48	6
4	HNO ₃ (<i>d</i> = 1.5)-H ₂ SO ₄ (<i>d</i> = 1.84)	BF ₃ , CH ₃ CO ₂ H, 0 °C	98.3	33.3	66.7	0.50	6
5	HNO ₃ -H ₂ SO ₄	No solvent, −55 °C	—	22	78	0.28	7
6	KNO ₃ -H ₂ SO ₄ (<i>d</i> = 1.84)	No solvent, 0 °C, 24 h	—	36	64	0.56	8
7	HNO ₃ (<i>d</i> = 1.42)	No solvent, reflux	— ^a	37.2	56.5	0.66	9
8 ^c	53%-HNO ₃ -98%H ₂ SO ₄	No solvent, 37–80 °C	93.7 ^b	27.4	72.6	0.37	11
9 ^c	NO ₂ -BF ₃	CH ₃ NO ₂ , −5 °C	—	39	61	0.64	9
10 ^c	fum-HNO ₃ -Claycop ^d	(CH ₃ CO) ₂ O-CCl ₄ , rt	93	25	75	0.33	12

^a Accompanied by 1,3,8-trinitronaphthalene. ^b Accompanied by 1,4,5-trinitronaphthalene. ^c Direct dinitration of naphthalene. ^d Montmorillonite K10 impregnated with anhydrous cupric nitrate.

Table 2 Nitration of 1-nitronaphthalene with nitrogen dioxide and ozone or dioxygen in the presence of solid inorganic catalysts^a

Run	Nitrating agent	Solvent	Catalyst ^b	Additive	Reaction time/h	Conversion ^c (%)	Yield ^d (%)	Isomer proportion (%) ^e		2 : 3 Ratio
								5-Nitro 2	8-Nitro 3	
1	NO ₂ -O ₃	CH ₂ Cl ₂	—	—	2	>99	89	23	77	0.30
2	NO ₂ -O ₃	CH ₂ Cl ₂	HBEA-25	—	2	>99	89	25	75	0.33
3	NO ₂ -O ₃	CH ₂ Cl ₂	K10 ^f	—	1	95	87	19	81	0.23
4	NO ₂ -O ₂	CH ₃ CN	K10	—	7	100	90	26	74	0.35
5	NO ₂ -O ₂ ^g	CH ₂ Cl ₂	—	—	48	51	17	45	55	0.82
6	NO ₂ -O ₂ ^h	CH ₂ Cl ₂	—	—	48	74	27	38	62	0.61
7	NO ₂ -O ₂ ⁱ	CH ₃ CN	—	—	48	89	58	40	60	0.67
8	NO ₂ -O ₂	CH ₃ CN	HZSM-5	—	7	35	21	52	48	1.08
9	NO ₂ -O ₂	CH ₃ CN	HBEA-25	—	7	94	69	68	32	2.12
10	NO ₂ -O ₂	CH ₃ CN	HBEA-25 ^j	—	7	82	65	67	33	2.03
11	NO ₂ -O ₂	CH ₃ CN	HBEA-25 ^k	—	48	82	51	66	34	1.94
12	NO ₂ -O ₂	CH ₃ CN	HBEA-25	CH ₃ SO ₃ H ^l	7	98	55	62	38	1.63
13	NO ₂ -O ₂	CH ₃ CN	HBEA-25	Fe(acac) ₃ ^m	7	96	55	58	42	1.38
14	NO ₂ -O ₂	CH ₃ CN	HBEA-25	H ₂ SO ₄	7	>99	77	64	36	1.78
15	NO ₂ -O ₂	CH ₃ CN	HBEA-25	FeCl ₃ ^l	7	>99	78	68	32	2.13
16	NO ₂ -O ₂	CH ₃ CN	HBEA-150	—	7	99	85	72	28	2.57
17	NO ₂ -O ₂	CH ₃ CN	FeBEA-150 ⁿ	—	7	98	86	66	34	1.94
18	NO ₂ -O ₂	CH ₃ CN	NaMFI-90	—	7	23	19	60	40	1.50
19	NO ₂ -O ₂	CH ₃ CN	HSZ-390HOA	—	7	55	33	71	29	2.45
20	NO ₂ -O ₂	CH ₃ CN	HSZ-690HOA	—	7	40	32	68	32	2.13

^a All reactions with the NO₂-O₃ system were conducted at −10 °C using substrate **1** (9–10 mmol), liquid NO₂ (2.0 mL), catalyst (1.0 g), and CH₂Cl₂ (30 mL) under the Kyodai conditions, while those with the NO₂-O₂ system were carried out at room temperature using **1** (1.0 mmol), liquid NO₂ (0.5 mL) and catalyst (2.0 g) in CH₃CN (5 mL) under an oxygen atmosphere, unless otherwise specified. ^b Zeolites and K10 clay were calcined at 500 and 300 °C, respectively, for 8 h in air prior to use. ^c Determined by GC. ^d Combined yield of **2** and **3** based on consumed **1**. ^e Calculated from GC peak areas, excluding other isomers and byproducts. ^f Double amount of catalyst was used. ^g Double amount of NO₂ was used. ^h Amount of NO₂ was increased to 2.0 mL. ⁱ Amount of substrate was increased to 2.9 mmol. ^j Amount of catalyst was decreased to 1.0 g. ^k Substrate (2.9 mmol) and catalyst (1.0 g) were used. ^l Around 20 mmol% amount to the substrate was added. ^m A 13 mmol% amount to the substrate was added. ⁿ Prepared by stirring a suspension of HBEA-150 (5 g) in 0.2 M aq. FeCl₃ solution (100 mL) for 24 h under reflux.

zeolite, the formation of **2** became highly favored, leading to the reversal of the 1,5- : 1,8-dinitro isomer ratios. Especially, in the presence of zeolite HBEA-150 or HSZ-390HOA, regioselection could be remarkably shifted towards the 1,5-dinitro isomer, the 1,5- : 1,8-dinitro isomer ratio reaching as high as 2.5–2.6 (runs 16 and 19). Polar solvent systems appeared to play a positive role in attaining a higher 1,5- : 1,8-dinitro isomer ratio. Both Brønsted and Lewis acids proved to be effective in the enhancement of reaction rate (runs 12–15). The H-type β-zeolite was superior to the Fe³⁺-exchanged one in regioselection, probably due to the high density of effective sites that could interact *via* the hydrogen bonding (runs 16 and 17). Zeolite HBEA-150 with a high Si : Al ratio showed a better 1,5-site selectivity than HBEA-25 with a lower ratio possibly due to the dealumination, which would modify the distribution of the size and shape of the pores in the zeolites. Y-type zeolite HSZ-390HOA with three-dimensional channels of a large pore size exhibited high regioselectivity, probably due to its structural similarity to

HBEA-150. Mordenite zeolite HSZ-690HOA with a linear array of large pores showed high regioselection but low conversion. Both selectivity and conversion were unsatisfactory with zeolites HZSM-5 and NaMFI-90 (runs 8 and 18). This may be attributed to the medium pore size structure, which would place more restriction on the transport of a substrate through the pores.


Montmorillonite K10 facilitated the reaction and gave a cleaner nitration product as compared with others. Unfortunately, the isomer ratio was not much different from those of classical nitration reactions. Air could be used as a partner of nitrogen dioxide, although oxygen shortened the reaction time much more. Recovered zeolites can be recycled without any appreciable loss of catalytic activity. It is most likely that the electrostatic interaction as well as the hydrogen bonding between the nitro group and the acidic sites forbid free mobility of **1** within the zeolite cages and render the *peri*-position of **1** less accessible toward NO₂⁺ ions, thus leading to the reversal of the 1,5 : 1,8-dinitro isomer ratio.

A similar nonacid nitration of 2-nitronaphthalene was not investigated because of the strong mutagenic and carcinogenic nature of this compound.

Nitration of naphthonitriles, methylbenzonitriles (tolunitriles), and dimethylbenzonitriles (xylonitriles)

As an extension, we have investigated a similar nonacid nitration of another type of deactivated arene, aromatic nitriles, which included 1- and 2-naphthonitriles **4** and **7**, 2-, 3- and 4-methylbenzonitriles (*o*-, *m*- and *p*-tolunitriles) **10**, **13** and **17**, and three isomeric dimethylbenzonitriles (2,4-, 2,5- and 3,4-xylonitriles) **19**, **23** and **26**. Nitro derivatives of these nitriles are versatile starting materials for a variety of nitrogen-containing heterocyclic compounds of pharmaceutical and agrochemical interest. However, they are usually obtained not by direct nitration of nitriles but by the dehydration of nitroamides and nitroaldoximes,¹⁴ treatment of diazotized nitroamines with CuCN¹⁵ and prolonged heating of halonitriles with CuCN in DMF.¹⁶ This could be due to the inconvenient separation of isomers which often form in comparable amounts as well as the susceptibility of the nitrile function toward strong protonic acids, especially at elevated temperatures.

Nitration of 1- and 2-naphthonitriles. Both 1- and 2-naphthonitriles **4** and **7** underwent a rapid Kyodai-nitration to give a mixture of 5-nitro and 8-nitro derivatives **5–6** and **8–9** in almost quantitative combined yield (Scheme 2). In contrast to 1-nitronaphthalene **1**, the 5-nitro isomers **5** and **8** were always favoured over the corresponding 8-nitro isomers **6** and **9**, the 5-nitro : 8-nitro isomer ratios being 1.27 and 1.33, respectively. The values are close to those (1.44 and 1.38, respectively) observed in the conventional nitration based on nitric and sulfuric acids (Table 3). Nitrile **4** was previously nitrated with nitronium tetrafluoroborate¹⁷ and fuming nitric acid.¹⁸ However, no information was available on the isomer composition of the products.



Scheme 2


The nitriles **4** and **7** were quite slow to react with nitrogen dioxide and dioxygen at room temperature to form a comparable amount of 5-nitro : 8-nitro isomers. The ratios of both isomers **5** : **6** and **8** : **9** fell in the range 1.1–1.2, the 5-nitro isomer being a little favoured over the 8-nitro isomer. In the presence of zeolites, however, the reaction became considerably facilitated and within 12 h it was almost complete. Rather surprisingly, the 5-nitro isomer **5** was remarkably favoured at the expense of the 8-nitro isomer **6** in the nitration of 1-naphthonitrile **4**, the 5-nitro : 8-nitro ratio reaching as high as 9.0 (Table 3). In contrast, the 8-nitro isomer **9** became more important in the similar reaction of 2-naphthonitrile **7** and the 5-nitro : 8-nitro ratio **8** : **9** was reversed from near the unity down to 0.72.

The isomer ratios were considerably dependent on the type of zeolites employed. The highest regioselectivity was attained

with HBEA-150 for nitrile **4**, while the maximum reversal of isomer ratios was observed with HSZ-960HOA for nitrile **7**. A protonic acid such as methanesulfonic acid was also found to facilitate the nitration by nitrogen dioxide and dioxygen, but under these conditions the regioselection became worse, the 5-nitro : 8-nitro isomer ratios approaching unity.

Methylbenzonitriles (tolunitriles). As previously reported,¹⁹ benzonitrile readily undergoes the Kyodai nitration to afford an isomeric mixture of nitrobenzonitriles (*ortho* : *meta* : *para* = 15–22 : 72–80 : 5–6) in high yield. However, benzonitrile was found to be almost inert toward the combined action of nitrogen dioxide and dioxygen at room temperature, little nitration products being obtained even after prolong contact in the presence of solid catalyst.

Three isomeric methylbenzonitriles **10**, **13** and **17** were also inert toward nitrogen dioxide and dioxygen under similar conditions. In the presence of zeolites, however, the reaction became considerably facilitated and satisfactory results were obtained after 24 h (Scheme 3). The most attractive feature of the present nitration methodology is the enhanced regioselectivity as compared with the traditional and the Kyodai-nitrations. The reaction proceeded slowly, but the product was simple and clean with little formation of higher nitration products.



Scheme 3

2-Methylbenzonitrile (*o*-tolunitrile) **10** reacted with nitric and sulfuric acids to give a mixture of 3-nitro and 5-nitro isomers **11** and **12** in a ratio of 37 : 63 (Table 4). The Kyodai nitration of nitrile **10** readily took place at lower temperature and gave an isomeric mixture of a similar composition (39 : 61). When the nitrile was reacted with nitrogen dioxide and dioxygen in the presence of zeolite, the regioselectivity was remarkably improved and the 5-nitro isomer became the predominant product. Especially, in the case of HBEA-25, the 5-nitro isomer was the exclusive product at 77% conversion level. HZS-960 and 980HOA were more effective in accelerating the reaction, but the regioselectivity was decreased.

Table 3 Nitration of 1- and 2-naphthonitriles with nitrogen dioxide and ozone or dioxygen in the presence of zeolites^a


Nitrile	Nitrating agent ^a	Catalyst	Temp./°C	Reaction time/h	Conversion (%) ^b	Isomer proportion (%) ^b		Isomer ratio
						5-nitro	8-nitro	
1-Naphthonitrile 4	NO ₂ -O ₃ ^c	—	−10	0.5	>99	5	6	5 : 6
	HNO ₃ (<i>d</i> = 1.5)–H ₂ SO ₄ (<i>d</i> = 1.84)	—	RT	8	>99	56	44	1.27
	NO ₂ -O ₂	—	RT	24	>99	59	41	1.44
	NO ₂ -O ₂	CH ₃ SO ₃ H	RT	24	56	54	46	1.17
	NO ₂ -O ₂	HBEA-150	RT	12	>99	50	50	1.00
	NO ₂ -O ₂	Hβ-25	RT	12	>99	90	10	9.00
	NO ₂ -O ₂	HSZ-930HOA	RT	12	>99	86	14	6.14
	NO ₂ -O ₂	HSZ-940HOA	RT	12	>99	73	27	2.70
	NO ₂ -O ₂	HSZ-960HOA	RT	12	>99	69	31	2.23
	NO ₂ -O ₂	HSZ-980HOA	RT	12	>99	84	16	5.25
2-Naphthonitrile 7	NO ₂ -O ₃ ^c	—	−10	0.5	>99	8	9	8 : 9
	HNO ₃ (<i>d</i> = 1.5)–H ₂ SO ₄ (<i>d</i> = 1.84)	—	36	3	>99	57	43	1.33
	NO ₂ -O ₂	HBEA-150	RT	12	>99	58	42	1.38
	NO ₂ -O ₂	HSZ-930HOA	RT	12	>99	47	53	0.89
	NO ₂ -O ₂	HSZ-940HOA	RT	12	>99	43	57	0.75
	NO ₂ -O ₂	HSZ-960HOA	RT	12	>99	47	53	0.89
	NO ₂ -O ₂	HSZ-980HOA	RT	12	>99	42	58	0.72
	NO ₂ -O ₂	HSZ-980HOA	RT	12	>99	47	53	0.89

^a All reactions were carried out in acetonitrile in the presence of an excess of NO₂, unless otherwise noted. For detail, see the Experimental Section.^b Determined by GC. ^c Nitromethane was used as the solvent.

The traditional nitration of 3-methylbenzonitrile (*m*-tolunitrile) **13** at room temperature gave a mixture of 6-nitro, 5-nitro and 4-nitro isomers **16**, **15** and **14** in a ratio of 40 : 6 : 54. Interestingly, the Kyodai-nitration of this nitrile gave an isomeric mixture of a different composition (14 : 63 : 23). The minor product **15** from the traditional nitration became most important in the Kyodai-nitration. When nitrile **13** was allowed to react with nitrogen dioxide and dioxygen in the presence of zeolites, the regioselectivity was remarkably improved, mainly leading to the 6-nitro isomer **16** at the expense of other two isomers **14** and **15**. A single recrystallization from ethanol of the crude nitration product gave the 6-nitro derivative, almost free from other isomeric impurities. HSZ-960HOA was most effective in accelerating the reaction and also exhibited better regioselection. 4-Methylbenzonitrile (*p*-tolunitrile) **17** gave the 3-nitro derivative **18** as a single product. In this case, HBEA-150 was found to be most effective in facilitating the reaction. Methylated benzonitriles were previously nitrated with fuming nitric acid alone or with sulfuric acid,²⁰ potassium nitrate–sulfuric acid,²¹ acetyl nitrate,^{3c} and nitronium tetrafluoroborate.¹⁷ However, the literature contains no detail as to the isomer proportion of the nitration products.

Dimethylbenzonitriles (xylonitriles). Three isomeric dimethylbenzonitriles **19**, **23** and **26** were found to react more readily with nitrogen dioxide and dioxygen in the presence of zeolites, giving the corresponding nitro derivatives as an isomeric mixture (Scheme 4). In every case, the regioselectivity was remarkably improved as compared with both the traditional and Kyodai nitrations. Noteworthy is the nitration of nitrile **19**, where the 6-nitro isomer **22** was obtained as the single product in the presence of HSZ-960–980HOA (Table 5). Both the traditional and Kyodai-nitration of nitrile **19** afforded three isomeric nitration products in comparable amounts. Such remarkable regioselection may be attributed to the increased acidity of active sites as well as the fitness of pore sizes in HSZ-960–980HOA, since the conversion became quite low with HSZ-930–940HOA.

The traditional nitration of 2,4-dimethylbenzonitrile **23** usually produces 3- and 5-nitro isomers **24** and **25** in an approximate ratio of 2 : 5. In the presence of HSZ-960HOA, the regioselectivity is remarkably improved in favour of the latter isomer. From a single recrystallization in ethanol, compound **25** was

**Scheme 4**

obtained in a high state of purity. Both the traditional and the Kyodai nitrations of nitrile **26** gave 3- and 6-nitro isomers **29** and **27** as the major products. Zeolite HSZ-930HOA was found

Table 4 Nitration of methylbenzonitriles with nitrogen dioxide and ozone or dioxygen in the presence of zeolites^a

Nitrile	Nitrating agent ^a	Catalyst	Temp./°C	Reaction time/h	Conversion (%) ^b	Isomer proportion (%) ^b					Isomer ratio
2-Methylbenzonitrile (<i>o</i> -tolunitrile) 10						3-Nitro 11	5-Nitro 12	12 : 11			
	NO ₂ –O ₃	—	–10	3	100 ^c	39	61 ^c	1.56			
	HNO ₃ (<i>d</i> = 1.5)–H ₂ SO ₄ (<i>d</i> = 1.84)	—	RT	24	76 ^c	37	63 ^c	1.70			
	HNO ₃ –(CH ₃ CO) ₂ O–TFAA	Hβ	–10	2	100	13	87	6.69 ^d			
	NO ₂	—	RT	24	0	—	—	—			
	NO ₂ –O ₂	HBEA-25	RT	24	77	0	100	—			
	NO ₂ –O ₂	HSZ-940HOA	RT	24	95	9	91	10.1			
	NO ₂ –O ₂	HSZ-960HOA	RT	24	100	25	75	3.00			
	NO ₂ –O ₂	HSZ-980HOA	RT	24	100	12	88	7.33			
3-Methylbenzonitrile (<i>m</i> -tolunitrile) 13						4-Nitro 14	5-Nitro 15	6-Nitro 16	14 : 16	15 : 16	
	NO ₂ –O ₃	—	–10	3	100 ^c	14	63	23 ^c	0.61	2.74	
	HNO ₃ (<i>d</i> = 1.5)–H ₂ SO ₄ (<i>d</i> = 1.84)	—	RT	24	76 ^c	40	6	54 ^c	0.74	0.11	
	NO ₂	—	RT	24	0	—	—	—	—	—	
	NO ₂ –O ₂	HBEA-25	RT	24	23	8	Trace	92	0.09	—	
	NO ₂ –O ₂	HSZ-940HOA	RT	24	88	Trace	5	95	—	0.05	
	NO ₂ –O ₂	HSZ-960HOA	RT	24	100	Trace	6	94	—	0.06	
	NO ₂ –O ₂	HSZ-980HOA	RT	24	27	Trace	5	95	—	0.05	
4-Methylbenzonitrile (<i>p</i> -tolunitrile) 17						3-Nitro 18					
	NO ₂ –O ₃	—	–10	3	100 ^c	~100					
	HNO ₃ (<i>d</i> = 1.5)–H ₂ SO ₄ (<i>d</i> = 1.84)	—	RT	24	86 ^c	~100					
	NO ₂	—	RT	24	0	—					
	NO ₂ –O ₂	HBEA-150	RT	24	94	~100					
	NO ₂ –O ₂	Hβ-25	RT	24	64	~100					
	NO ₂ –O ₂	HSZ-960HOA	RT	24	46	~100					
	NO ₂ –O ₂	HSZ-980HOA	RT	24	Trace	~100					

^a See footnotes in Table 3. ^b See footnotes in Table 3. ^c Determined by ¹H-NMR. ^d Ref. 3c.

Table 5 Nitration of dimethylbenzonitriles with nitrogen dioxide and ozone or dioxygen in the presence of zeolites^a

Nitrile	Nitrating agent ^a	Catalyst	Temp./°C	Reaction time/h	Conversion (%) ^b	Isomer proportion (%) ^b			Remark
3,4-Dimethylbenzonitrile (3,4- <i>o</i> -xylonitrile) 19						2-Nitro 20	5-Nitro 21	6-Nitro 22	
	NO ₂ -O ₃	—	−10	1.5	89	31	49	20 ^c	
	HNO ₃ (<i>d</i> = 1.5)–H ₂ SO ₄ (<i>d</i> = 1.84)	—	RT	15	100	33	45	22 ^c	
	HNO ₃ –(MeCO) ₂ O	—	−10	24	41	14	28	9 ^d	
	NO ₂	—	RT	15	0	—	—	—	
	NO ₂ -O ₂	HSZ-930HOA	RT	15	3	—	—	100 ^c	
	NO ₂ -O ₂	HSZ-940HOA	RT	15	8	—	17	83 ^c	
	NO ₂ -O ₂	HSZ-960HOA	RT	15	100	—	—	100 ^c	
	NO ₂ -O ₂	HSZ-980HOA	RT	15	100	—	—	100 ^c	
2,4-Dimethylbenzonitrile (2,4- <i>m</i> -xylonitrile) 23						3-Nitro 24	5-Nitro 25		Isomer ratio 25 : 24
	NO ₂ -O ₃	—	−10	1.5	100	26	74		2.84
	NO ₂ -O ₃	— ^e	RT	1.5	100	18	82		4.56
	HNO ₃ (<i>d</i> = 1.5)/H ₂ SO ₄ (<i>d</i> = 1.84)	—	RT	15	100	30	70		2.33
	NO ₂	—	RT	15	—	—	—		—
	NO ₂ -O ₂	HSZ-930HOA	RT	15	95	4	96		24.0
	NO ₂ -O ₂	HSZ-940HOA	RT	15	83	7	93		13.3
	NO ₂ -O ₂	HSZ-960HOA	RT	15	99	3	97		24.3
	NO ₂ -O ₂	HSZ-980HOA	RT	15	66	6	94		15.7
2,5-Dimethylbenzonitrile (2,5- <i>p</i> -xylonitrile) 26						3-Nitro 27	4-Nitro 28	6-Nitro 29	Dinitration products
	NO ₂ -O ₃	—	−10	1.5	100	49	6	43	2
	NO ₂ -O ₃	— ^e	−10	1.5	100	32	16	48	4
	HNO ₃ (<i>d</i> = 1.5)–H ₂ SO ₄ (<i>d</i> = 1.84)	—	RT	15	100	34	7	59	0
	NO ₂	—	RT	15	3	69	31	—	0
	NO ₂ -O ₂	HSZ-930HOA	RT	15	96	17	39	44	0
	NO ₂ -O ₂	HSZ-940HOA	RT	15	94	11	44	45	0
	NO ₂ -O ₂	HSZ-960HOA	RT	15	83	29	27	44	0
	NO ₂ -O ₂	HSZ-980HOA	RT	15	97	19	18	63	Trace


^a See footnotes in Table 3. ^b Determined by ¹H-NMR. ^c Determined by GC. ^d Ref. 31, accompanied by 59% of addition products. ^e Dichloromethane was used as the solvent.

to favour the 4-nitro isomer **28** at the expense of the 6-nitro isomer **27**. The proportion of the 6-nitro isomer **29** was not much affected. In the presence of HSZ-960HOA, however, the isomer **27** became important at the expense of isomer **28**. In the absence of zeolites, compound **26** was quite slow to react. Interestingly, the 3-nitro isomer **27** became more important in this case and little or no 6-nitro isomer **29** was detected at this low conversion stage.

In the Kyodai-nitration of dimethylbenzonitriles, the product was always accompanied by small amounts of dinitration products. However, no such higher nitration products were observed in the zeolite-assisted nitration using nitrogen dioxide and dioxygen.

Kyodai-nitration of benzenedicarbonitriles

As expected, benzenedicarbonitriles were completely inert toward the combined action of nitrogen dioxide and dioxygen irrespective of the type of solid catalyst employed. However, with the intention of confirming the scope of the Kyodai-nitration, we have treated three isomeric benzenedicarbonitriles **30**, **33** and **35** with nitrogen dioxide in the presence of ozone. The reaction was quite slow and incomplete for all substrates even in the presence of a protonic acid or a Lewis acid as the catalyst, which included methanesulfonic acid, trifluoromethanesulfonic acid, and a variety of transition metal chlorides. However, a noticeable improvement was observed when a combination of protonic acid and metal chloride was employed as the catalyst. Thus, treatment of benzene-1,2-dicarbonitrile (phthalonitrile) **30** with nitrogen dioxide and ozone in the presence of methanesulfonic acid and iron(III) chloride at room temperature for 24 h gave a comparable mixture of 3- and 4-nitrobenzene-1,2-dicarbonitrile **31** and **32** (Scheme 5). The conversion was not satisfactory, but the yield based on it was almost quantitative (Table 6). Dinitriles **31** and **32** are important as the starting material for a variety of phthalocyanines with peripheral substituents on the macrocyclic ring. However, they are usually prepared by a multi-step procedure starting from nitrophthalides.^{22–24} Direct nitration of highly deactivated dinitrile **30** required heating with concentrated nitric and sulfuric acids at elevated temperatures, under which conditions the vicinal nitrile functions underwent extensive decomposition.



Scheme 5

5-Nitrobenzene-1,3-dicarbonitrile **34** is attractive as a starting material for iodine-based imaging agents. Thus, the Kyodai-nitration of benzene-1,3-dicarbonitrile (isophthalonitrile) **33** was investigated in some detail. Although a protonic acid or a

metal halide alone proved to be ineffective as the catalyst, the combined use of methanesulfonic acid and iron(III) chloride was found to give an acceptable result. Compound **36** was obtained as the single product and the yield based on conversion was nearly quantitative. Under similar conditions, benzene-1,4-dicarbonitrile **35** gave 2-nitro derivative **36**. The ease of three isomeric benzenedicarbonitriles toward the Kyodai-nitration followed the order **33** > **30** > **35**. Dinitrile **35** was the least reactive of all and the attempted nitration of **35** by heating with nitric and sulfuric acids only led to an intractable tarry substance.

In summary, moderately deactivated arenes such as 1-nitronaphthalene, naphthonitriles, and methylated benzonitriles are nitrated with comparative ease by a combination of nitrogen dioxide and molecular oxygen in the presence of zeolite. The present results well demonstrate the great potential of zeolites in controlling the isomer proportion of nitration products, usually quite difficult or infeasible by traditional nitration methodologies, by modifying the framework topology and distribution of the size and shape of pores in zeolites.

Experimental

General

All reagents except compounds **7**, **19**, **23** and **26** were reagent-grade commercial products. The above four nitriles were prepared according to the reported procedure.²⁵ Montmorillonite K10 and zeolite HZSM-5 were purchased from Aldrich and Acros, respectively. Zeolites HBEA-25, 150 and NaMFI-90 were a gift from the Catalyst Society of Japan. Zeolites HSZ-930, 940, 960 and 980HOA were a gift from the Research Institute of TOSOH Corporation, Ltd. The SiO₂–Al₂O₃ ratios of these products were 27.0, 37.1, 110 and 480, respectively.²⁶ Acetonitrile and dichloromethane were distilled from CaH₂ before use and stored over 4 Å molecular sieves under argon. Nitrogen dioxide (99% pure) was purchased from Sumitomo Seika Co., Ltd. in a cylinder and used after transfer distillation. Melting points were determined on a Yanaco MP S 3 apparatus and are uncorrected. Gas–liquid chromatographic analyses were carried out with a Shimadzu GC-17A instrument fitted with an AOC-17 auto injector and FID detector, using a DB-5 capillary column (30 m × 0.247 mm id) and a 25 QC2 BPX5 (25 m × 0.25 mm id) capillary column. Temperature programmed heating between 50 °C and 250 °C was employed. The injection port and detector were at a temperature of 250 °C. Cyclododecane was used as an internal standard to evaluate the isomer composition and the identity of products was proven by direct comparison with authentic samples or by GC-MS analyses (Shimadzu GCMS-QP 5000 using isobutane as an ionizing gas). ¹H-NMR spectra were determined in CDCl₃ on a JEOL JNM-A-400 NMR spectrometer using tetramethylsilane as an internal standard. IR measurements were made on a JEOL FTIR-5300 spectrophotometer for KBr pellets and only characteristic peaks were recorded. Microanalyses were performed at the Advanced Instrumentation Center, Ehime University. All product yields presented in Tables 2–6 were not optimized.

Zeolite-assisted aromatic nitration using nitrogen dioxide and dioxygen

Nitration of nitronaphthalene and naphthonitriles. Typical procedure. 1-Naphthonitrile **4** (140 mg, 0.9 mmol), liquid NO₂ (0.5 mL, 15 mmol), zeolite (700 mg), and acetonitrile (20 mL) were placed in a flask. Oxygen was passed into the system to replace air, and a balloon filled with oxygen was connected. The mixture was stirred for 12 h at room temperature and progress of the reaction was monitored intermittently by TLC on a plastic sheet coated with silica gel 60 F₂₅₄ (Merck). When the reaction was almost complete, excess NO₂ was expelled by blowing air into the solution and was collected in a cold trap for reuse.

Table 6 Catalyzed Kyodai-nitration of benzenedicarbonitriles^a

Dinitrile	NO ₂ /mL	O ₃ /mmol	Catalyst	Conversion (%) ^b	Yield (%) ^c	
Benzene-1,2-dicarbonitrile (Phthalonitrile) 30	3.0	40	CH ₃ SO ₃ H/FeCl ₃	35	3-Nitro 31	4-Nitro 32
					50	50
Benzene-1,3-dicarbonitrile (isophthalonitrile) 33					5-Nitro 34	
	1.5	20	CH ₃ SO ₃ H	2	~100	
	1.5	20	FeCl ₃	0	~100	
	1.5	20	CH ₃ SO ₃ H–FeCl ₃	25	~100	
	1.5	20	CF ₃ SO ₃ H–FeCl ₃	18	~100	
	1.5	20	CF ₃ SO ₃ H–AlCl ₃	12	~100	
	1.5	20	CF ₃ SO ₃ H–ZnCl ₂	18	~100	
	1.5	20	CF ₃ SO ₃ H–BiCl ₃	11	~100	
	3.0	40	CH ₃ SO ₃ H–FeCl ₃	45	~100	
Benzene-1,4-dicarbonitrile (Terephthalonitrile) 35					2-Nitro 36	
	3.0	40	CH ₃ SO ₃ H–FeCl ₃	23	~100	

^a For detail, see the Experimental section. ^b Determined by ¹H-NMR. ^c Yield based on conversion.

The reaction mixture was evaporated under reduced pressure and the residue was extracted with ethyl acetate. The combined extracts were washed with brine, dried over MgSO₄, and evaporated to leave a solid residue, the composition of which was analyzed by GC and ¹H-NMR. For the isolation of each isomeric product, the residue was chromatographed on silica gel using hexane–EtOAc as the solvent to elute 5-nitro-1-naphthonitrile **5** and 8-nitro-1-naphthonitrile **6** in this order. A single crystallization of each fraction from ethanol gave **5** as light yellow crystals (58 mg, 32%) and **6** as red crystals (40 mg, 22%). The recovered zeolite was reused four times without appreciable loss of catalytic activity.

5-Nitro-1-naphthonitrile (5). Mp 205–207 °C (lit. 205²⁷ and 211–212 °C^{14b}); ν_{\max} (KBr)/cm^{−1} 2226 (CN), 1523 (NO₂), 1334 (NO₂), 794; δ_{H} (CDCl₃) 8.83 (d, 1H, *J* = 8.7), 8.59 (d, 1H, *J* = 8.1), 8.37 (dd, 1H, *J* = 7.8, 0.9), 8.09 (dd, 1H, *J* = 7.2, 0.9), 7.83 (t, 1H, *J* = 4.2), 7.79 (t, 1H, *J* = 7.2); *m/z* 198 (54), 152 (63), 140 (100), 125 (43).

8-Nitro-1-naphthonitrile (6). Mp 151–153 °C (lit.^{14b} 153–154 °C); ν_{\max} (KBr)/cm^{−1} 2226 (CN), 1523 (NO₂), 1536, 1338 (NO₂), 831, 756; δ_{H} (CDCl₃) 8.23 (dd, 1H, *J* = 8.4, 0.9), 8.16 (dd, 1H, *J* = 8.7, 1.5), 8.15 (dd, 1H, *J* = 7.2, 1.5), 8.02 (dd, 1H, *J* = 7.5, 1.2), 7.71 (m, 2H); *m/z* 198 (63), 168 (28), 152 (100), 140 (85), 125 (51).

5-Nitro-2-naphthonitrile (8). Mp 75–76 °C; (Found: C, 66.32; H, 2.96; N, 14.37. C₁₁H₆N₂O₂ requires C, 66.67; H, 3.05; N, 14.14%); ν_{\max} (KBr)/cm^{−1} 2227 (CN), 1521 (NO₂), 1340 (NO₂), 802; δ_{H} (CDCl₃) 8.73 (d, 1H, *J* = 8.8), 8.42 (dd, 1H, *J* = 8.0, 1.2), 8.37 (d, 1H, *J* = 1.2), 8.21 (d, 1H, *J* = 8.4), 7.87 (dd, 1H, *J* = 9.2, 1.6), 7.73 (t, 1H, *J* = 8.0); *m/z* 198 (65), 168 (54), 152 (65), 140 (100), 125 (40).

5-Nitro-2-naphthonitrile (9). Mp 66–67 °C; (Found: C, 66.69; H, 3.20; N, 14.25. C₁₁H₆N₂O₂ requires C, 66.67; H, 3.05; N, 14.14%); ν_{\max} (KBr)/cm^{−1} 2229 (CN), 1523 (NO₂), 1338 (NO₂), 842; δ_{H} (CDCl₃) 9.01 (s, 1H), 8.42 (dd, 1H, *J* = 8.0, 1.2), 8.21 (d, 1H, *J* = 8.4), 8.08 (d, 1H, *J* = 8.4), 7.77 (m, 2H); *m/z* 198 (78), 168 (25), 152 (95), 140 (100), 125 (58).

Nitration of methylated benzonitriles. General procedure. Methylbenzonitrile (30 mg, 0.3 mmol) or dimethylbenzonitrile (40 mg; 0.3 mmol), liquid NO₂ (0.5 mL, 15 mmol), zeolite (400 mg), and acetonitrile (20 mL) were placed in a flask. Oxygen was passed into the flask to replace air and a balloon containing oxygen was connected. The mixture was magnetically stirred for a given time at room temperature, while progress of the reaction was intermittently monitored by TLC. When the reaction was almost complete, unchanged NO₂ was expelled by blowing air into the solution and was collected in a cold trap for

reuse. The reaction mixture was evaporated under reduced pressure and the residue was extracted with ethyl acetate. The combined extracts were washed with brine, dried over MgSO₄, and evaporated. The product composition was analysed using GC and ¹H-NMR. For the isolation of each isomeric product, the residue was chromatographed on silica gel using hexane–EtOAc as the solvent.

2-Methyl-3-nitrobenzonitrile (11). Mp 68–69 °C (lit. 63–65^{28a} and 69.5 °C^{28b}); ν_{\max} (KBr)/cm^{−1} 2234 (CN), 1532 (NO₂), 1352 (NO₂); δ_{H} (CDCl₃) 8.12 (dd, 1H, *J* = 1.2, 8.2), 7.87 (dd, 1H, *J* = 1.2, 8.0), 7.50 (t, 1H, *J* = 8.2), 2.79 (s, 3H); *m/z* 162 (17), 146 (10), 145 (100), 118 (10), 117 (61), 116 (22).

2-Methyl-5-nitrobenzonitrile (12). Mp 106–107 °C (lit.^{29a} 104–106 °C); ν_{\max} (KBr)/cm^{−1} 2236 (CN), 1524 (NO₂), 1352 (NO₂); δ_{H} (CDCl₃) 8.48 (d, 1H, *J* = 2.8), 8.34 (dd, 1H, *J* = 2.8, 8.6), 7.54 (dd, 1H, *J* = 0.8, 8.6), 2.69 (s, 3H); *m/z* 162 (100), 132 (34), 117 (29), 116 (96), 114 (17), 104 (28).

3-Methyl-4-nitrobenzonitrile (14). Mp 80–81 °C (lit.^{29b} 80 °C); ν_{\max} (KBr)/cm^{−1} 2236 (CN), 1524 (NO₂), 1350 (NO₂); δ_{H} (CDCl₃) 8.02 (d, 1H, *J* = 8.0), 7.67 (m, 2H), 2.63 (s, 3H); *m/z* 162 (13), 146 (11), 145 (100), 118 (10), 117 (53), 116 (23).

3-Methyl-5-nitrobenzonitrile (15). Mp 100–102 °C (lit.^{16c} 104–105 °C); ν_{\max} (KBr)/cm^{−1} 2240 (CN), 1534 (NO₂), 1360 (NO₂); δ_{H} (CDCl₃) 8.34 (s, 1H), 8.28 (s, 1H), 7.79 (s, 1H), 2.55 (s, 3H); *m/z* 162 (77), 131 (11), 126 (10), 117 (16), 116 (100), 104 (24).

3-Methyl-6-nitrobenzonitrile (16). Mp 92–93 °C (lit.³⁰ 93–94 °C); ν_{\max} (KBr)/cm^{−1} 2236 (CN), 1524 (NO₂), 1350 (NO₂); δ_{H} (CDCl₃) 8.23 (d, 1H, *J* = 8.4), 7.70 (s, 1H), 7.59 (d, 1H, *J* = 8.4), 2.54 (s, 3H); *m/z* 162 (100), 132 (43), 116 (35), 104 (27).

4-Methyl-3-nitrobenzonitrile (18). Mp 106–107 °C (lit.³¹ 107 °C); ν_{\max} (KBr)/cm^{−1} 2234 (CN), 1526 (NO₂), 1352 (NO₂); δ_{H} (CDCl₃) 8.28 (d, 1H, *J* = 1.6), 7.87 (dd, 1H, *J* = 1.6, 7.8), 7.51 (d, 1H, *J* = 7.8), 2.69 (s, 3H); *m/z* 162 (21), 145 (100), 118 (15), 117 (57), 116 (22), 115 (10), 114 (11).

3,4-Dimethylbenzonitrile (19). Mp 67–68 °C (lit.²⁵ 65–66 °C); ν_{\max} (KBr)/cm^{−1} 2224 (CN); δ_{H} (CDCl₃) 7.41 (s, 1H), 7.39 (d, 1H, *J* = 7.5), 7.21 (d, 1H, *J* = 7.5), 2.32 (s, 3H), 2.29 (s, 3H).

3,4-Dimethyl-2-nitrobenzonitrile (20). Mp 79–80 °C (lit.³² 72–75 °C); ν_{\max} (KBr)/cm^{−1} 2237 (CN), 1529 (NO₂), 1359 (NO₂); δ_{H} (CDCl₃) 7.42 (d, 1H, *J* = 8.8), 7.55 (d, 1H, *J* = 8.8), 2.44 (s, 3H), 2.30 (s, 3H).

3,4-Dimethyl-5-nitrobenzonitrile (21). Mp 119–120 °C (lit.³² 116–118 °C); ν_{\max} (KBr)/cm^{−1} 2241 (CN), 1529 (NO₂), 1367 (NO₂); δ_{H} (CDCl₃) 7.91 (s, 1H), 7.65 (s, 1H), 2.46 (s, 3H), 2.40 (s, 3H).

3,4-Dimethyl-6-nitrobenzonitrile (22). Mp 170–171 °C (lit.³³ 179–181 °C); ν_{\max} (KBr)/cm^{−1} 2237 (CN), 1529 (NO₂), 1359

(NO₂); δ_{H} (CDCl₃) 8.11 (s, 1H), 7.64 (s, 1H), 2.45 (s, 3H), 2.42 (s, 3H).

2,4-Dimethylbenzonitrile (23). Oil.²⁵ ν_{max} (KBr)/cm⁻¹ 2220 (CN); δ_{H} (CDCl₃) 7.48 (d, 1H, $J = 7.9$), 7.12 (s, 1H), 7.07 (d, 1H, $J = 7.9$), 2.50 (s, 3H), 2.37 (s, 3H).

2,4-Dimethyl-3-nitrobenzonitrile (24). Mp 122–123 °C; (Found: C, 61.32; H, 4.75; N, 15.92. C₉H₈N₂O₂ requires C, 61.36; H, 4.58; N, 15.90%); ν_{max} (KBr)/cm⁻¹ 2230 (CN), 1527 (NO₂), 1373 (NO₂); δ_{H} (CDCl₃) 7.45 (d, 1H, $J = 7.9$), 7.41 (d, 1H, $J = 7.9$), 2.61 (s, 3H), 2.41 (s, 3H).

2,4-Dimethyl-5-nitrobenzonitrile (25). Mp 109–110 °C (lit.²¹ 108 °C); ν_{max} (KBr)/cm⁻¹ 2230 (CN), 1521 (NO₂), 1342 (NO₂); δ_{H} (CDCl₃) 7.87 (s, 1H), 7.60 (s, 1H), 2.60 (s, 3H), 2.57 (s, 3H).

2,5-Dimethylbenzonitrile (26). Oil.²⁵ ν_{max} (KBr)/cm⁻¹ 2236 (CN), 1524 (NO₂), 1352 (NO₂); δ_{H} (CDCl₃) 7.39 (s, 1H), 7.28 (d, 1H, $J = 8.3$), 7.19 (d, 1H, $J = 8.3$), 2.49 (s, 3H), 2.33 (s, 3H).

2,5-Dimethyl-6-nitrobenzonitrile (27). Mp 76–77 °C (lit.^{34a} 85 °C); ν_{max} (KBr)/cm⁻¹ 2234 (CN), 1528 (NO₂), 1356 (NO₂); δ_{H} (CDCl₃) 7.45 (d, 1H, $J = 8.3$), 7.41 (d, 1H, $J = 8.3$), 2.60 (s, 3H), 2.41 (s, 3H).

2,5-Dimethyl-4-nitrobenzonitrile (28). Mp 117–118 °C;^{34b} ν_{max} (KBr)/cm⁻¹ 2235 (CN), 1531 (NO₂), 1358 (NO₂); δ_{H} (CDCl₃) 7.87 (s, 1H), 7.60 (s, 1H), 2.60 (s, 3H), 2.57 (s, 3H).

2,5-Dimethyl-3-nitrobenzonitrile (29). Mp 117–118 °C (Found: C, 60.96; H, 4.89; N, 15.71. C₉H₈N₂O₂ requires C, 61.36; H, 4.58; N, 15.90%); ν_{max} (KBr)/cm⁻¹ 2235 (CN), 1533 (NO₂), 1350 (NO₂); δ_{H} (CDCl₃) 7.92 (s, 1H), 7.67 (s, 1H), 2.72 (s, 3H), 2.49 (s, 3H).

Kyodai-nitration of benzenedicarbonitriles. Typical procedure. Benzene-1,3-dicarbonitrile **33** (60 mg, 0.5 mmol), liquid NO₂ (3.0 mL, 90 mmol), methanesulfonic acid (52 mg), FeCl₃ (2.6 mg), and acetonitrile (30 mL) were placed in a two-necked flask fitted with an inlet tube and a vent which permits waste gas to escape. The mixture was cooled to –10 °C in an ice–salt bath, while a stream of ozonized oxygen was introduced under vigorous stirring through the gas inlet tube, which should dip just below the surface of the liquid in the flask. The progress of reaction was intermittently monitored by TLC. After 4 h, the cooling bath was removed, and the mixture was further stirred for 24 h at room temperature. Then, dry air was bubbled into the mixture to expel and recover unchanged nitrogen dioxide by cold trap. The mixture was evaporated under reduced pressure to leave a solid residue, which was extracted three times with dichloromethane. The combined extracts were washed with brine, dried, and evaporated. Chromatography of the resulting solid on silica gel using hexane–EtOAc as the solvent, followed by recrystallization from ethanol gave 5-nitrobenzene-1,3-dicarbonitrile **34** as colourless needles (35 mg, 41%). Unchanged dinitrile **33** was recovered in 55% yield.

3-Nitrobenzene-1,2-dicarbonitrile (31). Mp 165–166 °C (lit.¹⁴¹ 141³⁵ and 160 °C²²); ν_{max} (KBr)/cm⁻¹ 2240 (CN), 1541 (NO₂), 1354 (NO₂); δ_{H} (CDCl₃) 8.57 (dd, 1H, $J = 4.4, 1.2$), 8.15 (dd, 1H, $J = 8.0, 0.8$), 7.99 (t, 1H, $J = 8.4$); m/z 173 (84), 143 (84), 127 (68), 116 (49), 101 (51), 100 (100).

4-Nitrobenzene-1,2-dicarbonitrile (32). Mp 145–146 °C (lit.²³ 141 °C); ν_{max} (KBr)/cm⁻¹ 2240 (CN), 1535 (NO₂), 1356 (NO₂); δ_{H} (CDCl₃) 8.69 (dd, 1H, $J = 0.4, 2.0$), 8.59 (dd, 1H, $J = 8.4, 2.0$), 8.07 (dd, 1H, $J = 0.4, 8.4$); m/z 173 (65), 127 (100), 101 (12).

5-Nitrobenzene-1,3-dicarbonitrile (34). Mp 214–215 °C (lit.^{209–210} 36^a and 204–205 °C^{36b}); ν_{max} (KBr)/cm⁻¹ 2245 (CN), 1543 (NO₂), 1385, 1354 (NO₂); δ_{H} (CDCl₃) 8.75 (d, 2H, $J = 1.5$), 8.27 (t, 1H, $J = 1.5$); m/z 173 (64), 128 (11), 127 (100), 101 (10), 100 (77).

2-Nitrobenzene-1,4-dicarbonitrile (36). Mp 124–125 °C (Found: C, 55.59; H, 2.16; N, 24.33. C₉H₆N₂O₂ requires C, 55.50; H, 1.75; N, 24.27%); ν_{max} (KBr)/cm⁻¹ 2240 (CN), 1543 (NO₂), 1356 (NO₂); δ_{H} (CDCl₃) 8.63 (d, 1H, $J = 1.2$), 8.10 (m, 2H); m/z 173 (53), 127 (74), 100 (100).

Acknowledgements

This work was supported by a Grant-in-Aid (no. 12640576) from the Ministry of Education, Culture, Sports, Science, and Technology, Japan. The authors acknowledge the Catalytic Society of Japan and Dr M. Nakano of TOSOH Corporation, Ltd. for the generous gifts of zeolite samples. X. P. thanks the Japan Society for the Promotion of Science for a Fellowship (no. P99282).

References

- For a general survey of aromatic nitration, see: (a) L. F. Albright, R. V. C. Carr and R. J. Schmitt, eds. *Nitration*, ACS Symposium Series 623, American Chemical Society, Washington DC, 1996; (b) G. A. Olah, R. Malhotra and S. C. Narang, *Nitration: Methods and Mechanisms*, VCH Publishers Inc., New York, 1989; (c) K. Schofield, *Aromatic Nitration*, Cambridge University Press, London, 1980.
- H. Suzuki, S. Yonezawa, N. Nonoyama and T. Mori, *J. Chem. Soc., Perkin Trans. 1*, 1996, 2385.
- (a) K. Smith, S. Almeer and C. Peters, *Chem. Commun.*, 2001, 2748; (b) K. Smith, S. Almeer and S. J. Black, *Chem. Commun.*, 2000, 1571; (c) K. Smith, T. Gibbins, R. W. Millar and R. P. Claridge, *J. Chem. Soc., Perkin Trans. 1*, 2000, 2753.
- X. Peng and H. Suzuki, *Org. Lett.*, 2001, 3, 3431.
- E. R. Ward and J. G. Hawkins, *J. Chem. Soc.*, 1954, 2975.
- E. R. Ward, C. D. Johnson and L. A. Day, *J. Chem. Soc.*, 1959, 487.
- H. E. Fierz-David and R. Sponagel, *Helv. Chem. Acta*, 1943, **26**, 98.
- H. H. Hodgson and J. S. Whitehurst, *J. Chem. Soc.*, 1945, 202.
- An early claim¹⁰ that the reaction of naphthalene with excess NO₂–BF₃ in nitromethane at 0 °C led to a 5 : 3 mixture of **2** and **3** was later rectified; E. R. Ward and C. D. Johnson, *J. Chem. Soc.*, 1961, 4314.
- G. B. Bachman, H. Feuer, B. R. Bleustein and C. M. Vogt, *J. Am. Chem. Soc.*, 1955, **77**, 6188.
- N. Donaldson, *The Chemistry and Technology of Naphthalene Compounds*, Edward Arnold, London, 1958, p. 151.
- B. Gigante, A. O. Prezeres, M. J. Marcelo-Curto, A. Cornelis and P. Laszlo, *J. Org. Chem.*, 1995, **60**, 3449.
- (a) For a survey of the Kyodai-nitration, see: N. Nonoyama, T. Mori and H. Suzuki, *Zh. Org. Khim.*, 1998, **34**, 1591 (*Russ. J. Org. Chem.*, 1998, **34**, 1521); (b) T. Suzuki and R. Noyori, *Chemtracts*, 1997, **10**, 813; (c) T. Mori and H. Suzuki, *Synlett*, 1995, 383.
- (a) S. Paraskar, H. S. Jagtap and A. Sudalai, *J. Chem. Res. (S)*, 2000, 39; (b) M. Makosza, W. Danikiewicz and K. Wojciechowski, *Liebigs Ann. Chem.*, 1987, 711.
- (a) A. Rosowsky, R. A. Forsch, A. Null and R. G. Moran, *J. Med. Chem.*, 1999, **42**, 3510; (b) D. L. Boger, M. J. Kochanny and H. Cai et al., *Bioorg. Med. Chem.*, 1998, **6**, 643; (c) P. V. Divekar and L. C. Vining, *Can. J. Chem.*, 1964, **42**, 63.
- (a) E. S. Adams and K. L. Rinehart, *J. Antibiot.*, 1994, **47**, 1456; (b) T. H. Fischer, S. M. Dershem and M. L. Prewitt, *J. Org. Chem.*, 1990, **55**, 1040; (c) M. H. Gelb and R. H. Abeles, *J. Med. Chem.*, 1986, **29**, 585.
- (a) A. J. Bloom, M. Fleischmann and J. M. Mellor, *Electrochim. Acta*, 1978, **32**, 785; (b) S. J. Kuhn and G. A. Olah, *J. Am. Chem. Soc.*, 1961, **83**, 4564.
- I. I. Schuster, *Magn. Reson. Chem.*, 1996, **34**, 301.
- H. Suzuki, J. Tomaru and T. Mori, *J. Chem. Soc., Perkin Trans. 1*, 1994, 2413.
- (a) I. A. McCulloch, *Macromolecules*, 1994, **27**, 1697; (b) C. Bloomsfield, R. B. Moodie and K. Schofield, *J. Chem. Soc., Perkin Trans. 2*, 1983, 1003; (c) W. O. Kermack, *J. Chem. Soc.*, 1924, **125**, 2285.
- L. Chardonens and W. J. Kramer, *J. Am. Chem. Soc.*, 1957, **79**, 4955.
- R. D. George and A. W. Snow, *J. Heterocycl. Chem.*, 1995, **32**, 495.
- J. G. Young and W. Onyebugu, *J. Org. Chem.*, 1990, **55**, 2155.
- J. Griffiths and B. Roozpeikar, *J. Chem. Soc., Perkin Trans. 1*, 1976, 42.
- L. Friedman and H. Schechter, *J. Org. Chem.*, 1961, **26**, 2522.
- Detailed information is available from Dr Masao Nakano, Research Institute, TOSOH Corporation, Ltd., 4560 Kaisei-cho, Shinnanyo 746–8501, Japan.
- Elseviers Encyclopedia of Organic Chemistry*, vol. 12B, Elsevier, Amsterdam, 1953, 4164, taken from ref. 14b.
- (a) G. S. Ponticello and J. J. Baldwin, *J. Org. Chem.*, 1979, **44**, 4003; (b) E. Sorkin, W. Kraehenbuehl and H. Erlenmeyer, *Helv. Chim. Acta*, 1948, **31**, 65.

- 29 (a) *Dictionary of Organic Compounds*, ed. J. Buckingham, 5th ed., vol. 4, Chapman and Hall, New York, 1982, p. 3937; (b) *Dictionary of Organic Compounds*, ed. J. Buckingham, 5th ed., vol. 4, Chapman and Hall, New York, 1982, p. 3938.
- 30 T. H. Fischer and A. W. Meierhoefer, *J. Org. Chem.*, 1978, **43**, 220.
- 31 R. L. Chan and T. C. Bruice, *J. Am. Chem. Soc.*, 1977, **99**, 6721.
- 32 A. Fischer and C. C. Greig, *Can. J. Chem.*, 1974, **52**, 1231.
- 33 A. Brandstrom and S. A. I. Carlsson, *Acta Chem. Scand.*, 1967, **21**, 983.
- 34 T. R. Govindachari, S. Rajappa and V. Sudarsanam, *Indian J. Chem. Sect. B*, 1963, **1**, 247.
- 35 M. Scalera and R. E. Brouillard, US Patent, 2 525 620/1950, *Chem. Abstr.*, 1951, **45**, 2023c.
- 36 (a) J. H. Fendler, N. L. Arthur and C. E. Griffin, *J. Org. Chem.*, 1972, **37**, 812; (b) A. E. Senear, M. M. Rapport, J. F. Mead, J. T. Maynard and J. B. Koepfli, *J. Org. Chem.*, 1946, **11**, 378.