

Electron-Transfer Photoredox Catalysis: Development of a Tin-Free Reductive Dehalogenation Reaction

Jagan M. R. Narayanam, Joseph W. Tucker, and Corey R. J. Stephenson*


Department of Chemistry, Boston University, Boston, Massachusetts 02215

Received April 25, 2009; E-mail: crjsteph@bu.edu


The use of visible light as a promoter in synthesis is appealing because of its natural abundance, ease of use, and potential application on an industrial scale.¹ However, the lack of visible-light absorption by many organic molecules has limited the number of applications documented to date. A solution to this problem involves the use of photosensitive catalysts.² Recent efforts from the MacMillan^{3a} and Yoon^{3b} laboratories have focused on the use of the well-known photoredox catalyst tris(2,2'-bipyridyl)ruthenium(II) chloride, Ru(bpy)₃Cl₂,⁴ to promote the enantioselective alkylation of aldehydes and [2 + 2] cycloaddition reactions, respectively. In this paper, we describe an environmentally benign,⁵ tin-free reductive dehalogenation⁶ system that is based on the combination of Ru(bpy)₃Cl₂ and an amine as the hydrogen atom source. In addition, this photoredox system gives superior chemoselectivity compared with traditional methodologies based on the chemistry of radicals. Furthermore, photoredox catalysis is presented as a potential means of accessing traditional radical chemistry while avoiding the toxicity and purification problems associated with tin hydrides (Scheme 1).^{7–9}

Scheme 1. Photoredox Catalytic Reduction and Potential C–C Bond Formation

Commonly Used Methodology for Radical Generation Using Trialkyltin Hydride



Tin-Free Alternative Using Ru(II) Photo-Redox Catalysis



Our preliminary studies focused on the reductive debromination of bromopyrrolindoline **1a**.^{10,11} Initial success was obtained upon subjecting **1a** to irradiation by a 14 W fluorescent lamp¹² with Ru(bpy)₃Cl₂ (5 mol %) as a catalyst in the presence of ⁱPr₂NEt (2 equiv) in DMF for 24 h.¹³ These conditions provided the reduced product in 25% yield (40% conversion). Subsequent optimization revealed two sets of conditions of broad utility for reductive debromination: Ru(bpy)₃Cl₂ (2.5 mol %) in DMF with (a) ⁱPr₂NEt (10 equiv) and formic acid (10 equiv) or (b) ⁱPr₂NEt (2 equiv) and Hantzsch ester (**3**) (1.1 equiv). In the case of **1a**, both systems cleanly produced the debrominated product in >90% yield. In addition, lower catalyst loading had little effect on the reaction: catalyst loadings as low as 0.05 mol % gave efficient reductive dehalogenation without prolonging the reaction time (Table 1).


Table 1. Optimization of the Reductive Debromination of **1a**

entry	conditions	yield (%) ^a
1	ⁱ Pr ₂ NEt (2 equiv), HCOOH (2 equiv), Ru ²⁺ (5 mol %), 24 h	25
2	ⁱ Pr ₂ NEt (10 equiv), HCOOH (10 equiv), Ru ²⁺ (2.5 mol %), 4 h	90
3	Et ₃ N (10 equiv), HCOOH (10 equiv), Ru ²⁺ (5 mol %), 24 h	20 ^b
4	ⁱ Pr ₂ NEt (2 equiv), 3 (1.1 equiv), Ru ²⁺ (2.5 mol %), 4 h	95
5	ⁱ Pr ₂ NEt (10 equiv), HCOOH (10 equiv), Ru ²⁺ (1 mol %), 4 h	90
6	ⁱ Pr ₂ NEt (10 equiv), HCOOH (10 equiv), Ru ²⁺ (0.05 mol %), 4 h	75 ^c

^a Isolated yield after purification by chromatography on SiO₂. ^b 25% conversion. ^c Reaction conducted on a 2.0 g (4.6 mmol) scale.

The scope of this process is illustrated by the examples detailed in Table 2. Utilizing our optimized conditions, we were able to successfully reduce both C–Br and C–Cl bonds in the presence of other functional groups such as free hydroxyls (entries 4 and 6), olefins (entry 9), and alkynes (entry 10). Furthermore, we found excellent selectivity for the reductive dehalogenation of alkyl bromides and chlorides α to electron-withdrawing groups over vinyl iodides (entry 9) and aryl bromides and iodides (entries 2, 3, 7, and 8). These conditions were also tolerant of protecting groups, including silyl ethers (entry 5) and carbamates (entries 1–3). In the case of less reactive chlorinated compounds, S_N2 substitution by the formate ion was competitive with the dehalogenation reaction. For these substrates, the simple substitution of **3** in place of formic acid as the hydrogen atom source provided a complementary set of reaction conditions for the α-chlorocarbonyl compounds (entries 7–10). Furthermore, reduction of simple alkyl halides was not observed.¹⁴

Throughout our investigations, we conducted a series of control experiments using **1a** as a representative substrate. Exclusion of any of the reaction components, including visible light, Ru(bpy)₃Cl₂, ⁱPr₂NEt, or combinations thereof, afforded only trace amounts of the reduced product **2a** after 48 h. Furthermore, to add credence to the intermediacy of a radical, we utilized an α-bromo-α-cyclopropyl imide substrate as a radical clock (eq 1).¹⁵ The corresponding



cyclopropyl ring fragmentation product was obtained in 52% isolated yield. This result strongly suggests that the reaction proceeds via a radical mechanism.


To further support our mechanistic hypothesis, we performed labeling studies using either DCO₂D or DCO₂H (98% D, 10 equiv) with ⁱPr₂NEt (10 equiv) and Ru(bpy)₃Cl₂ (2.5 mol %) in DMF for the reduction of **1a** (eq 2).¹⁶ Both afforded a **2a_H**/**2a_D** product ratio

Table 2. Reductive Dehalogenation Using Photoredox Catalysis

entry	substrate	product	yield ^a
1			95 ^b
2			92 ^b
3			91 ^b
4			99 ^b
5			79 ^b
6			80 ^b
7			78 ^c
8			88 ^c
9			81 ^c
10			89 ^c

^a Isolated yield after purification by chromatography on SiO₂.


^b Ru(bpy)₃Cl₂·6H₂O (2.5 mol %), ⁱPr₂NEt (10 equiv), HCO₂H (10 equiv), DMF, *hν*, r.t., 4 h. ^c Ru(bpy)₃Cl₂·6H₂O (2.5 mol %), ⁱPr₂NEt (2 equiv), **3** (1.1 equiv), DMF, *hν*, r.t., 4–24 h.



of ~4:1 (¹H NMR), indicating that ⁱPr₂NEt likely is the major hydrogen atom source in the reduction. In addition, no deuterium incorporation was observed in **2a** when the reduction reaction was conducted in DMF-*d*₇.

We also performed a reaction using formic acid (10 equiv), Et₃N-*d*₁₅ (10 equiv), and Ru(bpy)₃Cl₂ (2.5 mol %) in DMF, which led to only 25% conversion of starting material over 24 h and no incorporation of deuterium into the reduced product **2a**. This suggests that in the absence of ⁱPr₂NEt, the formate ion can successfully compete with the triethylammonium radical cation as the major hydrogen atom source (Scheme 2, Path A).

Accordingly, we propose a mechanism wherein single-electron transfer from the ammonium formate complex to the excited Ru(II)* occurs, forming Ru(I) and the radical cation of the complex. Reduction of the carbon–halogen bond by the electron-rich Ru(I) forms the alkyl radical, which can be reduced by abstraction of a hydrogen atom from one of the methine carbons of the radical cation of ⁱPr₂NEt (Scheme 2, Path B). The proposed hydrogen abstraction from the radical cation of ⁱPr₂NEt also accounts for the reductive dehalogenation of **1a** by 3° amines alone in the presence of Ru(bpy)₃Cl₂.

Scheme 2. Plausible Mechanism for the Reductive Dehalogenation

In conclusion, we have developed a catalytic, tin-free method for reductive dehalogenation utilizing visible-light photocatalysis. The reaction is highlighted by its chemoselectivity, affording the reduced compounds in high yields under mild reaction conditions. Further development and application of photoredox catalysis in the context of radical chemistry and its application to C–C bond-forming reactions are currently underway.

Acknowledgment. This work was supported by startup funds from Boston University. J.M.R.N. thanks the Swiss National Science Foundation for a postdoctoral fellowship. The authors thank Cambridge Isotope Laboratories for a gift of Et₃N-*d*₁₅.

Supporting Information Available: Experimental procedures and ¹H and ¹³C NMR spectra for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) For a recent review of chemistry using visible light, see: Sala, X.; Romero, I.; Rodríguez, M.; Lluís, E.; Llobet, A. *Angew. Chem., Int. Ed.* **2009**, *48*, 2842.
- (2) For selected early examples of photoactivation of metal catalysts, see: (a) Srinivasan, R. *J. Am. Chem. Soc.* **1963**, *85*, 3048. (b) Baldwin, J. E.; Greeley, R. H. *J. Am. Chem. Soc.* **1965**, *87*, 4514. (c) Salomon, R. G.; Kochi, J. K. *J. Am. Chem. Soc.* **1974**, *96*, 1137.
- (3) (a) Nicewicz, D. A.; MacMillan, D. W. C. *Science* **2008**, *322*, 77. (b) Ischay, M. A.; Anzovino, M. E.; Du, J.; Yoon, T. P. *J. Am. Chem. Soc.* **2008**, *130*, 12886.
- (4) Juris, A.; Balzani, V.; Barigelletti, F.; Campagna, S.; Belser, P.; Von Zelewsky, V. *Coord. Chem. Rev.* **1988**, *84*, 85.
- (5) For reviews of sustainable chemistry, see: (a) Albini, A.; Fagnoni, M. *Green Chem.* **2004**, *6*, 1. (b) Jenck, J. F.; Agterberg, F.; Droscher, M. *J. Green Chem.* **2004**, *6*, 544.
- (6) For a review of reductive dehalogenation, see: Alonso, F.; Beletskaya, I. P.; Yus, M. *Chem. Rev.* **2002**, *102*, 4009.
- (7) For the use of dihydroacridines as a hydrogen atom source for the reduction of α-haloacetophenones using photoredox catalysis, see: Fukuzumi, S.; Mochizuki, S.; Tanaka, T. *J. Phys. Chem.* **1990**, *94*, 722.
- (8) For selected recent examples of Sn-free radical reactions, including dehalogenations, see: (a) Gong, H.; Andrews, R. S.; Zuccarello, J. L.; Lee, S. J.; Gagné, M. R. *Org. Lett.* **2009**, *11*, 879. (b) Spiegel, D. A.; Wiberg, K. B.; Schacherer, L. N.; Medeiros, M. R.; Wood, J. L. *J. Am. Chem. Soc.* **2005**, *127*, 12513. (c) Hayashi, N.; Shibata, I.; Baba, A. *Org. Lett.* **2005**, *7*, 3093.
- (9) Ryu, I.; Noboru, S.; Curran, D. P. *Chem. Rev.* **1996**, *96*, 177.
- (10) It has been previously demonstrated that cation formation at the benzylic position of these substrates is disfavored. See: Bruncko, M.; Crich, D.; Samy, R. *J. Org. Chem.* **1994**, *59*, 5543.
- (11) For a recent example of Co-mediated radical dimerization for the preparation of bispyrrolidinoindoline alkaloids, see: Kim, J.; Ashenhurst, J. A.; Movassaghi, M. *Science* **2009**, *324*, 238.
- (12) The photochemistry setup can be purchased from retail stores for <\$15. See the Supporting Information for further details.
- (13) Trialkylamines have been found to disproportionate to the corresponding trialkylammonium salts and enamines upon visible-light irradiation in the presence of Ru(bpy)₃Cl₂. See: DeLaive, P. J.; Sullivan, B. P.; Meyer, T. J.; Whitten, D. G. *J. Am. Chem. Soc.* **1979**, *101*, 4007.
- (14) Examination of other Ru catalysts with larger reduction potentials to address the reduction of unactivated halides is currently underway.
- (15) (a) Newcomb, M. *Tetrahedron* **1993**, *49*, 1151. (b) Griller, D.; Ingold, K. U. *Acc. Chem. Res.* **1980**, *13*, 317.
- (16) See the Supporting Information for details.

JA9033582