

Effective One-pot Synthesis of H type 1 and 2 Trisaccharide Derivatives Using Glycal Epoxide


Hiroshi Tanaka, Nobuatsu Matoba, and Takashi Takahashi*

Department of Applied Chemistry, Tokyo Institute of Technology, 2-12-2 Ookayama, Meguro, Tokyo 152-5882

(Received December 21, 2004; CL-041577)

We describe an efficient synthesis of H type 1 and 2 trisaccharides by one-pot glycosylation involving glycosidation of glycal epoxide.

Oligosaccharides play important roles in biological processes on cell surface and have served as important tumor markers. β -Galactoside **1** attached with α -fucoside at the C2 position (H-disaccharide) is often found in biologically active oligosaccharides such as H type 1 and 2 epitopes (**1a**) and (**1b**), and is known to be an appropriate tumor antigen (Scheme 1).¹ In order to develop chemical probes based on the structure of the H-disaccharide, an effective methodology for the synthesis of glycoconjugates containing the H-disaccharide is required.²




One-pot sequential glycosylation to form two and more glycosidic bonds, is an effective approach for the liquid-phase oligosaccharide synthesis.³ This approach involves sequential chemo- and regio-selective glycosylations without any protecting group manipulations and purification of each intermediate. In the one-pot glycosylation, reagents and resulting products should not interfere any following glycosylations. We have investigated one-pot glycosylation based on the chemoselective activation of various glycosyl donors with an appropriate activator, and recently reported the synthesis of a protected linear and branched trisaccharide libraries by the one-pot glycosylation method.⁴ Most of synthetic strategies are based on the in situ synthesis of oligosaccharides with a leaving group. Therefore, the synthesis of the trisaccharides **1** based on the one-pot glycosylation strategy requires the glycosidation of glycosyl donor attached with saccharide at the C2 position to form 1,2-*trans*-glycosidic bonds. However, the absence of the participating substituents of the glycosyl donors at the C2 position makes it difficult to stereoselectively form the 1,2-*trans*-glycosidic bond. Therefore, an effective one-pot glycosylation method for the

synthesis of various C2 glycosylated oligosaccharides is required. Herein we report the one-pot synthesis of H Type 1 and 2 trisaccharide units using a glycal epoxide.

Our strategy for the one-pot synthesis of H Type 1 and 2 trisaccharide units **1** is based on the preparing glycosyl acceptors **5** by glycosylation of acceptor **3** with glycal epoxide **2**. The glycal epoxides are known to undergo stereoselective glycosidation to provide glycosides **5** bearing a hydroxy group at the C2 position linked through a 1,2-*trans*-glycosidic bond.⁵ Subsequent glycosylation of the hydroxy group with glycosyl donor **1** would provide the protected H Type 1 and 2 trisaccharides **1**.

We first conducted the one-pot synthesis of H type 1 trisaccharide **9aA** bearing a primary amino group using the three building blocks **2**, **3a**, and **4A** (Scheme 2). Our initial investigation involves the stepwise synthesis of the protected trisaccharide **7aA** as shown in Scheme 2. Treatment of glucosamine **3a** with 1.2 equiv. of the glycal epoxide **2** in the presence of ZnCl_2 in CH_2Cl_2 provided the desired disaccharide **6** in 74% yield with complete β -selectivity. Use of TMSOTf as an activator resulted in a significant amount of silylated products. Fucosylation of the resulting disaccharide **6** with thiofucose **4A** smoothly proceeded in stereoselective manner to provide α -fucose **7aA** in good yield with complete α -selectivity.



Next, we examined one-pot glycosylation using **2**, **3a**, and **4A**. To glucosamine **3a** was added 1.2 equiv. of the glycol epoxide **2** and 2.5 equiv. of ZnCl_2 at -78°C . The reaction mixture was stirred at 0°C for 2 h. Subsequently, 2.5 equiv. of thiofucose **4A**, 3.0 equiv. of NIS, and a catalytic amount of TfOH at 0°C were added to the reaction mixture. After stirring at the same temperature for 2 h, the reaction mixture was quenched. After removal of the solvent, the residue was purified by silica gel chromatography and gel permeable chromatography to provide trisaccharide **7aA** in 46% yield based on **3a**. The analytical data of trisaccharide **7aA**, synthesized by one-pot glycosylation were identical with those of trisaccharide **7aA** by the stepwise synthesis.

Deprotection of the protected trisaccharide **7aA** was investigated. Treatment of **7aA** with $\text{NH}_2\text{NH}_2 \cdot \text{H}_2\text{O}$ in EtOH at reflux for 18 h, followed by acetylation of the amine provided acetamide **8aA** in 70% yield. Hydrogenolysis of both benzyl ethers and an azido group with H_2 in the presence of Pd/C provided the H type 1 trisaccharide **9aA** bearing with an amino alkyl chain at the reducing end in quantitative yield.

In order to demonstrate the feasibility of the method, we planned the combinatorial synthesis of a small oligosaccharide library **7aA-cB** based on the structure of H type 1 and 2 trisaccharide by one-pot glycosylation. (Figure 1 and Scheme 3). Six building blocks **2**, **3a-3c**, and **4A-4B** were designed for





Figure 1.



Scheme 3. Reagents and conditions: a) ZnCl_2 , CH_2Cl_2 , -78 to -20°C , 2.0 h; b) NIS, TfOH, CH_2Cl_2 , 0°C , 2.0 h.

the library synthesis.

The parallel synthesis of the 6 oligosaccharides **7** by one-pot glycosylation was performed utilizing Carusel[®], which controls the reaction temperature and the stirring rate in 10 reaction vessels. The six reaction vessels were set up with activated MS-4Å. Each acceptor **3a-3c** was added to the two reaction vessels, respectively and the reaction vessels were cooled to -78°C . The glycol epoxide **2** (1.2 equiv.) and ZnCl_2 (2.5 equiv.) were added to all the vessels at -78°C . The reaction mixtures were warmed to -20°C and stirred for 2 h at the same temperature. Subsequently, 2.5 equiv. of thiofucose **4A** (2.5 equiv.) or thiorthamnoside **4B** (2.5 equiv.), NIS (3.0 equiv.), and a catalytic amount of TfOH at 0°C were added to the reaction mixture. After stirring for 2 h at the same temperature, the reaction mixture was quenched with NEt_3 . The residues were purified by silica gel chromatography, followed by gel permeable chromatography to provide trisaccharide **7aA-bC** in moderate yields (42–60% yields) based on **3**.

In conclusion, we have demonstrated one-pot synthesis of H type 1 and 2 trisaccharides **7** using the glycol epoxide **2**. The glycosylation of glycol epoxide **2** with ZnCl_2 provided disaccharide possessing the hydroxy-free C2. The secondary hydroxy group subsequent undergo glycosylation to provide trisaccharide in good yield in one-pot. The one-pot sequential glycosylation should be useful to prepare oligosaccharides containing the H-disaccharide moiety.

References

- a) S. Hakomori and Y. Zhang, *Chem. Biol.*, **4**, 97 (1997).
b) D. S. Newburg, *Curr. Med. Chem.*, **6**, 117 (1999).
- a) S. J. Danishefsky, V. Behar, J. T. Randolph, and K. O. Lloyd, *J. Am. Chem. Soc.*, **117**, 5701 (1995). b) K. R. Love, R. B. Andrade, and P. H. Seeberger, *J. Org. Chem.*, **66**, 8165 (2001).
- a) S. Raghavan and D. Kahne, *J. Am. Soc. Chem.*, **115**, 1580 (1993). b) S. V. Ley and H. W. M. Pripke, *Angew. Chem., Int. Ed.*, **33**, 2292 (1994). c) H. K. Chenault and A. Castro, *Tetrahedron Lett.*, **35**, 9145 (1994). d) T. Tsukida, M. Yoshida, K. Kurosawa, Y. Nakai, T. Achiha, T. Kiyoi, and H. Kondo, *J. Org. Chem.*, **62**, 6876 (1997). e) X.-S. Ye and C.-H. Wong, *J. Org. Chem.*, **65**, 2410 (2000).
- a) H. Yamada, T. Harada, and T. Takahashi, *J. Am. Chem. Soc.*, **116**, 7919 (1994). b) H. Yamada, T. Harada, T. Miyazaki, and T. Takahashi, *Tetrahedron Lett.*, **35**, 3979 (1994). c) H. Yamada, T. Kato, and T. Takahashi, *Tetrahedron Lett.*, **40**, 4581 (1999). d) T. Takahashi, M. Adachi, A. Matsuda, and T. Doi, *Tetrahedron Lett.*, **41**, 2599 (2000). e) H. Yamada, H. Takimoto, T. Ikeda, H. Tsukamoto, T. Harada, and T. Takahashi, *Synlett*, **2001**, 1751. f) H. Tanaka, M. Adachi, H. Tsukamoto, T. Ikeda, H. Yamada, and T. Takahashi, *Org. Lett.*, **4**, 4213 (2002). g) H. Tanaka, M. Adachi, and T. Takahashi, *Tetrahedron Lett.*, **45**, 1433 (2004). h) M. Adachi, H. Tanaka, and T. Takahashi, *Synlett*, **2004**, 609. i) H. Tanaka, M. Adachi, and T. Takahashi, *Chem.—Eur. J.*, (2005), in press.
- a) J. T. Randolph and S. J. Danishefsky, *Angew. Chem., Int. Ed.*, **33**, 1470 (1994). b) M. T. Bilodeau, T. K. Park, S. Hu, J. T. Randolph, S. J. Danishefsky, P. O. Livingston, and S. Zhang, *J. Am. Chem. Soc.*, **114**, 7840 (1995). c) S. J. Danishefsky and M. T. Bilodeau, *Angew. Chem., Int. Ed.*, **35**, 1380 (1996). d) D. Sames, X.-T. Chen, and S. J. Danishefsky, *Nature*, **389**, 587 (1997). e) J. T. Randolph, K. F. McClure, and S. J. Danishefsky, *J. Am. Chem. Soc.*, **117**, 5712 (1995).