


Stereoselective Formation of Alkenyl Halides via Magnesium Halide Promoted Ring Opening of Bis-Activated Cyclopropanes

Yi Wang and Hon Wai Lam*

School of Chemistry, University of Edinburgh,
The King's Buildings, West Mains Road,
Edinburgh, EH9 3JJ, United Kingdom

h.lam@ed.ac.uk

Received November 5, 2008




In the presence of stoichiometric magnesium halides, a range of bis-activated cyclopropanes undergo highly stereoselective ring-opening reactions to produce multisubstituted alkenyl halides. More highly functionalized compounds may be obtained by trapping of the magnesium enolate intermediates *in situ*.

Cyclopropanes, the smallest unsaturated carbocycles, are an important class of building blocks for organic synthesis.¹ The high strain energy stored within the three-membered ring of cyclopropanes may be harnessed in the design of novel chemical transformations that are often unavailable to simple alkenes.^{1,2} For example, cyclopropanes are highly susceptible to a broad range of addition reactions across the double bond^{1b} to provide functionalized cyclopropanes, which themselves are important


targets and common structural components in numerous biologically active natural products and pharmaceuticals.³ The strain energy of cyclopropanes may also be exploited in a host of cycloaddition and rearrangement reactions, providing numerous other molecules of interest.¹

During our recent investigations into the development of stereoselective carbometallation–ring-opening reactions of bis-activated cyclopropanes to produce multisubstituted alkenes,^{2d} the reaction of cyclopropane **1a** with commercially available Grignard reagent **2** was conducted (eq 1). Instead of furnishing products derived from addition of the nucleophilic alkyl functionality of **2**, the major product isolated in this reaction was the alkenyl bromide **3a**. We attributed this initially surprising result to reaction of cyclopropane **1a** with magnesium bromide produced from the Schlenk equilibrium⁴ of the Grignard reagent. This hypothesis was supported by reaction of **1a** with MgBr_2 itself, which also produced **3a**, but in 96% yield (eq 2).



This process is closely related to the results of Ma and co-workers, who described a series of alkali metal halide induced cyclopropane ring-opening–alkylation reactions that required an alkali metal carbonate additive for highest yields (representative example shown in eq 3).^{5a} However, their process differs from ours in that simple protonolysis of the initial ring-opened species was not described in their work.^{5a} The majority of examples described by Ma were conducted using a bis-activated cyclopropane that was unsubstituted at the alkene, although four results using trisubstituted cyclopropanes were reported (as in eq 3).^{5a}

Ma (2003):



To establish whether the serendipitous result of eq 2 could be translated into a general and efficient process, a range of bis-activated cyclopropanes **1a–1g** were reacted with stoichiometric quantities of magnesium halides, and these results are presented in Table 1. Using MgBr_2 , cyclopropanes **1a–1e** containing alkyl or aryl functionality on the alkene efficiently underwent the ring-opening reaction to provide alkenyl bromides


(1) For reviews, see: (a) Binger, P.; Büch, H. M. *Top. Curr. Chem.* **1987**, 135, 77–151. (b) Baird, M. S. *Cyclopropanes: Transformations* (Houben-Weyl); Thieme: Stuttgart, Germany, 1997; Vol. E17d/2, pp 2781–2790. (c) Nakamura, M.; Isabe, H.; Nakamura, E. *Chem. Rev.* **2003**, 103, 1295–1326. (d) Fox, J. M.; Yan, N. *Curr. Org. Chem.* **2005**, 9, 719–732. (e) Rubin, M.; Rubina, M.; Gevorgyan, V. *Synthesis* **2006**, 1221–1245. (f) Rubin, M.; Rubina, M.; Gevorgyan, V. *Chem. Rev.* **2007**, 107, 3117–3179. (g) Marek, I.; Simaan, S.; Masarwa, A. *Angew. Chem., Int. Ed.* **2007**, 46, 7364–7376.

(2) For recent, selected examples of investigations into the reactivity of cyclopropanes, see: (a) Fisher, L. A.; Fox, J. M. *J. Org. Chem.* **2008**, 73, 8474–8478. (b) Sherill, W. M.; Rubin, M. *J. Am. Chem. Soc.* **2008**, 130, 13804–13809. (c) Fordyce, E. A. F.; Luebbers, T.; Lam, H. W. *Org. Lett.* **2008**, 10, 3993–3996. (d) Wang, Y.; Fordyce, E. A. F.; Chen, F. Y.; Lam, H. W. *Angew. Chem., Int. Ed.* **2008**, 47, 7350–7353. (e) Pallerla, M. K.; Yap, G. P. A.; Fox, J. M. *J. Org. Chem.* **2008**, 73, 6137–6141. (f) Alnasleh, B. K.; Sherrill, W. M.; Rubin, M. *Org. Lett.* **2008**, 10, 3231–3234. (g) Yan, N.; Liu, X.; Fox, J. M. *J. Org. Chem.* **2008**, 73, 563–568. (h) Fordyce, E. A. F.; Wang, Y.; Luebbers, T.; Lam, H. W. *Chem. Commun.* **2008**, 1124–1126. (i) Rubina, M.; Woodward, E. W.; Rubin, M. *Org. Lett.* **2007**, 9, 5501–5504. (j) Masarwa, A.; Stanger, A.; Marek, I. *Angew. Chem., Int. Ed.* **2007**, 46, 8039–8042. (k) Trofimov, A.; Rubina, M.; Rubin, M.; Gevorgyan, V. *J. Org. Chem.* **2007**, 72, 8910–8920. (l) Chuprakov, S.; Malyshev, D. A.; Trofimov, A.; Gevorgyan, V. *J. Am. Chem. Soc.* **2007**, 129, 14868–14869. (m) Hirashita, T.; Shiraki, F.; Onishi, K.; Ogura, M.; Araki, S. *Org. Biomol. Chem.* **2007**, 5, 2154–2158. (n) Giudici, R. E.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2007**, 129, 3824–3825.

(3) (a) Liu, H. W.; Walsh, C. T. *Biochemistry of the Cyclopropyl Group In The Chemistry of the Cyclopropyl Group*; Rappoport, Z., Ed.; Wiley: Chichester, UK, 1987; pp 959–1025. (b) Salaun, J. *Top. Curr. Chem.* **2000**, 207, 1–67.


(4) Schlenk, W.; Schlenk, W., Jr. *Chem. Ber.* **1929**, 62, 920–924.


TABLE 1. Magnesium Halide Mediated Ring Openings of Various Cyclopropenes^a




$\text{R}^1\text{O}_2\text{C}$ EWG R^2 $\xrightarrow[\text{THF, rt}]{\text{MgX}_2 (1 \text{ equiv})}$ X CO_2R^1 EWG R^2 **3a-3l**

Cyclopropenes













1a R = Ph **1d** R = CH₂CH₂Ph
1b R = *p*-Tol **1e** R = CH₂CH₂OBz
1c R = *n*-Bu


1f

1g

entry	MgX ₂	substrate	product	yield (%)
1	MgBr ₂	1a		96 ^b
2		1b		72
3		1c		75
4		1d		87
5		1e		84
6	MgI ₂	1a		97
7		1b		94
8		1d		99
9		1g		77
10 ^d	MgCl ₂	1b		83
11 ^d		1c		85
12 ^d		1f		62

^a Reactions were carried out using 0.20 mmol of cyclopropene in THF (2 mL). ^b A preparative scale reaction conducted using 2.00 mmol of cyclopropene **1a** in THF (20 mL) also provided **3a** in 96% yield. ^c Product isolated as an inseparable 9:1 *Z:E* mixture. ^d Reaction conducted at 40 °C. ^e The product was accompanied by ca. 6% of an inseparable regioisomeric impurity.

3a–3e, respectively, in good yields and with high stereoselectivities⁶ (entries 1–5). In all cases, the bromide anion attacked the least substituted carbon of the alkene with high selectivity, though small quantities of regioisomeric products were observed in a few cases (entries 10 and 11). This regioselectivity is opposite to that seen in the large majority of cyclopropene carbometallation and hydrometalation reactions described previously^{1d} but is consistent with the results of Ma and co-workers (eq 3).^{5a} Ring openings with MgI₂ and MgCl₂ also occurred smoothly to furnish alkenyl halides **3f–3l** (entries 6–12), though a reaction temperature of 40 °C was required in the case of MgCl₂ (entries 10–12). The ability to prepare alkenyl chlorides is notable since this feature was not reported previously.^{5a} In addition to dimethyl malonate derived substrates **1a–1e**, cyclopropenes **1f** and **1g** containing ethyl esters or a phenylsulfone, respectively, were competent substrates (entries

SCHEME 1. Possible Mechanism

9 and 12). Unfortunately, no reaction occurred when MgF₂ was employed, presumably due to the low solubility of this salt.⁷

The stereochemical outcome of these reactions, where inversion of configuration at the electrophilic vinylic center is observed, suggests that an in-plane S_N2-type mechanism^{8,9} (also referred to as S_NVσ^{9e}) is operative, rather than an addition–elimination process similar to that proposed in related cyclopropene carbometallation–ring-opening reactions.^{2d} Although S_NVσ mechanisms had long been considered to be energetically unfavorable compared to alternative pathways,¹⁰ a growing body of theoretical⁸ and experimental⁹ work suggests that this process is favorable in certain reactions.

Therefore, a possible mechanism for magnesium halide promoted cyclopropene ring opening, using a dimethyl malonate derived substrate for illustrative purposes, is outlined in Scheme 1. It is likely that bidentate coordination of magnesium halide to the cyclopropene (as in **4**) promotes nucleophilic attack by the halide ion to provide (via transition state **5**) magnesium enolate **6** that is protonated upon workup to furnish the product. This mechanistic scenario suggested that trapping of the magnesium enolates **6** to generate more highly functionalized products might be possible, in similar fashion to the results of Ma and co-workers (eq 3).^{5a} Accordingly, various one-pot multicomponent coupling reactions of bis-activated cyclopropenes, magnesium halides, and different electrophiles were investigated. However, we found these enolates to be rather unreactive, though Michael additions to β-unsubstituted enones using acetonitrile as the solvent were possible, without the requirement for additional basic additives (Scheme 2).^{5a}

In summary, magnesium halide salts promote ring-opening reactions of bis-activated cyclopropenes to provide trisubstituted alkenyl halides with high stereoselectivities. The magnesium enolates that are produced in these reactions may be trapped with enones to result in more highly functionalized products. Compared with related reactions,^{5a} this work extends the scope of the nucleophilic and electrophilic components to encompass chloride anions and enones, respectively, increasing the range of products that may be accessed.

Experimental Section

General Procedure A: Magnesium Halide Mediated Ring Opening of Cyclopropenes.

(7) Reactions conducted using other fluoride salts such as ZnF₂, KF, and CsF in THF or MeCN at elevated temperatures also returned unchanged starting material.


(8) (a) Glukhovtsev, M. N.; Pross, A.; Radom, L. *J. Am. Chem. Soc.* **1994**, *116*, 5961–5962. (b) Luchini, V.; Modena, G.; Pasquato, L. *J. Am. Chem. Soc.* **1995**, *117*, 2297–2300. (c) Kim, C. K.; Hyun, K. H.; Kim, C. K.; Lee, I. *J. Am. Chem. Soc.* **2000**, *122*, 2294–2299. (d) Bach, R. D.; Baboul, A. G.; Schlegel, H. B. *J. Am. Chem. Soc.* **2001**, *123*, 5787–5793.

(9) (a) Ochiai, M.; Oshima, K.; Masaki, Y. *J. Am. Chem. Soc.* **1991**, *113*, 7059–7061. (b) Okuyama, T.; Takino, T.; Sato, K.; Ochiai, M. *J. Am. Chem. Soc.* **1998**, *120*, 2275–2282. (c) Ochiai, M.; Nishi, Y.; Hirobe, M. *Tetrahedron Lett.* **2005**, *46*, 1863–1866. (d) Shiers, J. J.; Shipman, M.; Hayes, J. F.; Slawin, A. M. Z. *J. Am. Chem. Soc.* **2004**, *126*, 6868–6869. (e) Miyauchi, H.; Chiba, S.; Fukamizu, K.; Ando, K.; Narasaka, K. *Tetrahedron* **2007**, *63*, 5940–5953. See also ref 5a.

(10) Kelsey, D. R.; Bergman, R. G. *J. Am. Chem. Soc.* **1971**, *93*, 1953–1961.

(5) (a) Ma, S.; Zhang, J.; Cai, Y.; Lu, L. *J. Am. Chem. Soc.* **2003**, *125*, 13954–13955. For halide-catalyzed cyclopropene ring-opening–cycloaddition reactions with imines, see: (b) Ma, S.; Zhang, J.; Lu, L.; Jin, X.; Cai, Y.; Hou, H. *Chem. Commun.* **2005**, 909–911.

(6) The stereochemistries of alkenyl halides **3a**, **3d**, **3k**, and **7a** were determined on the basis of NOESY experiments. See Supporting Information for details.

SCHEME 2. One-Pot Cyclopropene Ring-Opening–Michael Reactions


^a The product was accompanied by ca. 5% of an inseparable regioisomeric impurity. ^b Yield in parentheses refers to a reaction conducted using 8.00 mmol of cyclopropene **1a**.

propene (0.20 mmol) in THF (1 mL + 1 mL rinse) was added via cannula to a vial containing the appropriate magnesium halide (0.20 mmol) and a stirrer bar. The resulting mixture was stirred at the indicated temperature for the indicated time and then filtered through a short plug of SiO_2 (ca. 4 cm high $\times$ 2 cm diameter) using EtOAc as eluent (ca. 50 mL). After the filtrate was concentrated in vacuo, purification of the residue by column chromatography afforded the alkenyl halide product.

(Z)-Dimethyl 2-(2-bromo-1-phenylvinyl)malonate (3a). The title compound was prepared according to General Procedure A from cyclopropene **1a** (46 mg, 0.20 mmol) and $MgBr_2$ (37 mg, 0.20 mmol) at room temperature for 2 h and purified by column chromatography (hexane $\rightarrow$ 5% EtOAc/hexane) to give a colorless oil (60 mg, 96%): IR (film) 2954, 2846, 1738 (C=O), 1620, 1491, 1437, 1265, 1201, 1151, 1076 cm^{-1} ; 1H NMR (360 MHz, $CDCl_3$) δ 7.41–7.38 (2H, m), 7.36–7.31 (3H, m), 6.75 (1H, d, $J = 0.6$ Hz), 4.50 (1H, d, $J = 0.6$ Hz), 3.76 (6H, s); ^{13}C NMR (62.9 MHz, $CDCl_3$) δ 167.1 (2 $\times$ C), 138.1 (C), 137.2 (C), 128.4 (2 $\times$ CH), 128.3 (2 $\times$ CH), 128.2 (CH), 110.6 (CH), 58.8 (CH), 52.9 (2 $\times$ CH_3); HRMS (EI) exact mass calcd for $C_{13}H_{13}^{79}BrO_4$ [M^+] 311.9992, found 311.9992.

General Procedure B: One-Pot Magnesium Halide Mediated Ring-Opening–Michael Reactions of Cyclopropenes. A solution of the appropriate cyclopropene (0.20 mmol) in MeCN (1 mL + 1 mL rinse) was added via cannula to a vial containing the appropriate enone (0.40 mmol), $MgBr_2$ (37 mg, 0.20 mmol), and a stirrer bar. The resulting mixture was stirred at 40 °C for 18 h. After cooling to room temperature, the mixture was filtered through a short plug of SiO_2 (ca. 4 cm high $\times$ 2 cm diameter) using EtOAc as eluent (ca. 50 mL), and the filtrate was concentrated in vacuo. Purification of the residue by column chromatography afforded the alkenyl halide product.

(Z)-Dimethyl 2-(2-bromo-1-phenylvinyl)-2-(3-oxobutyl)malonate (7a). The title compound was prepared according to General Procedure B from cyclopropene **1a** (46 mg, 0.20 mmol) and methyl vinyl ketone (32 μ L, 0.40 mmol) and purified by column chromatography (10–30% EtOAc/hexane) to give a colorless oil (60 mg, 78%): IR ($CHCl_3$) 2953, 1735 (C=O), 1613, 1491, 1437, 1366, 1254, 1171, 1092, 1030 cm^{-1} ; 1H NMR (360 MHz, $CDCl_3$) δ 7.38–7.31 (3H, m), 7.07–7.05 (2H, m), 6.92 (1H, s), 3.71 (6H, s), 2.53–2.50 (2H, m), 2.27–2.24 (2H, m), 2.09 (3H, s); ^{13}C NMR (62.9 MHz, $CDCl_3$) δ 206.6 (C), 169.5 (2 $\times$ C), 141.4 (C), 137.1 (C), 128.8 (2 $\times$ CH), 128.3 (2 $\times$ CH), 128.2 (CH), 112.3 (CH), 63.7 (C), 52.8 (2 $\times$ CH_3), 39.0 (CH_2), 29.9 (CH_3), 28.2 (CH_2); HRMS (ES) exact mass calcd for $C_{17}H_{20}^{79}BrO_5$ [$M + H$]⁺ 383.0489, found 383.0492.

Acknowledgment. This work was supported by EaStCHEM (The Edinburgh and St. Andrews Research School of Chemistry). The EPSRC National Mass Spectrometry Service Centre at the University of Wales, Swansea, is thanked for providing high-resolution mass spectra.

Supporting Information Available: Experimental procedures and full spectroscopic data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO802475X