This article was downloaded by: [Dalhousie University] On: 12 March 2013, At: 07:58 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

Crystal Structure of the First Monomeric Organotellurium(IV) Oxide: Bis(2dimethylaminomethylphenyl)tellur Oxide

Thomas M. Klapötke^a, Burkhard Krumm^a & Matthias Scherr^a

^a Department of Chemistry and Biochemistry, Ludwig-Maximilian University of Munich, Munich, Germany Version of record first published: 06 Jun 2009.

To cite this article: Thomas M. Klapötke, Burkhard Krumm & Matthias Scherr (2009): Crystal Structure of the First Monomeric Organotellurium(IV) Oxide: Bis(2dimethylaminomethylphenyl)tellurium(IV) Oxide, Phosphorus, Sulfur, and Silicon and the Related Elements, 184:6, 1347-1354

To link to this article: http://dx.doi.org/10.1080/10426500902947476

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <u>http://www.tandfonline.com/page/terms-and-conditions</u>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.



Crystal Structure of the First Monomeric Organotellurium(IV) Oxide: Bis(2-dimethylaminomethylphenyl)tellurium(IV) Oxide

Thomas M. Klapötke, Burkhard Krumm, and Matthias Scherr

Department of Chemistry and Biochemistry, Ludwig-Maximilian University of Munich, Munich, Germany

The crystal structure of $(2 \cdot Me_2 N CH_2 C_6 H_4)_2$ TeO was determined by X-ray diffraction on a single crystal. The crystal symmetry is monoclinic, space group P2₁, with unit cell dimensions a =9.2274(2), b = 10.1749(2), c = 11.5339(3) Å, β = 109.952°, V = 1017.90(4) Å³ and Z = 2. The structure has been solved and refined down to wR₂ = 0.0625, using 3675 independent reflections. The compound consists of isolated molecules with two intramolecular Te···N interactions of 2.565(4) and 2.755(6) Å stabilizing the tellurium atom.

Keywords Crystal structure; donor-stabilization; tellurium

INTRODUCTION

Organotellurium compounds containing intramolecular Te···N interactions have been investigated over the past years.¹⁻¹⁶ In a recent report, we were able to provide clear evidence for the existence of the tellurenyl fluoride $2-Me_2NCH_2C_6H_4TeF$, with the help of the stabilizing 2-dimethylaminomethylphenyl ligand.¹⁷ Besides other synthesized organotellurium(IV) compounds with this special nitrogen donor substituent, we also presented crystal structures of the organotellurium(IV) trifluoride $2-Me_2NCH_2C_6H_4TeF_3$,¹⁷

Received 29 April 2008; accepted 30 April 2008.

Dedicated to Professor Marian Mikołajczyk, CBMiM PAN in Łódź, Poland, on the occasion of his 70th birthday.

Financial support of this work by the University of Munich, the Fonds der Chemischen Industrie and the Deutsche Forschungsgemeinschaft (KL 636/10-1) is gratefully acknowledged.

Address correspondence to Thomas M. Klapötke, Department of Chemistry and Biochemistry, Ludwig-Maximilian University of Munich, Butenandtstr. 5–13(D), D-81377 Munich, Germany. E-mail: tmk@cup.uni-muenchen.de

the organotellurium(VI) oxofluoride salt $[(2-Me_2NCH_2C_6H_4)_2TeF_3O]_2$ -[SiF₆],¹⁷ and the organotelluronium salt $[(2-Me_2NCH_2C_6H_4)_2-TeOH]_2[SiF_6]$,¹⁸ all characterized by intramolecular Te···N secondary interactions. To the best of our knowledge only a few examples for crystal structures of organotellurium(IV) oxides exist, among others Ph₂TeO, $(C_6F_5)_2$ TeO and $(p-MeOC_6H_4)_2$ TeO.^{19–28} However, all structures have in common that they exist as dimers, tetramers, or polymers with $[Te(\mu-O)]_n$ (n = 2, 4, ∞) units. In this article, we present a monomeric crystal structure of an organotellurium(IV) oxide.

RESULTS AND DISCUSSION

The title compound $(2-Me_2NCH_2C_6H_4)_2$ TeO was obtained as a hydrolysis product of the corresponding organotellurium(IV) diazide, which was prepared from the organotellurium(IV) diffuoride with two equivalents of Me₃SiN₃ according to Scheme 1.¹⁷

The hydrolysis of the diazide, resulting from an incompletely closed vessel stored over a period of about 6 months, furnished the organotellurium(IV) oxide $(2Me_2NCH_2C_6H_4)_2$ TeO. After a period of about 2 months, partially decomposition was already noticeable, which could be proven by ¹²⁵Te NMR spectroscopy. An additional resonance at $\delta = 1160$ ppm was found, which is shifted to lower field compared to the corresponding organotellurium(IV) diazide ($\delta = 1090$ ppm¹⁷) and was, after the structure elucidation, assigned to the title compound.

Suitable single crystals were obtained from a solution of $(2Me_2NCH_2C_6H_4)_2Te(N_3)_2$ in dichloromethane in a loosely closed vessel at ambient temperature. The compound crystallizes in the monoclinic crystal system, space group $P2_1$, with Z = 2 (see Figure 1).

The molecules exist as discrete monomers without significant intermolecular interactions below the sum of the tellurium–oxygen van der Waals radii (vdWr TeO 3.58 Å).²⁹ This is in contrast to known structures of organotellurium(IV) oxides, which were reported to exist as dimers (Ph₂TeO,²⁸ (C₆F₅)₂TeO,^{22,23} and OC(μ -N^tBu)₂TeO²⁴) with two different Te–O distances, one characteristic double bond and an elongated coordinative bond, forming Te₂O₂ rings comprising bridging



SCHEME 1 Preparation of the tellurium(IV) diazide and the tellurium(IV) oxide as the product of hydrolysis.



FIGURE 1 ORTEP plot of the crystal structure of $(2-Me_2NCH_2C_6H_4)_2$ TeO; H atoms omitted for clarity. Selected bond lengths [Å] and angles [°]: Te1-O1 1.8293(10), Te1-C1 2.118(5), Te1-C10 2.134(5), Te1···N1 2.755(6), Te1···N2 2.566(5), C1-Te1-C10 97.7(2), C1-Te1-O1 92.60(18), C10-Te1-O1 89.69(19), O1-Te1···N1 82.40(13), O1-Te1···N2 159.66(17).

oxygen atoms. In case of $(p-\text{MeOC}_6\text{H}_4)_2\text{TeO}$, a polymeric $(\text{TeO})_{\infty}$ chain with two slightly different Te–O distances is reported.²¹ The Te–O distance in the title compound (1.829(1) Å) is slightly elongated compared to the calculated value of monomeric Ph₂TeO (1.814 Å),³⁰ but significantly shorter compared to the values for the tellurium oxygen double bonds in the crystal structures of Ph₂TeO $(1.871(2) \text{ Å})^{28}$ and $(C_6F_5)_2\text{TeO} (1.872(2)/1.87(1) \text{ Å})$.^{22,23} Due to the bridging character of the oxygen atom in these compounds, the double bond character is reduced. This explains the elongated Te–O bonds compared to the bond distance in the title compound and also compared to the calculated monomeric Ph₂TeO. The same findings with an elongated Te–O bond distance with the 2-dimethylaminomethylphenyl substituent compared to the calculated value in the monomeric phenyl substituted compound Ph2TeO were also reported previously for the hydroxytelluronium cation [2-Me₂NCH₂C₆H₄TeOH]⁺ (calc.: 1.937 Å³⁰ vs. exp.: 1.961(2) Å).¹⁸ Two intramolecular Te···N contacts of 2.755(6) Å (Te1···N1) and 2.565(4) Å (Te1···N2), slightly elongated compared to the contact distance in 2-Me₂NCH₂C₆H₄TeF₃ (2.405(4) Å)¹⁷ and considerably shorter compared to the distances in $(2-Me_2NCH_2C_6H_4)_2Te$ (3.048/3.145 Å),⁸ are stabilizing the tellurium atom. These secondary interactions, significantly shorter than the sum of the tellurium-nitrogen van der Waals radii (vdWr TeN 3.61 Å²⁹), are due to their sterical effects on the tellurium atom likely the reason for the monomeric structure. The Te-C bonds are in the range of common tellurium-carbon bond lengths (Te1-C1 2.117(5)/Te1-C10 2.135(5) Å vs. 2.099(4) Å in $2-Me_2NCH_2C_6H_4TeF_3^{17}$ and 2.12-2.20 Å in $(C_6F_5)_2TeO^{22,23}$). The C1-Te1-C10 angle (97.78(18)°) is more bent compared to the values in the previously reported compounds (Ph₂TeO: 91.8(6)^{\circ 28} and (C₆F₅)₂TeO: $88.2(1)^{\circ}/88.6(5)^{\circ 22,23}$, which can also be explained as a result of the two secondary contacts of the coordinating substituent. This leads to a distorted octahedral arrangement around the tellurium atom, taking the free valence electron pair of the tellurium atom into account. The coordination of the aminomethyl nitrogen atoms creates two five-membered heterocycles with four-coordinated amine moieties.

A view of the unit cell along the *a*-axis of the title compound is shown in Figure 2. The packing can best be described as a layered structure along the 2_1 screw axes in the [010] direction.

EXPERIMENTAL SECTION

General Remarks

All manipulations of air- and moisture-sensitive materials were performed under an inert atmosphere of dry argon using flame-dried glass vessels.³¹ The solvent dichloromethane was dried by standard methods stored under N₂ and freshly distilled prior to use. All reagents were used as received (Sigma-Aldrich, Fluka) if not stated otherwise. NMR spectra were recorded on a JEOL Eclipse 400 instrument at 25°C, and chemical shifts were determined with respect to Me₂Te (¹²⁵Te, 126.1 MHz).

X-Ray Crystallography

For the compound, an Oxford Xcalibur3 diffractometer with a CCD area detector was employed for data collection using $Mo-K_{\alpha}$ radiation



FIGURE 2 View of the unit cell of $(2-Me_2NCH_2C_6H_4)_2$ TeO along [100]; H atoms omitted and only selected atoms labeled for clarity.

 $(\lambda = 0.71073 \text{ Å})$. The structure was solved using direct methods (SIR97³²) and refined by full-matrix least-squares on F^2 (SHELXL³³). All non-hydrogen atoms were refined anisotropically. Satisfactory atomic positions of the solvent molecules could not reliably be determined, therefore the disordered solvent was treated as a diffuse contribution using the program SQUEEZE. 34,35 SQUEEZE calculated 157 Å³ void space per unit cell and 43 electrons; one molecule of dichloromethane requires 42 electrons per unit cell. The absolute structure was determined by refinement of the Flack parameter (Flack parameter = 0.04(3), 1585Friedel pairs).^{36,37} See Table I. ORTEP plots are shown with thermal ellipsoids at the 50% probability level. Crystallographic data for the structure has been deposited with the Cambridge Crystallographic Data Centre under the depository number CCDC-681994. Copies of the data can be obtained free of charge upon application to The Director, CCDC, 12 Union Road, Cambridge CB2 1EZ; UK (Fax: int. code +(1223)336-033; e-mail for inquiry: fileserv@ccdc.cam.ac.uk; e-mail for deposition: deposit@ccdc.cam.ac.uk).

	$(2-Me_2NCH_2C_6H_4)_2TeO$
Empirical formula	$\mathrm{C}_{18}\mathrm{H}_{24}\mathrm{N}_{2}\mathrm{O}_{1}\mathrm{Te}$
Formula mass	411.99
Temperature [K]	100
Crystal size [mm]	0.21 imes 0.19 imes 0.10
Crystal description	colorless block
Crystal system	monoclinic
Space group	$P2_1$
a [Å]	9.2274(2)
<i>b</i> [Å]	10.1749(2)
c [Å]	11.5339(3)
β[°]	109.952(3)
V [Å ³]	1778.0(2)
Ζ	2
$ ho_{ m calc} \ [m g \ m cm^{-3}]$	1.344
$\mu \; [\mathrm{mm}^{-1}]$	1.465
F(000)	412
θ range [°]	3.7 - 26.0
Index ranges	$-10 \leq h \leq 11$
	$-12 \leq k \leq 12$
	$-14 \leq l \leq 12$
Reflcns collected	5227
Reflcns observed	3205
Reflcns unique	$3675 (R_{int} = 0.0245)$
R1, $wR2$ (2σ data)	0.0273, 0.0615
R1, wR2 (all data)	0.0316, 0.0625
Max./min. transm.	0.851/0.754
Data/restr./param.	3675/0/203
GOOF on F^2	0.984
Larg. diff. peak/hole [e/Å ³]	0.307/-0.577

TABLE I Crystal and Structure Refinement Data

Formation of $(2-Me_2NCH_2C_6H_4)_2$ TeO

The reaction of $(2\text{-}Me_2NCH_2C_6H_4)_2TeF_2$ with two equivalents of Me_3SiN_3 furnished the tellurium(IV) diazide $(2\text{-}Me_2NCH_2C_6H_4)_2$ $Te(N_3)_2.^{17}$ Single crystals of the tellurium oxide were obtained by storing a solution of the diazide in dichloromethane for 6 months in a loosely closed glass vessel at ambient temperature.

 ^{125}Te NMR (CDCl_3) δ 1160 ppm.

REFERENCES

L. Engman, A. Wojton, B. J. Oleksyn, and J. Sliwinski, *Phosphorus, Sulfur, and Silicon*, 179, 285 (2004).

- [2] G. Mugesh, A. Panda, S. Kumar, S. D. Apte, H. B. Singh, and R. J. Butcher, Organometallics, 21, 884 (2002).
- [3] J. E. Drake, M. B. Hursthouse, M. Kulcsar, M. E. Light, and A. Silvestru, J. Organomet. Chem., 623, 153 (2001).
- [4] A. Panda, G. Mugesh, H. B. Singh, and R. J. Butcher, *Organometallics*, 18, 1986 (1999).
- [5] T. S. Butcher and M. R. Detty, J. Org. Chem., 63, 177 (1998).
- [6] Z. Majeed, W. R. McWhinnie, and T. A. Hamor, J. Organomet. Chem., 549, 257 (1997).
- [7] T. A. Hamor, H. Chen, W. R. McWhinnie, S. L. McWhinnie, and Z. Majeed, J. Organomet. Chem., 523, 53 (1996).
- [8] R. Kaur, H. B. Singh, and R. J. Butcher, Organometallics, 14, 4755 (1995).
- [9] M. R. Detty, A. J. Williams, J. M. Hewitt, and M. McMillan, Organometallics, 14, 5258 (1995).
- [10] M. R. Detty, A. E. Friedman, and M. McMillan, Organometallics, 13, 3338 (1994).
- [11] A. G. Maslakov, W. R. McWhinnie, M. C. Perry, N. Shaikh, S. L. W. McWhinnie, and T. A. Hamor, J. Chem. Soc., Dalton Trans., 619 (1993).
- [12] H. B. Singh, N. Sudha, and R. T. Butcher, Inorg. Chem., 31, 1431 (1992).
- [13] V. I. Minkin, I. D. Sadekov, A. A. Maksimenko, O. E. Kompan, and Y. T. Struchkov, J. Organomet. Chem., 402, 331 (1991).
- [14] H. B. Singh, N. Sudha, A. A. West, and T. A. Hamor, J. Chem. Soc., Dalton Trans., 907 (1990).
- [15] N. Al-Salim, A. A. West, W. R. McWhinnie, and T. A. Hamor, J. Chem. Soc., Dalton Trans., 2363 (1988).
- [16] M. A. K. Ahmed, A. E. McCarthy, W. R. McWhinnie, and F. J. Berry, J. Chem. Soc., Dalton Trans., 771 (1986).
- [17] T. M. Klapötke, B. Krumm, and M. Scherr, Z. Anorg. Allg. Chem., 633, 1618 (2007).
- [18] T. M. Klapötke, B. Krumm, and M. Scherr, Acta Crystallogr., E63, 04189 (2007).
- [19] T. M. Klapötke, B. Krumm, and M. Scherr, Acta Crystallogr., C64, m17 (2008).
- [20] G. Schatte, T. Chivers, H. M. Tuononen, R. Suontamo, R. Laitinen, and J. Valkonen, *Inorg. Chem.*, 44, 443 (2005).
- [21] J. Beckmann, D. Dakternieks, A. Duthie, F. Ribot, M. Schürmann, and N. A. Lewcenko, Organometallics, 22, 3257 (2003).
- [22] D. Naumann, W. Tyrra, R. Hermannn, I. Pantenburg, and M. S. Wickleder, Z. Anorg. Allg. Chem., 628, 833 (2002).
- [23] T. M. Klapötke, B. Krumm, P. Mayer, H. Piotrowski, and O. P. Ruscitti, Z. Naturforsch., B57, 145 (2002).
- [24] G. Schatte, T. Chivers, C. Jaska, and N. Sandblom, Chem. Commun., 1657 (2000).
- [25] K. Kobayashi, N. Deguchi, O. Takahashi, K. Tanaka, E. Horn, O. Kikuchi, and N. Furukawa, Angew. Chem., Int. Ed., 38, 1638 (1999).
- [26] T. Chivers, M. Parvez, and G. Schatte, Inorg. Chem., 38, 5171 (1999).
- [27] K. V. Domasevitch, V. V. Skopenko, and E. B. Rusanov, Z. Naturforsch., B51, 832 (1996).
- [28] N. W. Alcock and W. D. Harrison, J. Chem. Soc., Dalton Trans., 709 (1982).
- [29] A. Bondi, J. Phys. Chem., 68, 441 (1964).
- [30] J. Beckmann, D. Dakternieks, A. Duthie, N. A. Lewcenko, C. Mitchell, and M. Schürmann, Z. Anorg. Allg. Chem., 631, 1856 (2005).
- [31] D. F. Shriver and M. A. Drezdzon, *The Manipulation of Air Sensitive Compounds* (Wiley, New York, 1986).

- [32] A. Altomare, G. Cascarano, C. Giacovazzo, A. Guagliardi, A. G. G. Moliterni, M. C. Burla, G. Polidori, M. Camalli, R. Spagna, SIR97 (1997).
- [33] G. M. Sheldrick, SHELXL, University of Göttingen, Germany (1997).
- [34] A. L. Spek, SQUEEZE, University of Utrecht, The Netherlands (2001).
- [35] P. Van der Sluis and A. L. Spek, Acta Crystallogr., A46, 194 (1990).
- [36] H. D. Flack and G. Bernardinelli, J. Appl. Crystallogr., 33, 1143 (2000).
- [37] H. D. Flack, Acta Crystallogr., A39, 876 (1983).