

Electrogenerated Chemiluminescent Chemodosimeter Based on a Cyclometalated Iridium(III) Complex for Rapid and Sensitive Detection of Thiophenol

Kyoung-Rok Kim, Hoon Jun Kim, and Jong-In Hong

Anal. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.analchem.8b03445 • Publication Date (Web): 14 Dec 2018

Downloaded from <http://pubs.acs.org> on December 14, 2018

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



Electrogenerated Chemiluminescent Chemodosimeter Based on a Cyclometalated Iridium(III) Complex for Rapid and Sensitive Detection of Thiophenol

Kyoung-Rok Kim, Hoon Jun Kim, and Jong-In Hong*

Department of Chemistry, College of Natural Sciences, Seoul National University, Seoul 08826, Republic of Korea

ABSTRACT: Thiophenol is the simplest aromatic thiol that is utilized for various applications in industry and agriculture. However, it should be used with care because thiophenol is readily absorbed into the human body by inhalation and ingestion, which leads to serious internal injuries. Thus, there is an urgent need for real-time and accurate monitoring of thiophenol. Despite remarkable advantages of electrogenerated chemiluminescence (ECL) analysis, ECL thiophenol probes have never been reported. Herein, a new strategy for the rapid detection of thiophenol using an ECL turn-on chemodosimeter based on a cyclometalated Ir(III) complex is described. This analytical system showed superior sensitivity (limit of detection (LOD) value, 3.8 nM) in comparison to the conventional fluorescence method. In addition, our system exhibited remarkable selectivity and reaction rate towards thiophenol over other analytes. Moreover, it was successfully applied to quantify thiophenol in real water samples, providing a new proof-of-concept for field-monitoring based on ECL.


Electrogenerated chemiluminescence (ECL) is a process in which sequential electron transfer reactions on the surface of the electrode induce luminescence.¹ It is superior in comparison to conventional fluorescence assays because of its high sensitivity and low background signal. Additionally, ECL provides the possibility of potential point-of-care-testing (POCT) and field-monitoring with the simplicity of the equipment and method.¹⁻⁴ For these reasons, ECL has emerged as one of the most powerful techniques for the detection of various analytes.⁵⁻¹⁴ Therefore, it is possible to develop efficient environmental pollutant sensors for on-site real sample monitoring by introducing the intrinsic features of ECL to a single-molecule receptor. In this regard, our research group has developed single-molecule ECL sensors for various targets.¹⁵⁻¹⁸ Until now, most of ECL-based sensors were developed by using tris(2,2'-bipyridine)ruthenium(II) complexes ([Ru(bpy)₃]²⁺) and ruthenium(II) polyimine complexes with characteristic orange/red emissions due to the limited ligand-field splitting energy of the ruthenium metal center.¹⁹ To overcome this limitation, several groups introduced new ECL luminophores with high quantum efficiencies and various emission wavelengths.²⁰⁻²⁶ Among them, cyclometalated iridium(III) complexes have been most successfully employed as ECL luminophores because Ir(III) complexes allow tuning the emission wavelength through the ligand modification, along with remarkable ECL quantum yields.^{16,27-33}

Thiophenol (PhSH) is an important chemical reagent that is utilized as a raw material and intermediate to produce pharmaceuticals, agrochemicals, polymers, etc.³⁴⁻³⁶ Although thiophenol is important for various applications, it

should be used with care because of its high toxicity. Thiophenol has a median lethal concentration (LC₅₀) of 0.01–0.4 mM in fish and a median lethal dose (LD₅₀) of 46.2 mg/kg in mouse.^{37,38} In addition, thiophenol is readily assimilated into the human body by inhalation and ingestion, which leads to serious systemic injury, central nervous system damage, muscular weakness, coma, and even death.^{39,40} Therefore, there is a compelling need to develop a real-time and accurate detection method for thiophenol. To date, various methods have been developed for the detection of thiophenol, including surface-enhanced Raman scattering (SERS) spectroscopy,⁴¹ chromatography-coupled spectrometry,⁴² and colorimetric⁴³ and fluorescence assays.⁴⁴⁻⁴⁸ Among them, the fluorescence assay has been widely used over the past few decades because it is a non-destructive analytical process and can be applied for real-time monitoring.⁴⁹⁻⁵⁵ Most of the previously reported fluorescence assays could selectively detect thiophenol (pK_a 6.5) over aliphatic thiols (pK_a ~8.5) under physiological conditions because of the different degree of deprotonation. However, this method requires not only heavy equipment but also the expertise to conduct a complicated analytical procedure, which restricts rapid on-site detection.

In this study, a turn-on ECL chemodosimeter (probe **2**) based on a cyclometalated Ir(III) complex was developed for rapid and quantitative detection of thiophenol. Probe **2** selectively reacted with two molecules of thiophenol to produce **2**-PhS⁻ with ECL turn-on. The design concept of probe **2** toward thiophenol was based on the following (Scheme 1a): (i) (piq)₂Ir(acac) (piq = 1-phenylisoquinoline,

Scheme 1. (a) Design of probe 2 for rapid detection of thiophenol. (b) The proposed sensing mechanism of probe 2 for detection of thiophenol through ECL analysis.



acac = acetylacetone) was selected as a luminophore as it is known to exhibit strong ECL emission,²⁷ (ii) 2,4-dinitrophenyl (DNP) group, a well-known photo-induced electron transfer (PET) quencher, was introduced as a reaction site for thiophenol,⁵⁶ and (iii) an electron-withdrawing formyl group was introduced to accelerate the reaction rate.^{57–59} Based on this rationale, it was expected that probe 2 would rapidly react with the thiophenolate moiety via nucleophilic aromatic substitution (S_NAr) to generate 2-PhS⁻. Next, 2-PhS⁻ was directly oxidized on the Pt electrode to generate (2-PhS⁻)^{•+}, which was converted to the singlet excited state, (2-PhS⁻)^{*}, by one electron transfer from the tripropylamine (TPrA) radical (TPrA[•]), which eventually emitted ECL, as shown in Scheme 1b.^{1,2} The present ECL analysis system for detection of thiophenol showed superior sensitivity and lower limit of detection (LOD) value (3.8 nM) than the conventional photoluminescence method. It also showed remarkable selectivity and reaction rate toward thiophenol over other analytes, which is highly desirable for on-site detection. Moreover, probe 2 was successfully utilized to quantify thiophenol in real water samples, suggesting a new proof-of-concept for ECL-based field-monitoring.


EXPERIMENTAL SECTION

Materials and instruments. All reagents were purchased from either TCI (Tokyo Chemical Industry, Tokyo, Japan), Alfa (Alfa Aesar, MA, USA) or Sigma-Aldrich (Sigma-Aldrich Corp., MO, USA). All reagents were used without further purification. Deuterated solvents for NMR spectra were acquired from CIL (Cambridge Isotopic Laboratories, Ma, USA). Analytical thin layer chromatography was carried out with Kiesegel 60F-254 plates of Merck. Column chromatography was performed on silica gel 60 (230–400 mesh) from SILICYCLE. High-resolution mass spectral (HRMS) data (JEOL, JMS-700) with fast atom bombardment (FAB) positive mode were received directly from the National Center for Inter-University Research Facilities (NCIRF). Mass spectra were recorded with MALDI-TOF Microflex from Bruker. ¹H and ¹³C NMR spectra were

recorded in CDCl₃ or DMSO-*d*₆ using a Bruker Avance DPX-300 or Varian/Oxford As-500 instrument. All chemical shifts were reported in parts per million (ppm) using the residual proton signals of deuterated solvents as an internal reference. All absorption spectra were obtained on a JASCO V-730 spectrometer, and fluorescence spectra were obtained on a JASCO FP-6500 spectrometer and a Spectra-Max M2 spectrophotometer. The band width of all spectrometers for excitation and emission was fixed to 5 nm except for SpectraMax M2 spectrophotometer (sensitivity: high). For photophysical experiments, 2 mM stock solutions of probes 1 and 2 were prepared in DMSO and diluted with acetonitrile (CH₃CN). Various analyte stock solutions were prepared in deionized water and diluted to 10 mM concentration, while thiophenol was dissolved and diluted with acetonitrile.

Electrochemical and electrochemiluminescent (ECL) measurement. Electrochemical studies were performed with a CH Instruments 660 Electrochemical Analyzer (CH Instruments, Inc., TX, USA). Cyclic voltammetry (CV) was applied to individual solutions in order to investigate electrochemical properties with a CH Instruments 650B Electrochemical Analyzer (CH Instruments, Inc., TX, USA). Electrochemical properties were examined by cyclic voltammetry (CV) with a scan rate of 0.1 V/s, and differential pulse voltammetry (DPV) with the following conditions: sample width, 17 ms; pulse amplitude, 50 mV; pulse width, 50 mV; pulse period, 200 ms; and quiet time, 2 s. All potential values were adjusted relative to the ferrocene/ferrocenium (F_c/F_c^+) redox couple by measuring the oxidation potential of 1 mM ferrocene (vs Ag/Ag⁺) as a standard.

The ECL intensity profile was obtained using a low-voltage photomultiplier tube module (H-6780, Hamamatsu photonics K.K., Tokyo, Japan) during the cyclic voltammetry process in the range of 0–1.6 V vs. Ag/AgCl (scan rate: 0.1 V/s). A 25 μ L volume ECL cell was directly mounted on the PMT module with home-made mounting support during the experiments. The ECL spectrum was obtained by measuring ECL intensities with a series of optical filters (550, 580, 600, 620, 650, 670 and 690 nm).^{60,61} All the ECL

Scheme 2. synthetic route of probes 1 and 2^a

^aReagents and conditions: a) bis(pinacolato)diboron, KOAc, Pd(dppf)Cl₂, 1,4-dioxane, 90 °C; b) 1-chloroisoquinoline, Pd(PPh₃)₄, NaHCO₃, THF, H₂O, 80 °C; c) IrCl₃·xH₂O, 2-ethoxyethanol, H₂O, 100 °C; d) BBr₃, CH₂Cl₂, 0 °C → r.t.; e) acetylacetonate, Na₂CO₃, 2-ethoxyethanol, 80 °C; f) 1-chloro-2,4-dinitrobenzene, K₂CO₃, DMF, 100 °C. (Pd(dppf)Cl₂ = [1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium(II), KOAc = potassium acetate, THF = tetrahydrofuran, DMF = dimethylformamide)

data were collected by the simultaneous cyclic voltammetry in the solution. The ECL solutions commonly contained 100 mM TPrA (tripropylamine, Sigma-Aldrich, MO, USA) and 0.1 M tetrabutylammonium perchlorate (TBAP, TCI) as the supporting electrolyte in acetonitrile (CH₃CN, spectroscopy grade, ACROS). Especially, TPrA was selected and used as an ECL co-reactant. All the electrochemical and ECL experiments were referenced with respect to an Ag/Ag⁺ reference electrode in organic solvents, or to an Ag/AgCl reference electrode in aqueous solution. Cyclic voltammetry for ECL experiments was applied to the solutions at the scan rate of 0.1 V/s. The electrochemical and ECL solutions were freshly prepared in each experiment,

and Pt working electrode was polished with 0.05 μm alumina (Buehler, IL, USA) on a felt pad. Then the electrode was blown by ultra-pure N₂ gas for 1 min. A single solution was only used for one experiment and discarded after collecting data. The reported ECL values were obtained by averaging the values from at least three experiments with a good reliability.

Synthesis of probes 1 and 2. Probes 1 and 2 were synthesized in six steps as shown in Scheme 2. Synthesis began with commercially available 4-bromoanisole and 5-bromoanisole to afford boron compounds for next steps. After main ligands with the methyl ether protecting group were obtained through the conventional borylation/Suzuki cross-coupling reactions, the corresponding Ir(III) complex dimers were synthesized in refluxing 2-ethoxyethanol. Then, demethylation of aryl methyl ethers was conducted by boron tribromide (BBr₃), followed by chelation of acetylacetonate (acac) as an ancillary ligand under basic conditions to furnish 1-PhS⁻ and 2-PhS⁻. Finally, probes 1 and 2 were synthesized by introduction of the DNP group via S_NAr. Details of synthesis as well as spectral data (¹H NMR, ¹³C NMR and HRMS) of new compounds were described in the Supporting Information.

RESULTS AND DISCUSSION

Time-dependent photoluminescence response and the sensing mechanism. To prove the rate-accelerating effect of the formyl group, control probe, 1, was synthesized. Probes 1 and 2 showed significant turn-on photoluminescence (PL) response upon reaction with thiophenol (Figure 1a and 1b). The reaction rates of probes 1 and 2 were compared by measuring the time-dependent changes of PL in the presence of 40 equivalent of thiophenol in aqueous media (pH 7.4, HEPES buffer/CH₃CN = 1:1, v/v). As shown in Figure 1c, probe 2 reached saturation within 25 min, whereas probe 1 took 270 min to reach a plateau (Figure S1).




Figure 1. Time-dependent measurements of the photoluminescence (PL) response ($\lambda_{\text{ex}} = 450 \text{ nm}$) of probes 1 (10 μM) (a) and 2 (10 μM) (b) in the presence of thiophenol (400 μM) in H₂O/CH₃CN (1:1 v/v, pH 7.4, 10 mM HEPES). (c) Relative PL intensity changes at 613 nm for probe 1 (black line) and at 589 nm for probe 2 (red line) in the presence of thiophenol. (d) Relative PL intensity of probes 1 and 2 before (black bar) and after (red bar) the reaction with 400 μM thiophenol for 25 min in CH₃CN/H₂O (1:1 v/v, pH 7.4, 10 mM HEPES).




Figure 2. PL emission spectra ($\lambda_{\text{ex}} = 450 \text{ nm}$) of probe **2** ($10 \mu\text{M}$) in the presence of increasing amount of thiophenol. ($\text{CH}_3\text{CN}/\text{H}_2\text{O} = 1:1 \text{ v/v}$, pH 7.4, 10 mM HEPES) (b) Linear plot of the PL intensity at 589 nm upon addition of varying concentrations of thiophenol.

In addition, the PL intensity of probe **2** at $\sim 25 \text{ min}$ after addition of thiophenol showed a 31-fold increase at 589 nm, while probe **1** showed only a 4-fold increase in the PL intensity at 613 nm (Figure 1d). This indicated that the formyl group adjacent to DNP played a significant role in improving the reactivity of the probe. The strong electron-withdrawing formyl group assists in an efficient cleavage of the DNP group from the Meisenheimer type intermediate.⁵⁷⁻⁵⁹ It can also stabilize both the intermediate and the reaction product (**2-PhS**⁻) by a resonance-assisted hydrogen bond.⁶² Moreover, probe **2** exhibited stable PL emission over the pH range of 5–11 (Figure S2). These results indicated that our design concept of probe **2** was effective for the rapid detection of thiophenol.

To elucidate the sensing mechanism of probe **2** for thiophenol, ¹H NMR and MALDI-TOF analyses of probe **2** were performed in the presence of 40 equivalent of thiophenol. As depicted in Figure S3, the ¹H NMR spectrum of the reaction mixture was identical to the superimposition of those of **2-PhS**⁻, 2,4-dinitrophenyl phenyl sulfane (DN), and thiophenol. Next, the m/z signal of probe **2** was compared in the absence and presence of thiophenol by MALDI-TOF analysis. This comparison showed that the m/z signal (1121.533) of probe **2** disappeared and that

(789.118) of **2-PhS**⁻ appeared upon the addition of thiophenol (Figure S4). These observations indicated that the DNP group of probe **2** was cleaved through the $\text{S}_{\text{N}}\text{Ar}$ reaction with thiophenol, forming **2-PhS**⁻ and DN.

Photophysical properties of probe 2. The UV-visible absorption spectra of **1-PhS**⁻, **2-PhS**⁻, probe **1**, probe **2** and DN in $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ (1:1 v/v, pH 7.4, 10 mM HEPES) are shown in Figure S5a. All the iridium complexes displayed intense absorption bands below 375 nm, which correspond to spin-allowed ligand-centered (¹LC) transitions. In addition, the weak and broad absorption bands in the range of 380–550 nm were assigned to metal-to-ligand charge transfer (MLCT) transitions.⁶³ In case of DN, the intense absorption band appeared around 350 nm. However, the absorption spectra of probes in the presence of 40 equivalent of thiophenol did not show any noticeable changes (Figure S5b, c).

The sensitivity of probe **2** was investigated in aqueous media (pH 7.4, 10 mM HEPES buffer/ $\text{CH}_3\text{CN} = 1:1 \text{ v/v}$) by measuring the PL intensity at 589 nm with the concentration of thiophenol in the range of 0–300 μM . As shown in Figure 2a, the PL intensity of probe **2** increased significantly upon gradual addition of thiophenol until saturation was reached after the addition of more than 20 equivalent of thiophenol. The PL intensity showed an almost linear correlation ($R^2 = 0.962$) from 0 to 15 equivalent of thiophenol as depicted in Figure 2b and thus the LOD was calculated as 39 nM (signal-to-noise (S/N) ratio = 3). We then conducted a competitive PL analysis of probe **2** for thiophenol by adding various anions (counter ion, Na^+) and bi-thiols. As shown in Figure 3, the addition of a large excess (80 equivalent) of other analytes to probe **2** did not induce significant changes in the PL intensity. However, upon further addition of 40 equivalent of thiophenol to the mixtures, the intense PL intensities were observed. These results indicate that probe **2** is highly selective toward thiophenol without interference from other analytes.

ECL properties of probe 2. The ECL intensity of probe **2** was measured in aqueous media (pH 7.4, HEPES buffer/ $\text{CH}_3\text{CN} = 1:1 \text{ v/v}$, 0.1 M TPrA, 0.1 M HEPES, and 0.1 M tetrabutylammonium perchlorate (TBAP) as the supporting electrolyte). During cyclic voltammetry (CV) in the range of 0–1.6 V, no change occurred in the ECL intensity




Figure 3. PL emission intensities at 589 nm ($\lambda_{\text{ex}} = 450 \text{ nm}$) of probe **2** ($10 \mu\text{M}$) upon addition of each analyte ($800 \mu\text{M}$) in the absence (black bar) and presence of $400 \mu\text{M}$ thiophenol (red bar) in $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ (1:1 v/v, pH 7.4, 10 mM HEPES), [(1) None; (2) CO_3^{2-} ; (3) $\text{C}_2\text{O}_4^{2-}$; (4) CN^- ; (5) HCO_3^- ; (6) N_3^- ; (7) NO_3^- ; (8) OAc^- ; (9) F^- ; (10) Cl^- ; (11) Br^- ; (12) I^- ; (13) SO_4^{2-} ; (14) HS^- ; (15) Cys; (16) GSH; (17) Hcy].




Figure 4. (a) ECL intensities of probe **2** (10 μM) upon the addition of thiophenol in $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ (1:1 v/v, pH 7.4, 100 mM TPrA, 100 mM HEPES, and 0.1 M TBAP as the supporting electrolyte) while the potential is swept at a Pt disk electrode (diameter 2 mm) in the range 0–1.6 V (scan rate: 0.1 V/s). (b) Linear plot of the ECL intensity at 1.4 V upon addition of varying concentrations of thiophenol.

for probe **2**. However, it did show a concentration-dependent increase in the ECL intensity (up to 52-fold) until more than 20 equivalent of thiophenol had been added (Figure 4a). In particular, the ECL intensity at 1.4 V had an almost linear relationship ($R^2 = 0.995$) with the concentration of thiophenol in the range of 0–200 μM (Figure 4b). Furthermore, the ECL of probe **2** showed sufficiently stable maximum intensities at 1.4 V under various concentrations of thiophenol (Figure S6). The LOD was estimated to be 3.8 nM (signal-to-noise (S/N) ratio = 3), which is superior to that (39 nM) determined from the PL assay. To assess the detection sensitivity of probe **2**, we examined the changes in the ECL intensity of probe **2** in the presence of low concentrations of thiophenol. As shown in Figure S7, probe **2** showed a linear increase in the ECL intensity upon addition of thiophenol in the range of 0–100 nM. This is the first ECL method capable of detecting thiophenol at nanomolar concentrations.

A competitive ECL analysis was carried out to evaluate the selectivity of probe **2** for thiophenol by adding various anions (counter ion, Na^+) and biothiols. As depicted in Figure 5a, the addition of a large excess (80 equivalent) of other analytes to probe **2** did not induce any significant changes in the ECL intensity. However, upon the addition of 40 equivalent of thiophenol to the mixtures, remarkable changes in the ECL intensities were observed, except in the case of iodide, which is an oxidation-sensitive anion (Figure 5b).^{64,65} Because competitive PL analysis revealed that there was no interference from anions including iodide in the detection of thiophenol, this result confirmed that the




Figure 5. (a) ECL intensities of probe **2** (10 μM) in the presence of various analytes (800 μM each; thiophenol 400 μM) in $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ (1:1 v/v, pH 7.4, 0.1 M TPrA, 0.1 M HEPES, and 0.1 M TBAP as the supporting electrolyte) while the potential is swept at a Pt disk electrode (diameter 2 mm) in the range 0–1.6 V (scan rate: 0.1 V/s). (b) ECL intensities at 1.4 V of probe **2** (10 μM) upon addition of 800 μM various analytes in the absence (black bar) and presence of 400 μM thiophenol (red bar), [(1) None; (2) CO_3^{2-} ; (3) $\text{C}_2\text{O}_4^{2-}$; (4) CN^- ; (5) HCO_3^- ; (6) N_3^- ; (7) NO_3^- ; (8) OAc^- ; (9) F^- ; (10) Cl^- ; (11) Br^- ; (12) I^- ; (13) SO_4^{2-} ; (14) HS^- ; (15) Cys; (16) GSH; (17) Hcy].

iodide anion genuinely had no effect on the PL selectivity of probe **2** but did affect the electrochemical process.

ECL mechanism studies. Density functional theory (DFT) calculations were applied to probes **1** and **2**, and the reaction products (**1**-PhS $^-$ and **2**-PhS $^-$) to support the PET sensing mechanism (Figure S8). The highest occupied molecular orbitals (HOMOs) of probes **1** and **2** were mainly delocalized over the iridium centre and phenyl group, and their lowest occupied molecular orbitals (LUMOs) were localized on the DNP group, while the LUMOs of **1**-PhS $^-$ and **2**-PhS $^-$ were localized primarily on the isoquinoline group. This strongly suggested that the turn-on response of probes **1** and **2** relied on the PET mechanism. In addition, **2**-PhS $^-$ had a larger HOMO-LUMO gap than **1**-PhS $^-$ because of the strongly electron-withdrawing formyl groups on the phenyl rings of the piq ligands. These groups strongly stabilized the HOMO/LUMO levels (0.66 eV and 0.51 eV, respectively), increasing the HOMO-LUMO energy gap. This result also explains the blue-shifted emission maximum of **2**-PhS $^-$ (589 nm) as compared to that of **1**-PhS $^-$ (613 nm).

To confirm the theoretical predictions and ECL mechanism, CV and DPV were conducted to estimate the experimental HOMO/LUMO levels of **1**-PhS $^-$ and **2**-PhS $^-$ (Table S2). It was found that the LUMO of DNP (−4.36 eV)⁴⁷ lay between the HOMO and LUMO levels of **1**-PhS $^-$ and **2**-PhS $^-$,




Figure 6. HOMO/LUMO energy levels calculated from CV and DPV measurements and the electronic distributions of **1-PhS**^{••} and **2-PhS**^{••}, and generation of TPA^{••} through the catalytic pathway.

which implied that their PL would be quenched by DNP via the PET process (Figure S9a). The HOMO/LUMO energy levels of **1-PhS**^{••} and **2-PhS**^{••} were compared with the singly occupied molecular orbital (SOMO) energy level of the TPrA radical (TPrA[•]) and the HOMO energy level of TPrA to clarify the ECL mechanism.²⁷ As indicated in Figure S9b, the LUMO levels of **1-PhS**^{••} (−3.28 eV) and **2-PhS**^{••} (−3.31 eV) were sufficiently lower than the SOMO level of TPrA[•] so that an electron can be smoothly transferred from TPrA[•] to the LUMO of **1-PhS**^{••} and **2-PhS**^{••}, which produced singlet excited states in both **1-PhS**^{••} and **2-PhS**^{••}. Additionally, it was investigated whether **1-PhS**^{••} and **2-PhS**^{••} were capable of oxidizing TPrA through the catalytic pathway, which is critical for high efficiency of ECL. The HOMO level of **1-PhS**^{••} (−5.39 eV) was similar to that of TPrA (−5.38 eV), while the HOMO level of **2-PhS**^{••} (−5.88 eV) lay at a lower level than that of TPrA (Figure 6). Therefore, it was expected that **2-PhS**^{••} would produce TPrA[•] more efficiently than **1-PhS**^{••} via a smooth catalytic process. Ir(III) complexes with a formyl group on the main ligand exhibited weaker PL intensities compared to those without a formyl group.^{16,66–68} As expected, **2-PhS**^{••} exhibited much weaker PL intensity than **1-PhS**^{••}. However, the ECL intensity of **2-PhS**^{••}

was comparable to that of **1-PhS**^{••}, which could be explained by the smooth catalytic process of **2-PhS**^{••} (Figure S10). It is noteworthy that the relative maximum ECL intensities of **1-PhS**^{••} and **2-PhS**^{••} were about twice that of Ru(bpy)₃²⁺ (bpy = 2,2'-bipyridyl). Furthermore, the ECL efficiencies of **1-PhS**^{••} and **2-PhS**^{••} were higher than that of Ru(bpy)₃²⁺. The ECL spectrum of **2-PhS**^{••} was similar to the PL spectrum of **2-PhS**^{••}, which indicates that the ECL signal genuinely originated from **2-PhS**^{••} (Figure S11).

Quantification of thiophenol in water samples. The ECL assay coupled with probe **2** was applied to the quantification of thiophenol in real water samples by the standard addition method. Environmental water samples were collected from the Han River in Seoul and all ECL measurements were performed in the same manner as described above. The concentration of the diluted analytical solution was determined by comparing the ECL intensity with the titration curve in Figure 2. The actual sample concentrations before dilution were successfully calculated by multiplying with the dilution factor (2.5) (Table 1). Probe **2** showed good recovery of thiophenol in the range from 96% to 98%. All these results revealed that the present ECL system has great potential for field-monitoring of thiophenol in real samples.

CONCLUSIONS

In conclusion, a turn-on cyclometalated Ir(III) complex-based ECL chemodosimeter was developed for the detection of thiophenol. To the best of our knowledge, this is the first example of thiophenol detection based on ECL. Probe **2** was rationally designed to improve the reactivity toward thiophenol without loss of selectivity. The present system showed superior sensitivity and a lower LOD value (3.8 nM) compared to the values obtained by the conventional PL methods. In addition, probe **2** could be successfully used for the quantification of thiophenol in real water samples, which provides a new proof-of-concept for field-monitoring based on ECL. It is expected that our strategy will be helpful for developing ECL-based analysis systems for the detection of other biologically and environmentally important small molecules.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Detailed procedures for synthesis, UV-Vis absorption spectra, additional photoluminescence spectra, MALDI-TOF mass spectra, additional electrochemiluminescence studies, DFT calculations, electrochemical properties and NMR spectra.

AUTHOR INFORMATION

Corresponding Author

* Prof. Dr. J.-I. Hong, Department of Chemistry, Seoul National University, Seoul 08826 (Republic of Korea), Fax: (+82) 2-889-1568, E-mail: jihong@snu.ac.kr

Author Contributions

Table 1. Quantification of thiophenol in water samples from the Han River.^{a,b}

[PhSH] ^c	Rel. ECL intensity (Experiment) (n=3)	Rel. ECL intensity (Calculation)	Recovery (%)
0 μM	0.0031 ± 0.0002	-	-
50 μM	0.1174 ± 0.0021	0.1216	95.94 ± 1.72%
100 μM	0.2152 ± 0.0053	0.2218	97.12 ± 2.40%
200 μM	0.4151 ± 0.0107	0.4222	98.20 ± 2.53%

^aAll ECL intensities of probe **2** (10 μM) were measured during the cyclic voltammetry (CV) process in the range of 0–1.6 V vs. Ag/AgCl (scan rate: 0.1 V/s). ^bAnalytical solutions were prepared by diluting 0.8 mL of Han River samples containing various concentrations of thiophenol by 0.2 mL of 1.0 M HEPES buffer (pH 7.41, 1.0 M TPrA) and 1.0 mL of 0.20 M TBAP in acetonitrile. ^cConcentration of thiophenol in Han River samples.

All authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

This work was supported by the NRF grant (No. 2018R1A2B2001293) funded by the MSIP.

REFERENCES

- (1) Richter, M. M. *Chem. Rev.* **2004**, *104*, 3003-3036.
- (2) Miao, W. *Chem. Rev.* **2008**, *108*, 2506-2553.
- (3) Hu, L.; Xu, G. *Chem. Soc. Rev.* **2010**, *39*, 3275-3304.
- (4) Hesari, M.; Ding, Z. *J. Electrochem. Soc.* **2016**, *163*, H3116-H3131.
- (5) Michael, S.; Heng-Wei, L. *Angew. Chem., Int. Ed.* **2007**, *46*, 893-896.
- (6) Emanuela, B.; Isabelle, G.; Denis, B.; Paolo, P.; Neso, S.; Sandra, P. *Chem. - Eur. J.* **2009**, *15*, 5145-5152.
- (7) Oh, J.-W.; Kim, T. H.; Yoo, S. W.; Lee, Y. O.; Lee, Y.; Kim, H.; Kim, J.; Kim, J. S. *Sens. Actuators, B* **2013**, *177*, 813-817.
- (8) Chen, K.; Schmitt, M. *Chem. Commun.* **2014**, *50*, 5756-5759.
- (9) Liang, W.; Zhuo, Y.; Xiong, C.; Zheng, Y.; Chai, Y.; Yuan, R. *Anal. Chem.* **2015**, *87*, 12363-12371.
- (10) Li, P.; Jin, Z.; Zhao, M.; Xu, Y.; Guo, Y.; Xiao, D. *Dalton Trans.* **2015**, *44*, 2208-2216.
- (11) Li, H.; Sedgwick, A. C.; Li, M.; Blackburn, R. A. R.; Bull, S. D.; Arbault, S.; James, T. D.; Sojic, N. *Chem. Commun.* **2016**, *52*, 12845-12848.
- (12) Xu, Y.; Zhang, L.; Liu, Y.; Jin, Z.; Zhao, Q.; Yang, F.; Xiao, D. *Biosens. Bioelectron.* **2016**, *77*, 182-187.
- (13) Nasiri Khonsari, Y.; Sun, S. *Chem. Comm.* **2017**, *53*, 9042-9054.
- (14) Wu, F.-F.; Zhou, Y.; Zhang, H.; Yuan, R.; Chai, Y.-Q. *Anal. Chem.* **2018**, *90*, 2263-2270.
- (15) Shin, I.-S.; Bae, S. W.; Kim, H.; Hong, J.-I. *Anal. Chem.* **2010**, *82*, 8259-8265.
- (16) Kim, H. J.; Lee, K.-S.; Jeon, Y.-J.; Shin, I.-S.; Hong, J.-I. *Biosens. Bioelectron.* **2017**, *91*, 497-503.
- (17) Kim, S.-Y.; Kim, H. J.; Hong, J.-I. *RSC Adv.* **2017**, *7*, 10865-10868.
- (18) Rhee, H.; Kim, T.; Hong, J.-I. *Dalton Trans.* **2018**, *47*, 3803-3810.
- (19) Wei, H.; Wang, E. *Luminescence* **2011**, *26*, 77-85.
- (20) Richter, M. M.; Bard, A. J. *Anal. Chem.* **1996**, *68*, 2641-2650.
- (21) Richter, M. M.; Debad, J. D.; Striplin, D. R.; Crosby, G. A.; Bard, A. J. *Anal. Chem.* **1996**, *68*, 4370-4376.
- (22) McCall, J.; Bruce, D.; Workman, S.; Cole, C.; Richter, M. M. *Anal. Chem.* **2001**, *73*, 4617-4620.
- (23) Bruce, D.; Richter, M. M.; Brewer, K. J. *Anal. Chem.* **2002**, *74*, 3157-3159.
- (24) Anderson, J. D.; McDonald, E. M.; Lee, P. A.; Anderson, M. L.; Ritchie, E. L.; Hall, H. K.; Hopkins, T.; Mash, E. A.; Wang, J.; Padias, A.; Thayumanavan, S.; Barlow, S.; Marder, S. R.; Jabbour, G. E.; Shaheen, S.; Kippelen, B.; Peyghambarian, N.; Wightman, R. M.; Armstrong, N. R. *J. Am. Chem. Soc.* **1998**, *120*, 9646-9655.
- (25) Kulmala, S.; Håkansson, M.; Spehar, A. M.; Nyman, A.; Kankare, J.; Loikas, K.; Ala-Kleme, T.; Eskola, J. *Anal. Chim. Acta* **2002**, *458*, 271-280.
- (26) Gross, E. M.; Armstrong, N. R.; Wightman, R. M. *J. Electrochem. Soc.* **2002**, *149*, E137-E142.
- (27) Kim, J. I.; Shin, I.-S.; Kim, H.; Lee, J.-K. *J. Am. Chem. Soc.* **2005**, *127*, 1614-1615.
- (28) Fernandez-Hernandez, J. M.; Longhi, E.; Cysewski, R.; Polo, F.; Josel, H.-P.; De Cola, L. *Anal. Chem.* **2016**, *88*, 4174-4178.
- (29) Kerr, E.; Doeven, E. H.; Wilson, D. J. D.; Hogan, C. F.; Francis, P. S. *Analyst* **2016**, *141*, 62-69.
- (30) Zhou, Y.; Xie, K.; Leng, R.; Kong, L.; Liu, C.; Zhang, Q.; Wang, X. *Dalton Trans.* **2017**, *46*, 355-363.
- (31) Smith, Z. M.; Kerr, E.; Doeven, E. H.; Connell, T. U.; Barnett, N. W.; Donnelly, P. S.; Haswell, S. J.; Francis, P. S. *Analyst* **2016**, *141*, 2140-2144.
- (32) Zhou, Y.; Kong, L.; Xie, K.; Liu, C. *J. Organomet. Chem.* **2017**, *846*, 335-342.
- (33) Kapturkiewicz, A. *Anal. Bioanal. Chem.* **2016**, *408*, 7013-7033.
- (34) Shimada, K.; Mitamura, K. *J. Chromatogr. B: Biomed. Sci. Appl.* **1994**, *659*, 227-241.
- (35) Eychmüller, A.; Rogach, A. L. *Pure Appl. Chem.* **2000**, *72*, 179-188.
- (36) Love, J. C.; Estroff, L. A.; Kriebel, J. K.; Nuzzo, R. G.; Whitesides, G. M. *Chem. Rev.* **2005**, *105*, 1103-1169.
- (37) Hell, T. P.; Lindsay, R. C. *J. Environ. Sci. Health, Part B* **1989**, *24*, 349-360.
- (38) Munday, R. *Free Radical Biol. Med.* **1989**, *7*, 659-673.
- (39) Munday, R.; Manns, E. J. *Appl. Toxicol.* **1985**, *5*, 414-417.
- (40) Amrolia, P.; Sullivan, S. G.; Stern, A.; Munday, R. *J. Appl. Toxicol.* **1989**, *9*, 113-118.
- (41) Onuegbu, J.; Fu, A.; Glembocki, O.; Pokes, S.; Alexson, D.; Hosten, C. M. *Spectrochim. Acta, Part A* **2011**, *79*, 456-461.
- (42) Wang, T.; Chamberlain, E.; Shi, H.; Adams, C. D.; Ma, Y. *Int. J. Environ. Anal. Chem.* **2010**, *90*, 948-961.
- (43) Kim, H.; Swamy, K. M. K.; Kwon, N.; Kim, Y.; Park, S.; Yoon, J. *ChemPhysChem* **2017**, *18*, 1752-1754.
- (44) Jiang, W.; Fu, Q.; Fan, H.; Ho, J.; Wang, W. *Angew. Chem., Int. Ed.* **2007**, *46*, 8445-8448.
- (45) Jiang, W.; Cao, Y.; Liu, Y.; Wang, W. *Chem. Comm.* **2010**, *46*, 1944-1946.
- (46) Kand, D.; Mishra, P. K.; Saha, T.; Lahiri, M.; Talukdar, P. *Analyst* **2012**, *137*, 3921-3924.
- (47) Xiong, L.; Yang, L.; Luo, S.; Huang, Y.; Lu, Z. *Dalton Trans.* **2017**, *46*, 13456-13462.
- (48) Zhang, M.; Leng, T.; Shen, Y.; Wang, C. *Analyst* **2018**, *143*, 756-760.
- (49) Lin, W.; Long, L.; Tan, W. *Chem. Comm.* **2010**, *46*, 1503-1505.
- (50) Yu, D.; Huang, F.; Ding, S.; Feng, G. *Anal. Chem.* **2014**, *86*, 8835-8841.
- (51) Liu, X.; Qi, F.; Su, Y.; Chen, W.; Yang, L.; Song, X. *J. Mater. Chem. C* **2016**, *4*, 4320-4326.
- (52) Sun, Q.; Yang, S.-H.; Wu, L.; Yang, W.-C.; Yang, G.-F. *Anal. Chem.* **2016**, *88*, 2266-2272.
- (53) Yue, Y.; Huo, F.; Zhang, Y.; Chao, J.; Martínez-Máñez, R.; Yin, C. *Anal. Chem.* **2016**, *88*, 10499-10503.
- (54) Xie, X.; Li, M.; Tang, F.; Li, Y.; Zhang, L.; Jiao, X.; Wang, X.; Tang, B. *Anal. Chem.* **2017**, *89*, 3015-3020.
- (55) Yang, L.; Su, Y.; Geng, Y.; Zhang, Y.; Ren, X.; He, L.; Song, X. *ACS Sens.* **2018**, *3*, 1863-1869.
- (56) Zhang, R.; Ye, Z.; Yin, Y.; Wang, G.; Jin, D.; Yuan, J.; Piper, J. A. *Bioconjugate Chem.* **2012**, *23*, 725-733.
- (57) Um, I.-H.; Kim, M.-Y.; Kang, T.-A.; Dust, J. M. *J. Org. Chem.* **2014**, *79*, 7025-7031.
- (58) Kang, T.-A.; Cho, H.-J.; Um, I.-H. *Bull. Korean Chem. Soc.* **2014**, *35*, 2135-2138.
- (59) Murto, J. M.; Murto, M.-L. *Acta Chem. Scand.* **1966**, *20*, 297-302.
- (60) Cui, H.; Zou, G.-Z.; Lin, X.-Q. *Anal. Chem.* **2003**, *75*, 324-331.
- (61) Li, L.; Liu, H.; Shen, Y.; Zhang, J.; Zhu, J.-J. *Anal. Chem.* **2011**, *83*, 661-665.
- (62) Kim, H.; So, S. M.; Chin, J.; Kim, B. M. *Aldrichimica Acta* **2008**, *41*, 77-88.

- (63) Lamansky, S.; Djurovich, P.; Murphy, D.; Abdel-Razzaq, F.; Kwong, R.; Tsyba, I.; Bortz, M.; Mui, B.; Bau, R.; Thompson, M. E. *Inorg. Chem.* **2001**, *40*, 1704-1711.
- (64) Zu, Y.; Bard, A. J. *Anal. Chem.* **2000**, *72*, 3223-3232.
- (65) Schmittl, M.; Qinghai, S. *Chem. Comm.* **2012**, *48*, 2707-2709.
- (66) Chen, H.; Zhao, Q.; Wu, Y.; Li, F.; Yang, H.; Yi, T.; Huang, C. *Inorg. Chem.* **2007**, *46*, 11075-11081.
- (67) Wu, Y.; Wu, R.; Li, H.; Zeng, H.; Li, Y.; Wang, Q.; Shi, M.; Fan, X. *RSC Adv.* **2017**, *7*, 52621-52625.
- (68) Gao, H.; Li, Z.; Zhao, Y.; Qi, H.; Zhang, C. *Sens. Actuators, B* **2017**, *245*, 853-859.

