


This article was downloaded by: [Chulalongkorn University]

On: 03 January 2015, At: 07:20

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

### FACILE AND EFFICIENT METHOD FOR DEPROTECTION OF 1,3-OXATHIOLANES WITH N,N'-DIBROMO-N,N'-1,2-ETHANEDIYLBIS(p-TOLUENESULPHONAMIDE)

Ramin Ghorbani-Vaghei<sup>a</sup> & Ardeshir Khazaei

<sup>a</sup> Department of Chemistry, Faculty of Science, Bu-Ali Sina University, Hamadan, Iran

Published online: 16 Aug 2010.

To cite this article: Ramin Ghorbani-Vaghei & Ardeshir Khazaei (2004) FACILE AND EFFICIENT METHOD FOR DEPROTECTION OF 1,3-OXATHIOLANES WITH N,N'-DIBROMO-N,N'-1,2-ETHANEDIYLBIS(p-TOLUENESULPHONAMIDE), Phosphorus, Sulfur, and Silicon and the Related Elements, 179:9, 1787-1791, DOI: [10.1080/10426500490466508](https://doi.org/10.1080/10426500490466508)

To link to this article: <http://dx.doi.org/10.1080/10426500490466508>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

## FACILE AND EFFICIENT METHOD FOR DEPROTECTION OF 1,3-OXATHIOLANES WITH N,N'-DIBROMO-N,N'-1,2-ETHANEDIYLBIS(P- TOLUENESULPHONAMIDE)

Ramin Ghorbani-Vaghei and Ardeshir Khazaei  
Department of Chemistry, Faculty of Science, Bu-Ali Sina  
University, Hamadan, Iran

(Received January 6, 2004; accepted January 8, 2004)

*Aliphatic and aromatic 1,3-oxathiolanes are readily oxidized to aldehydes and ketones in good yields under mild conditions by N,N'-dibromo-N,N'-1,2-ethanediybis(p-toluenesulphonamide) [BNBTS].*

**Keywords:** BNBTS; deprotection; oxathiolanes; oxidation

### INTRODUCTION

The electrophilic nature of the carbonyl groups is a dominant feature of their extensive chemistry. One of the major challenging problems during many multistep syntheses is how to protect a carbonyl from nucleophilic attack until its electrophilic properties can be exploited. For this reason, deprotection of the carbonyl groups remains a crucial challenge to organic chemists. Among various functional groups, protection of the carbonyl groups as a 1,3-oxathiolane is important for the following reasons: they can be used as acyl carbanion equivalents<sup>1a,b,c</sup> for C–C bond formation; the chiral 1,3-oxathiolanes are valuable for synthesis of  $\alpha$ -hydroxyaldehydes;<sup>1a</sup> and 1,3-oxathiolanes are comparatively more stable than *O,O*-acetals in acidic conditions, and much easier to remove than *S,S*-acetals and as protecting groups of carbonyl compounds.<sup>2</sup> A variety of methods for the deprotection of oxathiolanes are reported employing isoamylnitrite,<sup>3</sup> chloramine-T,<sup>4</sup> TMSOTf alone<sup>5</sup> or in the presence of *p*-nitrobenzaldehyde,<sup>6</sup> polymer-supported *p*-nitrobenzaldehyde,<sup>7</sup> NCS-AgNO<sub>3</sub>,<sup>8</sup> I<sub>2</sub>-AgNO<sub>3</sub>,<sup>9</sup>


Address correspondence to Ramin Ghorbani-Vaghei, Department of Chemistry, Faculty of Science, Bu-Ali Sina University, Hamadan, Iran. E-mail: ghorbani@basu.ac.ir

NBS in acetone,<sup>10</sup> glyoxylic acid in the presence of Amberlyst 15 in a microwave oven<sup>11</sup> and ammonium tribromide.<sup>12</sup> However, these methods suffer from drawbacks such as the difficulty of removing the byproduct oxathioacetals derived from *p*-nitrobenzaldehyde,<sup>6</sup> the use of expensive polymer-supported reagent,<sup>7</sup> on the requirement of a large molar excess of reagents such as expensive silver salts.<sup>8,9</sup> Consequently, it seems that there is still a need for development of newer methods that can proceed under mild and economically appropriate conditions. Therefore, we decided to study the application of *N,N'*-dibromo-*N,N'*-1,2-ethanediybis(*p*-toluenesulphonamide) [**BNBTS**],<sup>13</sup> which is relatively easy to make.


## RESULTS AND DISCUSSION

We now report a convenient method for conversion of 1,3-oxathiolanes to aldehydes and ketones using **BNBTS** (Figure 1).

Aliphatic and aromatic 1,3-oxathiolanes were found to be oxidized to aldehydes and ketones using **BNBTS** in aqueous acetonitrile at room temperature (Scheme 1).



SCHEME 1



**FIGURE 1** Since **BNBTS** contain two bromine atoms that are attached to nitrogen atoms it is very possible that this reagent releases  $\text{Br}^+$  which can act as an electrophilic species,<sup>13</sup> in situ.

















The advantages of **BNBTS** are as follows:

1. The preparation of **BNBTS** is easy.
2. **BNBTS** is stable in atmospheric conditions for two months.
3. After completion of the reaction and evaporation of solvent, the sulphonamide is recovered and can be reused many times without decreasing the yield.

The results of the conversion of various 1,3-oxathiolanes to aldehydes and ketones are presented in Table I.

Since **BNBTS** contains two bromine atoms that are attached to nitrogen atoms, it is very possible that this reagent releases  $\text{Br}^+$  which can act as an electrophilic species,<sup>13</sup> in situ. The products of reaction

**TABLE I** Deprotection of 1,3-Oxathiolanes with **BNBTS** in Aqueous Acetonitrile at Room Temperature

Entry	Substrate	Product <sup>a</sup>	Time [min]/h	Yield (%)
1			4	90
2			[30]	93
3			2.5	84
4			[60]	89
5			[90]	92
6			[20]	82
7			7	85
8			10	90

<sup>a</sup>Products were characterized by their physical properties, comparison with authentic samples, and by spectroscopic methods.

with **BNBTS** were isolated simply by filtering off the sulfonamide **2** and evaporating the solvent from the filtrate. The method has advantages in terms of yields, simplicity of reaction conditions, short reaction times, and lack of side products. The recovered starting material **2** was rebrominated and used many times without reducing the yield.

## CONCLUSION

From the results obtained, we find that the described procedure and the reaction conditions are simple. The new reagent (**BNBTS**) is stable and the recovered reagent can be reused.

Infrared (IR) and nuclear magnetic resonance (NMR) spectra were recorded using a Shimadzu 435-U-04 spectrophotometer (KBr pellets) and a 90 MHz Jeol FT-NMR spectrometer, respectively.

## GENERAL PROCEDURE FOR DEPROTECTION OF 1,3-OXATHIOLANES USING BNBTS

A solution of the 1,3-oxathiolane (1.0 mmol) in acetonitrile (2 ml) was added quickly to a well-stirred solution of **BNBTS** (4 mmol) in aqueous 80% acetonitrile (10 ml) at room temperature. The mixture was stirred for the specified time (Table I). After completion of the reaction, the insoluble sulphonamide **2** was removed by filtration and washed with cold acetonitrile (5 ml). The solvent was then evaporated and ether was added. The ethereal phase was washed with saturated aqueous sodium carbonate and water and then was dried with  $\text{MgSO}_4$  and evaporated to give the product.

## REFERENCES

- [1] a) J. E. Lynch and E. L. Eliel, *J. Am. Chem. Soc.*, **106**, 2943 (1984); b) E. L. Eliel and S. Morris-Natsche, *J. Am. Chem. Soc.*, **106**, 2937 (1984); c) K. Utimoto, A. Nakamura, and S. Matsubara, *J. Am. Chem. Soc.*, **112**, 8189 (1990).
- [2] T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, 3rd ed. (John Wiley and Sons, New York, 1999), p. 329.
- [3] K. Fuji, K. Ichikawa, and E. Fujita, *Tetrahedron Lett.*, **19**, 3561 (1978).
- [4] D. W. Emerson and H. Wynberg, *Tetrahedron Lett.*, **12**, 3445 (1971).
- [5] T. Ravindranathan, S. P. Chavan, and S. W. Dantale, *Tetrahedron Lett.*, **36**, 2285 (1995).
- [6] a) T. Ravindranathan, S. P. Chavan, R. P. Tejwani, and J. P. Varghese, *J. Chem. Soc. Chem. Commun.*, **42**, 1750 (1991); b) T. Ravindranathan, S. P. Chavan, J. P.

- Varghese, S. W. Dantale, and R. B. Tejwani, *J. Chem. Soc. Chem. Commun.*, **45**, 1937 (1994).
- [7] T. Ravindranathan, S. P. Chavan, and M. M. Awachat, *Tetrahedron Lett.*, **35**, 8835 (1994).
- [8] a) S. V. Frye and E. L. Eliel, *Tetrahedron Lett.*, **26**, 3907 (1985); b) E. L. Eliel and K. Soai, *Tetrahedron Lett.*, **21**, 2859 (1981).
- [9] a) K. Nishide, K. Yokota, D. Nakamura, T. Sumiya, M. Node, M. Ueda, and K. Fuji, *Tetrahedron Lett.*, **34**, 3425 (1993); b) K. Nishide, K. Yokota, D. Nakamura, T. Sumiya, M. Node, M. Ueda, and K. Fuji, *Heterocycles*, **44**, 393 (1997).
- [10] B. Karimi, H. Seradj, and M. H. Tabaei, *Synlett.*, 1798 (2000).
- [11] S. P. Chavan, P. Soni, and S. K. Kamat, *Synlett.*, 1251 (2001).
- [12] E. Mondal, P. R. Sahu, G. Bose, and A. T. Khan, *Tetrahedron Lett.*, **43**, 2843 (2002).
- [13] a) A. Khazaei and A. Shirdarreh, *Synth. Commun.*, **29**, 4079 (1999); b) A. Khazaei, R. G. Vaghei, and M. Tajbakhsh, *Tetrahedron Lett.*, **42**, 5099 (2001).