Photolytic decomposition of some polycyclic α -diazoketones: generation and reactivity of 1,2-oxocarbenes in benzene with and without *tert*-butylamine



Peter J. N. Brown,^{*a*} J. I. G. Cadogan,^{*b*} Ian Gosney,^{*c*} Alexander Johnstone,^{*c*} R. Michael Paton^{*c*} and Norman H. Wilson^{*c*}

^a BP International Ltd., Research and Engineering Centre, Chertsey Road, Sunbury-on-Thames, Middlesex, UK TW16 7LN

^b Department of Chemistry, Imperial College of Science, Technology and Medicine,

South Kensington, London, UK SW7 2AY

^c Department of Chemistry, The University of Edinburgh, West Mains Road, Edinburgh, UK EH9 3JJ

Irradiation in benzene of the two isomeric 2,3-diazofluoranthenones 11 and 14 results in the formation of only products due to the formal insertion of the intermediate oxocarbenes into the C-H bonds of the solvent, unless the reactions are carried out in the presence of *tert*-butylamine, whereupon products of an apparent [3 + 2]-cycloaddition of the oxocarbene to the solvent are also obtained, *viz*. dihydrofurans 13 and 16, respectively. Wolff rearrangement of the generated oxocarbenes to give products derived from direct trapping of the ketene intermediates with *tert*-butylamine are not observed. The formal insertion product, 2-phenylfluoroanthen-3-ol 12 (from 11) undergoes partial oxidation during chromatographic purification on silica to generate a persistent free radical species 17 which has been detected and characterised by EPR spectroscopy. A reinvestigation of the photolytic decomposition of the related polycyclic α -diazoketone 5 under the same conditions has established that a more likely structure for the isolated dihydrofuran is 8, and not the reported isomeric dihydrofuran 20, which is formed from 8 by hydrogen migration during isolation by repeated thick-layer chromatography.

We reported some time ago^1 that photolysis of 2-diazoacenaphthen-1-one 1 gives rise to an oxocarbene 2, which can be trapped by aromatic receptors, *e.g. p*-xylene, to afford spiro-[acenaphthen-1,7'-*cis*-norcara-2',4'-diene]-2-one 3 (Scheme 1).



Scheme 1 Reagents and conditions: i, photolysis, p-xylene; ii, reflux, toluene, 18 h or catalytic amount of either AgClO₄ or p-CH₃C₆H₄SO₃H

The strain inherent in this system prompted isomersation to 2arylacenaphthen-1-one 4 when subjected to either heating in boiling toluene (18 h) or acid-catalysis with toluene-*p*-sulfonic acid. Significantly, aromatisation also occurred very rapidly with the production of a transient red colour under the influence of silver perchlorate (4%), thus providing the first extension of the silver-ion-catalysed cleavage of strained rings to the norcaradiene system.

We viewed the formation of the norcaradiene 3 from the

oxocarbene 2 as a function of the inability of the latter to undergo a Wolff rearrangement,² but it was of interest not only for its rarity,³ but also in comparison with the behaviour of oxocarbene 6, which is generated by photolytic decomposition 9-diazo-8,9-dihydro-4H-cyclopenta[d,e,f]phenanthren-8of one 5⁴ in benzene. Oxocarbene 6 is also reluctant to undergo a Wolff reaction, as evidenced by obtention of ca. 2% yield of carboxamide 10 from a trapping reaction of ketene intermediate 9 with tert-butylamine, but in sharp contrast to 2, its trapping to afford the corresponding norcaradiene 7 was not detected, and the major compound isolated was purported to be the dihydrofuran 8, which is the product of formal [3 + 2]cycloaddition of the oxocarbene 6 to the solvent (Scheme 2). We ascribed this difference in behaviour to the greater accessibility of the ketone group with respect to the bridgehead carbon atom [marked with an asterisk (*)] in 7 compared with 3.

In the present work, we have investigated the photolytic decomposition of the two isomeric diazofluoranthenones 11 and 14 in benzene and have found only the products of formal insertion of the intermediate oxocarbene into the C-H bond of benzene, unless the reactions are carried out in the presence of *tert*-butylamine, whereupon the products of apparent [3 + 2]-cycloaddition of the oxocarbene to the solvent are also obtained. Wolff rearrangement of the generated oxocarbenes to give products derived by direct trapping of ketene intermediates with *tert*-butylamine is not observed.

Results and discussion

Fluoranthene-2,3-quinone, prepared according to the method of Periasamy and Bhatt,⁵ was reacted with *p*-toluenesulfonylhydrazide to give a yellow crystalline material from which the light-sensitive diazoketones 11 and 14 could be isolated in 39 and 20% yield, respectively, by column



Scheme 2 Reagents and conditions: i, photolysis, benzene-Bu'NH₂ (ca. 30:1 v/v)

chromatography. The structural assignments are based largely on spectroscopic evidence, but are also consistent with what might be expected on the basis of prevailing attack by the hydrazide at the least sterically crowded carbonyl group of the quinone, and the spectral characteristics of the products obtained from the following chemical transformations.

On irradiation in benzene, 3-diazofluoranthen-2-(3H)-one 14 gave as the only isolable product 3-phenylfluoranthen-2-ol 15 in 38% yield. In the case of the isomeric 2-diazofluoranthen-3(2H)-one 11, photolysis under similar conditions, followed by rapid chromatography over silica, afforded a 74% yield of 2phenylfluoranthen-3-ol 12 as a fluorescent green crystalline solid which melted sharply at 140.5-141.5 °C. The ¹H NMR spectrum of this compound, unlike that of its isomer 15, was extremely broad at room temperature, and although the spectrum sharpened considerably at lower temperatures, or upon addition of D₂O, only progressive broadening occurred at temperatures up to 160 °C. EPR investigations indicated that this behaviour was associated with the presence in solution of a small amount of persistent radical species ($t_1 > 2$ days at room temperature) believed to be aryloxy radical 17. The EPR spectrum [Fig. 1(a)] consisted of a series of overlapping multiplets giving overall at least 28 lines ($a_{\rm H} = 0.06 \,\mathrm{mT}$). The wealth of fine structure is consistent with extensive delocalisation of the unpaired electron throughout the multi-ring system. The concentration of the radical was temperature dependent. On cooling to -50 °C the signal became almost undetectable, but returned to its full magnitude on re-warming to room



temperature. The amount at room temperature was estimated to be *ca.* 0.6% of the solute by comparison with standard solutions of 2,2-diphenyl-1-picrylhydrazyl. These results are in accord with the sharpening of the ¹H NMR spectrum at low temperature and implies that the radical exists in equilibrium with one or more dimers, whose formation is favoured at low temperature.

Radical 17 is believed to result from aerial oxidation of 2phenylfluoranthen-3-ol 12 at the surface of the silica during chromatographic purification. To test this hypothesis a solution of compound 12 in toluene was saturated with oxygen and subsequently treated with silica. Only traces of the signal were detected initially, but addition of the silica generated an intense signal identical to that described above. No signal was detected on photolysis of 11 in de-gassed benzene.

Further evidence for the assignment of the spectrum to radical 17 was provided by a repeat experiment in which diazo compound 11 was photolysed in hexadeuteriobenzene. The



Fig. 1 EPR spectra of (a) radical 17 and (b) the deuteriated analogue 18

resulting signal [Fig. 1(b)] is assigned to radical **18** and shows loss of fine structure resulting from replacement of 2-phenyl ring protons by deuterions.

Photolysis of the isomeric diazoketones 11 and 14 in benzene containing 6% v/v tert-butylamine led to the concomitant formation of minor amounts of the dihydrofurans 13 (3%) and 16 (12%), respectively. The structure of these compounds rests on their spectral properties, but further evidence is provided by ¹H NMR spectral comparison with the aforementioned dihydrofuran 8 obtained by irradiation of 9-diazo-8,9-dihydro-4*H*-cyclopenta[*d*,*e*,*f*]phenanthren-8-one 5 under similar conditions. In particular, the striking similarities in the position of the resonances due to H-13a (δ 4.69) of 14 and H-12a (δ 4.67) of 8 may be cited as evidence in favour of the foregoing assignments to the structures of the diazoketones 11 and 14.

Note that in contrast to Trost and Kinson,⁴ we have found that irradiation of 9-diazo-8,9-dihydro-4H-cyclopenta[d,e,f]phenanthren-8-one 5 in benzene proceeds in a manner analogous to the photolyses of diazoketones 11 and 14 described above, i.e. the sole isolable product is 9-phenyl-4H-cyclopenta[d, e, f] phenanthren-8-ol 19, the product of formal insertion into benzene, unless tert-butylamine (1.1% v/v) is present in which case 8 can also be isolated together with small amounts of the trapped product 10 arising from Wolff rearrangement of oxocarbene 6. While our assignment of 8 is consistent with its ¹H NMR spectrum, the latter is not wholly identical to that reported previously by Trost and Kinson." Comparison of both data suggests that a more likely structure for their compound is the isomeric dihydrofuran 20 which is formed from 8 by hydrogen migration (1,3-shifts) during isolation by repeated thick-layer chromatography. Most significant is the chemical shift difference for the methylene protons at C-4. The chemical shift for the methylene protons in 20 come at lower chemical shift (δ 3.50 vs. 4.28 for 8) in keeping with the change in hybridization of the 7b and 12b carbon atoms. The observed δ values compare favourably with values of 3.72⁶ and 4.13^7 for the corresponding methylene protons in 21 and 22, respectively. Additional evidence that our assignment was correct came from the observed rearrangement of 8 to 19 in the presence of acid.

The formation of the dihydrofurans, *e.g.* 13, may be rationalised by direct cycloaddition of the dipolar form 23b of the intermediate oxocarbene 23a to the solvent, but the role of *tert*butylamine in this mechanism is obscure. It remains possible that the reaction proceeds *via* the intermediacy of a zwitterionic species 24^8 and that cyclisation to dihydrofuran 13 is aided by *tert*-butylamine as shown in Scheme 3. The question of the



mechanism of formation of the dihydrofuran 8 remains an interesting problem. It is still possible that it arises via the intermediacy of norcaradiene species 7, since the formal insertion product 19 could arise via the same intermediate. However, 19 could also arise by rearrangement of the dihydrofuran, as demonstrated by its acid-catalysed rearrangement to 19 (vide supra), so that isolation of these two species does not by itself imply a common intermediate norcaradiene, and a dipolar cycloaddition mechanism could still hold.

Experimental

Melting points are uncorrected. ¹H NMR spectra were recorded on a Bruker WP-200 spectrometer for solutions in CDCl₃, unless stated otherwise, with chemical shifts quoted relative to Me₄Si as internal standard, and coupling constants, *J* values, in Hz; c refers to complex multiplet. IR spectra were recorded for Nujol mulls. Mass spectra were recorded on an AEI MS 902 mass spectrometer at 70 eV.

Photolyses were carried out using a water-cooled, 100 W medium-pressure mercury vapour lamp (Applied Photophysics) through a Pyrex filter. The reactions were carried out under an atmosphere of dry nitrogen, unless otherwise stated, the reaction mixture being stirred continuously. The course of the reactions was monitored by thick-layer chromatography (tlc) and the visible evolution of nitrogen gas.

The EPR instrumentation and methods were as previously described.⁹

Preparation of 2- and 3-diazofluoranthenones

Fluoranthene-2,3-quinone and *p*-toluenesulfonylhydrazide were reacted according to the method of Reid and Dietrich.¹⁰ The quinone (1.12 g, 4.59 mmol) and the hydrazide (0.88 g, 4.91 mmol) were suspended in methanol (20 ml) and the mixture heated under reflux, under an atmosphere of dry N₂, for 20 min with occasional swirling. The mixture was cooled, and the solvent removed on a rotary evaporator. The residue was taken up in CH₂Cl₂ (20 ml) and applied to the top of an alumina column which was wrapped in aluminium foil to exclude light. Elution with CH₂Cl₂-petrol (50:50) gave 2-diazofluoranthen-3(2*H*)one 11 (455 mg, 39%), mp 171–176 °C (decomp.) (lit.,⁷ 165 °C); ν_{max} /cm⁻¹ 2125 (C=N₂) and 1630 (C=O); $\delta_{\rm H}$ 8.10–7.28 (c) (Found: M⁺, 244.0643, C₁₆H₈N₂O requires *M*, 244.0637). Elution with CH₂Cl₂ gave 3-diazofluoranthen-2(3*H*)-one 14 (236 mg, 20%), mp 153–155.5 °C (decomp.); ν_{max} 2090 (C=N₂) and 1630 cm⁻¹ (C=O); δ 7.75–6.99 (c) (Found: M⁺, 244.0644. C₁₆H₈N₂O requires *M*, 244.0637).

Photolysis of 3-diazofluoranthen-2(3H)-one in benzene

3-Diazofluoranthen-2(3*H*)-one 14 (437 mg, 1.79 mmol) was dissolved in benzene (80 ml) and photolysed for 3 h. The solvent was removed on a rotary evaporator and the dark red oil obtained adsorbed onto silica and chromatographed under low pressure. Elution with petrol–CH₂Cl₂ (60:40) gave a red oil (67 mg) which tlc indicated to be composed of a mixture of at least five components. This was not examined further. Further elution gave a reddish brown oil, which yielded cream-coloured crystals on trituration with petrol, 3-phenylfluoranthen-2-ol 15 (200 mg, 38%), mp 110–111 °C (from petrol–acetone) (Found: C, 89.9; H, 4.8. Calc. for C₂₂H₁₄O: C, 89.75; H, 4.8%), ν_{max}/cm^{-1} 3490 (OH), 1605, 1170, 1147, 777, 756 and 706; $\delta_{\rm H}$ 7.86–7.65 (4 H, c, H-1 and H-4–H-6), 7.53–7.27 (9 H, c, phenyl protons and H-7–H-10), and 5.42 (1 H, br, s, exchanges on shaking with D₂O, OH).

Photolysis of 3-diazofluoranthen-2(3H)-one in benzene-tertbutylamine

3-Diazofluoranthen-2(3H)-one 14 (920 mg, 3.77 mmol) was dissolved in benzene (75 ml) and tert-butylamine (5 ml) and the solution subjected to photolysis for 5.5 h. Removal of the solvent gave a dark green tarry material which was adsorbed onto silica and chromatographed under low pressure. Elution with petrol-CH₂Cl₂ (80:20) gave a fluorescent green solid, 9a,13adihydrobenzo[b]fluorantheno[3,2-d]furan 16 (132 mg, 12%), mp 162.5-164 °C (from toluene) (Found: C, 89.6; H, 4.8. Calc. for $C_{22}H_{14}O$: C, 89.75; H, 4.8%); ν_{max}/cm^{-1} 1225, 1180, 1170, 907, 814, 770, 747 and 690; $\delta_{\rm H}$ 7.86–7.54 (5 H, c, aromatic), 7.51 (1 H, s, H-8). 7.40-7.22 (2 H, c, aromatic), 6.32-5.86 (4 H, c, H-10-H-13), 5.60 (1 H, d of d, J 12, 4, H-9a) and 4.69 (1 H, c d, J 12, H-13a). Increasing the polarity of the solvent (petrol-CH₂Cl₂, 60:40) gave 73 mg of a red oil which tlc indicated was similar to the multi-component mixture described above. Further elution gave 3-phenylfluoranthen-2-ol 15, identical to that described above (257 mg, 23%). Back-flushing the column with ethyl acetate gave 624 mg of black tar.

Photolysis of 2-diazofluoranthen-3(2H)-one in benzene

2-Diazofluoranthen-3(2H)-one 11 (471 mg, 1.93 mmol) was dissolved in benzene (90 ml) and subjected to photolysis for 1 h. Removal of the solvent gave a dark brown oil which was adsorbed onto silica and chromatographed under low pressure. Elution with petrol- $CH_2Cl_2(80:20)$ gave a green solid, solutions of which exhibited green fluorescence, 2-phenylfluoranthen-3ol 12 (419 mg, 74%), mp 140.5-141.5 °C (from ethanol) (Found: C, 89.7; H, 4.65. Calc. for C₂₂H₁₄O: C, 89.75; H, 4.8%), v_{max}/cm^{-1} 3500 and 3400 (OH), 780, 760 and 705; δ_{H} 8.15–7.18 (13 H, br c, aromatic) and 5.92 (1 H, s, exchanges on shaking with D₂O, OH). Spectrum sharpened considerably on cooling to -40 °C, or on shaking with D₂O. $\delta_{\rm H}$ ([²H₆]DMSO) 8.34 (1 H, d, J 8.5, H-4), 8.02-7.84 (4 H, c, aromatic), 7.68-7.20 (8 H, c, aromatic) and 3.6 (1 H, br, s, OH). The spectrum became progressively broader on heating to 160 °C and sharpened on cooling again. EPR spectroscopy (vide infra) indicated that the broadening of the ¹H NMR spectrum was due to the presence of a small concentration of a long-lived free radical species 17.

Photolysis of 2-diazofluoranthen-3(2H)-one in benzene-tertbutylamine

2-Diazofluoranthen-3-(2H)-one 11 (666 mg, 2.73 mmol) was

dissolved in benzene (75 ml) and tert-butylamine (5 ml), and the mixture subjected to photolysis for 4.5 h. Removal of the solvent gave a dark brown tar which was adsorbed onto silica and chromatographed under low pressure. Elution with petrol-CH₂Cl₂ (95:5) gave a yellow solid, 8b,12a-dihydrobenzo-[b]fluorantheno[2,3-d]furan 13 (20 mg, 2.5%), mp 79-83 °C; v_{max} (CHCl₃)/cm⁻¹ 2950, 2920, 2845, 1445, 1260, 1185, 1105 and 1015; $\delta_{\rm H}$ 7.96–7.22 (8 H, c, aromatic), 6.16–5.94 (4 H, c, H-9–H-12), 5.72 (1 H, d of d, J 12, 3, H-12a) and 4.44 (1 H, d, J 12, H-8b) [Found: M⁺, 294.1032(26); (M⁺-2), 292.0876 (100). C₂₂H₁₄O requires M, 294.1045; (M-2) (C₂₂H₁₂O), 292.0888)]. Suitable analytical data could not be obtained from this compound because of the small amount of material available. Increasing the polarity of the solvent to petrol- $CH_2Cl_2(80:20)$ gave 2-phenylfluoranthen-3-ol 12 identical with that obtained previously (426 mg, 53%). Back-flushing the column with ethyl acetate gave a black tar (327 mg).

Photolysis of 9-diazo-8,9-dihydro-4*H*-cyclopenta[*d,e,f*]-phenanthren-8-one in benzene

9-Diazo-8,9-dihydro-4*H*-cyclopenta[*d*,*e*,*f*]phenanthren-8-one **5**⁴ (1.04 g, 4.48 mmol) was dissolved in benzene (190 ml) and the solution subjected to photolysis for 2.5 h. The solvent was removed and the residue adsorbed onto silica and chromatographed under low pressure. Elution with petrol-ether (90:10) gave a pale purple solid, 9-phenyl-4*H*-cyclopenta[*d*,*e*,*f*]phenanthren-8-ol **19** (740 mg, 58.5%), mp 108-109 °C (from ethanol) (Found: C, 89.1; H, 5.1; M⁺, 282.1032. C₂₁H₁₄O requires C, 89.35; H, 5.0%; M⁺, 282.1045); ν_{max} /cm⁻¹ 3490 and 3440 (OH), 1640, 1600, 1435, 1105, 1020, 773 and 702; δ_H 7.97 (1 H, m, H-7), 7.59-7.19 (10 H, c, aromatic), 5.76 (1 H, br s, exchanges on shaking with D₂O, OH) and 4.09 (2 H, br s, CH₂); δ_H ([²H₆]acetone) 8.09 (1 H, m, H-7), 7.90 (1 H, br s, exchanges on shaking with D₂O, OH), 7.63-7.22 (10 H, c, aromatic) and 4.21 (2 H, br s, CH₂).

Photolysis of 9-diazo-8,9-dihydro-4H-

cyclopenta[d,e,f]phenanthren-8-one in benzene-tert-butylamine 9-Diazo-8,9-dihydro-4H-cyclopenta[d,e,f]phenanthren-8-one 5 (973 mg, 4.19 mmol) was dissolved in benzene (90 ml), tertbutylamine (1 ml) was added, and the resulting solution photolysed for 2 h. The solvent was removed and the residue obtained adsorbed onto silica and chromatographed under low pressure. Elution with petrol gave a colourless solid, 8a,12a-dihydro-4Hbenzo[b]cyclopenta[4,4a,4b,5]phenanthro[9,10-d]furan 8 (120 mg, 10%), mp 112–113 °C (from petrol-benzene); v_{max}/cm^{-1} 1160, 1155, 822, 802, 769, 720 and 698; $\delta_{\rm H}$ 7.88–7.26 (6 H, c, aromatic), 6.33-6.08 (3 H, c, H-9-H-11), 5.93 (1 H, m, H-12), 5.70 (1 H, d of d, J 12, 4, H-8a), 4.67 (1 H, c d, J 12, H-12a) and 4.28 (2 H, br s, CH₂) (Found: M⁺, 282.1051. C₂₁H₁₄O requires M⁺, 282.1045). Satisfactory analytical data were not obtained for this compound because of the limited amount of pure material available after repeated recrystallisation. Elution with petrol-ether (90:10) gave 9-phenyl-4H-cyclopenta[d,e,f]phenanthren-8-ol 19 identical with that previously obtained (608 mg, 51%). Further elution gave a red oil (48 mg) which according to tlc contained at least five components and was not examined further. Increasing the polarity of the eluting solvent to petrol-ether (70:30) gave, as a red solid, N-tert-butyl-4,8-dihydrocyclopenta[d,e,f]fluorene-4-carboxamide 10 (33 mg, 3%), mp 124–127 °C (lit.⁴ not reported); v_{max}/cm^{-1} 3250 (NH) and 1640 (C=O); $\delta_{\rm H}$ (60 MHz) 8.0–7.0 (6 H, c, aromatic), 5.22 (1 H, br s, H₄), 4.25 (2, br s, CH₂), 3.99 (1 H, br s, NH) and 1.27 (9 H, s, Bu'); M⁺ 277.

Acid-catalysed rearrangement of 8a,12a-dihydro-4*H*benzo[*b*]cyclopenta[4,4*a*,4*b*,5]phenanthro[9,10-*d*]furan 8

A sample of the dihydrofuran was dissolved in $CDCl_3$ in an NMR tube and a single crystal of *p*-toluenesulfonic acid was added. After 75 min, the signals due to the protons of the di-

hydrofuran ring system had disappeared, and the spectrum indicated that a clean rearrangement to 9-phenyl-4H-cyclopental[d,e,f]phenanthren-8-ol 19 had occurred.

EPR spectroscopy studies

(a) EPR spectrum of radical 17 derived from 2-phenylfluoranthen-3-ol. A solution of 2-phenylfluoranthen-3-ol 12 in toluene (0.05 M) was prepared and a portion of the solution transferred to an EPR tube, degassed and the EPR spectrum observed at ambient temperature [Fig. 1(a)]. A strong signal was obtained which, when resolved, was found to be rather complex, consisting of a symmetrical multiplet centred on $g = 2.0072 \pm 0.0002$). The smallest observed splitting was 0.58 (± 0.02) mT and the total width of the spectrum was 15.5 (± 0.1) mT (total number of lines >28). By comparison of the intensity of the signal with those of signals obtained from solutions of diphenylpicrylhydrazyl (DPPH) of known concentrations, it was possible to estimate that at room temperature the radical species present constituted between 0.4 and 0.8% of the solute. The concentration of the radical was also shown to be temperature dependent, the intensity of the signal decreasing considerably on cooling to -50 °C and returning to its original level on returning to room temperatre.

(b) EPR spectrum of radical 18 derived from 2-phenyl-[²H₃]fluoranthen-3-ol. 2-Diazofluoranthen-3(2*H*)-one 11 (60 mg, 0.25 mmol) was dissolved in [²H₆]benzene (2 ml), which had been dried over a molecular sieve, and the solution placed in an NMR tube. The tube was attached to the cooling jacket of a photochemical reactor and the solution photolysed for 3 h, after which time the NMR spectrum indicated that 2-phenyl[²H₃]fluoranthen-3-ol had been produced. The solution was chromatographed over silica and eluted with petrol-CH₂CL₂ (50:50) to give a red oil. A repeat examination by ¹H NMR spectroscopy showed that exchange of the *O*-deuterium for hydrogen had occurred to give 2-phenyl[²H₃]fluoranthen-3-ol (48 mg, 65%) [$\delta_{\rm H}$ 8.4-7.0 (H, br c, aromatic) and 5.04 (1 H, br s, OH)] and that the spectrum sharpened considerably on cooling to -60 °C. A solution of this material in toluene (0.1 M) was prepared and a portion transferred to an EPR tube, degassed and its spectrum recorded. A strong symmetrical signal, consisting of five broad lines with the same centre as the EPR spectrum already described, was obtained [Fig. 1(*b*)].

Acknowledgements

We thank Mr A. P. Anderson for his assistance in EPR measurements. We also thank Dr A. D. McNaught of the Royal Society of Chemistry Information Services for helpful comments in connection with the naming of these compounds.

References

- I C. F. Bannerman, P. J. N. Brown, J. I. G. Cadogan, I. Gosney, A. Johnstone and N. H. Wilson, J. Chem. Soc., Chem. Commun., 1975, 618.
- 2 For a review on the Wolff rearrangement, including mechanism and competing reactions, see G. B. Gill, in *Comprehensive Organic Synthesis*, ed. B. M. Trost, I. Fleming and G. Pattenden, Pergamon Press, Oxford, 1991, vol. 3, p. 887.
- 3 See, W. Reid and H. Lohwasser, Liebigs Ann. Chem., 1965, 683, 118, for details of the failure of 2-diazoacenaphthen-1-one 1 to undergo Wolff rearrangement on thermolysis, instead giving the corresponding ketazine.
- 4 B. M. Trost and P. L. Kinson, J. Am. Chem. Soc., 1975, 97, 2438.
- 5 M. Periasamy and M. V. Bhatt, Synthesis, 1977, 330.
- 6 W. E. Barth and R. G. Lawton, J. Am. Chem. Soc., 1971, 93, 1730.
- 7 J. Douris and A. Mathieu, Bull. Soc. Chim. Fr., 1971, 3365.
- 8 M. E. Alonso, A. Morales and A. W. Chitty, J. Org. Chem., 1982, 47, 3747.
- 9 R. M. Paton and R. U. Weber, Org. Magn. Res., 1977, 9, 494.
- 10 W. Reid and R. Dietrich, Chem. Ber., 1961, 94, 387.

Paper 6/034961 Received 20th May 1996 Accepted 10th July 1996