

Palladium-Catalyzed Suzuki–Miyaura Cross-Couplings of Sulfonyl Chlorides and Boronic Acids


Srinivas Reddy Dubbaka and Pierre Vogel*

Institute of Molecular and Biological Chemistry, Swiss Institute of Technology,
BCH CH-1015 Lausanne, Switzerland

pierre.vogel@epfl.ch

Received October 31, 2003

ABSTRACT




Arene-, arylmethane, and alk-2-ene-1-sulfonyl chlorides undergo Suzuki–Miyaura cross-coupling with arene-, heteroarene-, and alkeneboronic acids in THF at reflux. The reactivity order is ArI > ArSO₂Cl > ArBr >> ArCl.

The palladium-catalyzed Suzuki–Miyaura reaction has emerged as an extremely powerful method for the cross-coupling of aryl bromides, iodides, and triflates with boronic acids.¹ Coupling between electron-rich or electron-neutral aryl chlorides and arene boronic acids is also possible by using palladium complexes with sterically hindered, electron-rich, or carbene ligands such as those developed by Buchwald,² Fu,³ and Herrmann,⁴ respectively. Aryl arenesulfonates can cross-couple with arylboronic acids.⁵ Vinylation,⁶ carbonylation⁷ and homocoupling⁸ of arenesulfonyl chlorides have already been described. Recently, we have disclosed

the palladium-catalyzed Stille cross-coupling of sulfonyl chlorides and organostannanes.⁹ We are reporting here that arene-, phenylmethane-, and 2-methylprop-2-enesulfonyl chlorides can be condensed with arene-, heteroarene-, and alkeneboronic acids in the presence of a palladium catalyst (Scheme 1).

Scheme 1. Cross-Coupling of Sulfonyl Chlorides with Boronic Acids



In our first search for optimal reaction conditions, we explored the condensation of *p*-toluenesulfonyl chloride

(1) For reviews, see for example: (a) Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, *95*, 2457–2483. (b) Littke, A. F.; Fu, G. C. *Angew. Chem., Int. Ed.* **2002**, *41*, 4176–4211. (c) Hassan, J.; Sévignon, M.; Gozzi, C.; Schulz, E.; Lemaire, M. *Chem. Rev.* **2002**, *102*, 1359–1469. (d) Kotha, S.; Lahiri, K.; Kashinath, D. *Tetrahedron* **2002**, *58*, 9633–9695. (e) Agrofoglio, L. A.; Gillaizeau, I.; Saito, Y. *Chem. Rev.* **2003**, *103*, 1875–1916.

(2) (a) Wolfe, J. P.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **1999**, *38*, 2413–2416. (b) Wolfe, J. P.; Singer, R. A.; Yang, B. H.; Buchwald, S. L. *J. Am. Chem. Soc.* **1999**, *121*, 9550–9561.

(3) (a) For a recent review, see: Littke, A. F.; Fu, G. C. *Angew. Chem., Int. Ed.* **2002**, *41*, 4176–4211. (b) Littke, A. F.; Fu, G. C. *Angew. Chem., Int. Ed.* **1998**, *37*, 3387–3388.

(4) (a) For recent review, see: Herrmann, W. A. *Angew. Chem., Int. Ed.* **2002**, *41*, 1290–1309. (b) Hillier, A. C.; Grasa, G. A.; Viciu, M. S.; Lee H. M.; Yang, C.; Nolan S. P. *J. Organomet. Chem.* **2002**, *653*, 69–82. (c) Herrmann, W. A.; Reisner, C. P.; Spieger, M. *J. Organomet. Chem.* **1998**, *557*, 93–96. (d) Zhang, C.; Huang, J.; Trudell, M. L.; Nolan S. P. *J. Org. Chem.* **1999**, *64*, 3804–3805.

(5) (a) Percec, V.; Bae, J.-Y.; Hill, D. H. *J. Org. Chem.* **1995**, *60*, 1060–1065. (b) Nguyen, H. N.; Huang, X.; Buchwald, S. L. *J. Am. Chem. Soc.* **2003**, *125*, 11818–11819.

(6) (a) Kasahara, A.; Izumi, T.; Kudou, N.; Azami, H.; Yamamoto, S. *Chem. Ind. (London)* **1988**, 51–52. (b) Miura, M.; Hashimoto, H.; Itoh, K.; Nomura, M. *Tetrahedron Lett.* **1989**, *30*, 975–976. (c) Miura, M.; Hashimoto, H.; Itoh, K.; Nomura, M. *J. Chem. Soc., Perkin. Trans. 1* **1990**, *8*, 2207–2211.

(7) Miura, M.; Itoh, K.; Nomura, M. *Chem. Lett.* **1989**, 77–78.


(8) Miura, M.; Hashimoto, H.; Itoh, K.; Nomura, M. *Chem. Lett.* **1990**, 459–460.

(9) Dubbaka, S. R.; Vogel, P. *J. Am. Chem. Soc.* In press.

(TsCl) with 3-nitrobenzeneboronic acid in the presence of various catalysts such as $\text{Pd}(\text{PPh}_3)_4$, $\text{PdCl}_2(\text{PhCN})_2$, $\text{Pd}_2(\text{dba})_3 + \text{P}(t\text{-Bu})_3$, $\text{Pd}_2(\text{dba})_3 + \text{tri-2-furylphosphine}$, and $\text{Pd}_2(\text{dba})_3 + 2\text{-(di-tert-butylphosphino)biphenyl}$ in boiling DME, DMF, THF, or dioxane.¹⁰ Cross-coupling occurred only in the presence of a base such as K_2CO_3 , Cs_2CO_3 , K_3PO_4 , or Et_3N . The best yield (60%) was obtained using 8 mol % $\text{Pd}(\text{PPh}_3)_4$ and 2.0–3.0 equiv of K_2CO_3 .

Then, we searched to optimize the proportion of palladium employed, screening different sterically hindered, electron-rich, or carbene ligands,¹¹ as well as bases and solvents (Table 1 and Figure 1) for 1-naphthalenesulfonyl chloride reacting with 4-methylbenzeneboronic acid.

Table 1. Screening of Bases for Suzuki–Miyaura Cross-Coupling of Sulfonyl Chloride and Boronic Acid



entry	base	yield ^a
1	Et_3N	10
2	NaOAc	5
3	Cs_2CO_3	0
4	K_3PO_4	19
5	K_2CO_3	41
6	Na_2CO_3	60
7	Li_2CO_3	28

^a Yields of cross-coupling products determined after flash chromatography.

We found that Na_2CO_3 is a better activator for the boronic acid (without destroying the sulfonyl chloride) compared to other bases such as Et_3N , Cs_2CO_3 , NaOAc , K_3PO_4 , K_2CO_3 , or Li_2CO_3 . When using Cs_2CO_3 as a base, the formation of the cross-coupling product was not observed, but all the starting sulfonyl chloride was consumed. When NaOAc was used as a base, the cross-coupling product was isolated in low yield (6%). Under these conditions, starting sulfonyl chloride was consumed in ca. 5 h. This is explained by the fact that the sulfonyl chloride is converted into the corresponding sulfonate, which retards the reaction.

A small library of ligands was tested, and it was found that the carbene derived by HCl α -elimination from **7** generates, with palladium, the best catalyst (Figure 1).

A wide range of electronically and structurally diverse sulfonyl chlorides and boronic acids can be cross-coupled efficiently under these conditions.¹² Thus, with respect to the electron-rich, electron-neutral, and electron-poor arene-sulfonyl chlorides, they react with the boronic acids to provide the corresponding biaryl products in good yields in most cases (Table 2). Self-coupling of boronic acids was






Figure 1. Screening of ligands for Suzuki–Miyaura cross-coupling of sulfonyl chloride and boronic acid.

observed in some cases as a concurrent reaction.¹³ The proportion of the latter was lower in boiling THF than in boiling dioxane, acetonitrile, 1,2-dimethoxyethane, or *p*-xylene. When $\text{Pd}(\text{PPh}_3)_4$ was used as a catalyst, traces of diarylsulfide were frequently observed. For instance, treatment of *p*-toluenesulfonyl chloride with 4-methoxybenzeneboronic acid in the presence of $\text{Pd}(\text{PPh}_3)_4$ and K_2CO_3 (3 equiv) led to Tol-S-Ar (Ar = 4-methoxyphenyl) that was isolated in 8% yield. This byproduct results probably from the reduction of the sulfonyl chloride into the corresponding sulfenyl chloride,^{9,14} which is then coupled with the boronic acid following the Suzuki–Miyaura catalytic cycle (Figure

(12) **Typical Experimental Procedure for Catalyst B (Table 3).** A round-bottom flask was dried under vacuum, and to the flask were added the corresponding sulfonyl chloride (1.00 mmol), Pd_2dba_3 (0.015 mmol), ligand **7** (0.06 mmol), Na_2CO_3 (3 mmol), and the boronic acid (2 mmol) (were weighed in a glovebox) under a nitrogen atmosphere. Then, the flask was connected to a vacuum line and then filled with argon (three times), and THF (5 mL) was added under argon. The reaction mixture was stirred at reflux for 15–35 h. After cooling to room temperature, the mixture was diluted with ether and washed with water. The aqueous layer was extracted again with ether (three times). The combined organic phases were dried (Na_2SO_4), filtered, and concentrated under reduced pressure (solvents were removed under reflux on cooling to -20°C in the cases of compounds with low boiling points or low molecular masses). The residue was purified by FC.

(13) Mañas, M. M.; Perez, M.; Pleixats, R. *J. Org. Chem.* **1996**, *61*, 2346–2351.

(14) Oae, S.; Togo, H. *Bull. Chem. Soc. Jpn.* **1983**, *56*, 3802–3812.

(10) See Supporting Information for screening of catalysts and solvents.

(11) All ligands were bought from Strem Chemicals, Inc.

Table 2. Palladium-Catalyzed Suzuki–Miyaura Cross-Couplings of Sulfonyl Chlorides and Boronic Acids

$\text{R-SO}_2\text{Cl} + \text{R}^1\text{-B(OH)}_2 \xrightarrow[\text{THF, reflux, 15–35 h}]{\text{Palladium Catalyst, Base (3 equiv)}} \text{R-R}^1$ $1.0 \text{ equiv} \quad 2.0 \text{ equiv}$			
entry	R	R ¹	yield* cat. A ^a /cat. B ^b
1	4-methylphenyl	phenyl	55
2	4-methylphenyl	4-methylphenyl	70 92
3	4-methylphenyl	3-nitrophenyl	60
4	4-methylphenyl	3-formylphenyl	55
5	4-methylphenyl	4-methoxyphenyl	50 65
6	4-methylphenyl	fur-2-yl	40
7	4-methylphenyl	(<i>E</i>)-2-phenylvinyl	35 48
8	4-methylphenyl	4-chlorophenyl	60 78
9	4-methylphenyl	1-naphthyl	64 76
10	4-chlorophenyl	3-nitrophenyl	65
11	4-chlorophenyl	4-methylphenyl	62 80
12	1-naphthyl	3-formylphenyl	55 72
13	1-naphthyl	3-nitrophenyl	61
14	1-naphthyl	4-methylphenyl	55 82
15	1-naphthyl	4-methylphenyl	58 ^c 14 ^d
16	1-naphthyl	4-methylphenyl	trace ^e
17	1-naphthyl	4-methylphenyl	42 ^f
18	3-nitrophenyl	3-nitrophenyl	78
19	3-nitrophenyl	4-methylphenyl	70
20	4-nitrophenyl	4-methylphenyl	64
21	benzyl	4-methylphenyl	30
22	benzyl	3-nitrophenyl	35 52
23	methallyl	3-nitrophenyl	35 50

* Catalyst A: 8–10 mol % Pd(PPh₃)₄ and 2–3 equiv of K₂CO₃. Catalyst B: 1.5 mol % Pd₂dba₃, 6 mol % ligand **7**, and 3 equiv of Na₂CO₃. ^a R–S–R was isolated as a side product. ^b R–R was isolated as a side product. ^c Catalyst A and activated charcoal. ^d Catalyst B with 1.0 equiv of TBAB. ^e Performed with 1.5 mol % Pd(OH)₂/C and Na₂CO₃ as a base. ^f Performed with 5 mol % Pd(OAc)₂, 10 mol % PPh₃, and K₂CO₃ as a base.

2). In contrast, when using carbene, sterically hindered, or electron-rich ligands, the arylpalladium chloride intermediates can give the corresponding homocoupled product¹⁵ (Figure 2).

After screening the different ligands (**1–9**) (Figure 1), we have found that catalyst B (although not tested for all the




Figure 2. Probable catalytic cycle for Suzuki–Miyaura cross-coupling of sulfonyl chlorides and boronic acids.

reactions described here) was more efficient than the classical catalyst A (Table 2).


Under nonbasic conditions, the cross-coupling reaction was more efficient in the presence of CuCl (3 equiv) than with CuBr·Me₂S. With 3 equiv of CuI, no reaction was observed.¹⁰

With *p*-halogenobenzenesulfonyl chlorides **10**, **13**, and **17**, we have observed that the sulfonyl chloride group is more reactive than the bromo- and chloroarene moieties (Scheme 2).¹⁶ However, iodoarenes are more reactive than arenesulfonyl chlorides.

Recently, Leadbeater and Marco have reported transition-metal-free Suzuki-type coupling reactions using tetrabutylammonium bromide (TBAB) as a phase transfer reagent.¹⁷ We have applied these conditions to our sulfonyl chlorides, but without success.¹⁸ When the reaction is performed in the presence of a palladium catalyst and TBAB, the corresponding cross-coupling product is obtained in a lower yield than in the absence of TBAB (Table 2, entry 15).

When TsCl was mixed with 1 equiv of PdCl₂(PhCN)₂ in THF-*d*₈, a signal (δ_c 145.2 ppm) typical of the C–Pd bond¹⁹

Scheme 2. Reactivity Order for Suzuki–Miyaura Cross-Coupling Is ArI > ArSO₂Cl > ArBr ≫ ArCl



appeared in the ^{13}C NMR spectra after 10 min at 45 °C. At least two mechanisms can be envisaged for this observation: (1) the metal inserts into the $\text{SO}_2\text{--Cl}$ bond first with rapid subsequent elimination of SO_2 (Figure 2), or (2) the oxidative addition of the palladium occurs first onto the C--S bond, with subsequent elimination of SO_2 .

We also examined nickel-catalyzed Suzuki–Miyaura cross-coupling reactions between various sulfonyl chlorides and areneboronic acids in the presence of a base, but none of our assays have been met with success.²⁰

In summary, a palladium-catalyzed cross-coupling reaction between arene-, phenylmethane-, 2-methylprop-2-enesulfonyl

chlorides, and arene-, heteroarene-, and alkeneboronic acids is possible. The reaction provides a useful complement to our recently described work of Stille cross-couplings of sulfonyl chlorides and organostannanes⁹ and should open new possibilities for medicinal chemistry and material science.

Acknowledgment. We thank the Swiss National Science Foundation (Grant No. 2000-20100002/1) and the Office Federal de l'Éducation et de la Sciences, (Grant No. COI. 0071, COST D13/010/01). We thank also Mr. Martial Rey and Francisco Sepúlveda for technical help.

Supporting Information Available: Experimental procedures, characterization of unknown compounds, and references to known compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL036131X

(15) Clark, F. R. S.; Norman, R. O. C.; Thomas, C. B. *J. Chem. Soc., Perkin Trans. 1* **1975**, 121–125.

(16) For more details, see also ref 9.

(17) Leadbeater, N. E.; Marco, M. *J. Org. Chem.* **2003**, 68, 5660.

(18) Acetonitrile was used as a solvent instead of water, and K_2CO_3 was used as a base.

(19) Bergbreiter, D. E.; Osburn, P. L.; Wilson, A.; Sink, E. M. *J. Am. Chem. Soc.* **2000**, 122, 9058–9064.

(20) See Supporting Information for reaction details.