

Halopalladation/Decarboxylation/ Carbon–Carbon Forming Domino Process: Synthesis of 5-Halo-6-substituted Benzo[*b*]naphtho[2,1-*d*]furans


Xiao-Cheng Huang,[†] Feng Wang,[†] Yun Liang,^{*,†} and Jin-Heng Li^{*,†,‡}

Key Laboratory of Chemical Biology & Traditional Chinese Medicine Research
(Ministry of Education), Hunan Normal University, Changsha 410081, China, and
College of Chemistry and Materials Science, Wenzhou University,
Wenzhou 325035, China

liangyun1972@126.com; jhli@hunnu.edu.cn

Received December 28, 2008


ABSTRACT



A novel and general halopalladation/decarboxylation/carbon–carbon forming domino protocol is described for the synthesis of 5-halo-6-substituted benzo[*b*]naphtho[2,1-*d*]furans. The protocol represents the first example of trapping the σ -vinylpalladium intermediate, generated from halopalladation of alkynes, by the decarboxylative coupling reaction.

Halopalladation of alkynes has been proven to be an extremely important and convenient method for the construction of both the carbon–carbon and carbon–halide bonds in one step.^{1–4} Halopalladation of alkynes will generate a versatile reactive σ -vinylpalladium intermediate **A** (Scheme 1).¹ Until now, how to capture the σ -vinylpalladium intermediate is still a highly challenging area because the capture reagents are often limited to allyl halides, alkynes, conjugated

Scheme 1. Mechanism of Halopalladation of Alkynes with Allyl Halides



dienes, and carbon monoxide. Recently, we found that the σ -vinylpalladium intermediate could be trapped by some nucleophilic *ortho*-groups in the arylalkyne moiety.³ This prompted us to explore other nucleophilic groups for the purpose. Accidentally, we found that arylcarboxylates were

[†] Hunan Normal University.

[‡] Wenzhou University.

(1) For reviews, see: (a) Heck, R. F. *Palladium Reagents in Organic Synthesis*; Academic Press: New York, 1985. (b) Tsuji, J. *Palladium Reagents and Catalysts: Innovations in Organic Synthesis*; Wiley and Sons: New York, 1995. (c) Li, J. J.; Gribble, G. W. *Palladium in Heterocyclic Chemistry*; Pergamon: New York, 2000. (d) Lu, X. *Handbook of Organopalladium Chemistry for Organic Synthesis*; Negishi, E., Ed.; Wiley-Interscience: New York, 2002; Vol. 2, pp 2267–2288. (e) Han, X. L.; Liu, G. X.; Lu, X. Y. *Chin. J. Org. Chem.* **2005**, *25*, 1182. (f) Tang, S.; Wang, N.-X.; Li, J.-H. *Chin. J. Org. Chem.* **2007**, *27*, 819.

(2) Phillips, F. C. Z. *Anorg. Allg. Chem.* **1894**, *6*, 229.

(3) (a) Liang, Y.; Tang, S.; Zhang, X.-D.; Mao, L.-Q.; Xie, Y.-X.; Li, J.-H. *Org. Lett.* **2006**, *8*, 3017. (b) Tang, S.; Xie, Y.-X.; Li, J.-H.; Wang, N.-X. *Synthesis* **2007**, 1841. (c) Tang, S.; Yu, Q.-F.; Peng, P.; Li, J.-H.; Zhong, P.; Tang, R.-Y. *Org. Lett.* **2007**, *9*, 3413.

an available alternative to the traditional capture reagents to trap the σ -vinylpalladium complexes.⁵ In the presence of PdCl₂ and CuX₂ (X = Cl, Br), a variety of 2-(2-(2-substituted ethynyl)phenyl)benzofuran-3-carboxylates could undergo an intramolecular decarboxylative coupling reaction to capture the σ -vinylpalladium intermediate in situ. In this paper, we report our preliminary results on the first example of trapping the σ -vinylpalladium intermediate by the decarboxylative coupling reaction (eq 1). It is noteworthy that the products, polycyclic aromatic furans, are prominent structural units found in a wide range of natural products, pharmaceuticals, and materials as well as important intermediates in organic synthesis.⁶

Ethyl 2-(2-(2-phenylethynyl)phenyl)benzofuran-3-carboxylate (**1a**) was treated with PdCl₂ and CuCl₂ to screen the optimal reaction conditions, and the results are summarized in Table 1. Initially, a set of solvents were examined, and benzene was found to be the most effective solvent (entries 1–6). Treatment of substrate **1a** with PdCl₂ and CuCl₂ in benzene afforded the corresponding target product **2a** in a 69% yield (entry 1). Identical results were obtained using toluene as the solvent (entry 2). However, other solvents, such as ClCH₂CH₂Cl, CH₃CN, THF, and cyclohexane, decreased the yield to some extent (entries 3–6). Subsequently, the amount of CuCl₂ was tested (entries 7–9). We found that 1 equiv of CuCl₂ reduced the yield to 55% (entry 7), and 5 equiv of CuCl₂ provided the same yield as that of 3 equiv of CuCl₂ (entry 8). However, a trace amount of **2a** was observed without either CuCl₂ or PdCl₂ (entries 9 and 11). It is worth noting that no reaction is observed using LiCl instead of CuCl₂ (entry 10), and room temperature gives a low yield after 24 h (entry 12). The screening results demonstrated that two analogs, *i*-propyl 2-(2-(2-phenylethynyl)phenyl)benzofuran-3-carboxylate (**1b**) or benzyl 2-(2-(2-phenylethynyl)phenyl)benzofuran-3-carboxylate (**1c**), were also suitable for the halopalladation/decarboxylation/carbon–carbon forming reaction with PdCl₂ and CuCl₂ in moderate yields (entries 13 and 14). To our delight, the reaction of substrate **1a** with PdBr₂ and CuBr₂ was conducted successfully in MeCN to afford the corresponding 5-bromo-6-phenylbenzo[*b*]naphtho[2,1-*d*]furan (**3a**) in 59% yield (entry 16).

(4) For other papers on the halopalladation reactions using PdX₂/CuX₂ as a catalytic system, see: (a) Bäckvall, J. E.; Nordberg, R. E. *J. Am. Chem. Soc.* **1980**, *102*, 393. (b) Ji, J.; Zhang, C.; Lu, X. *J. Org. Chem.* **1995**, *60*, 1160. (c) Ma, S.; Lu, X. *J. Org. Chem.* **1993**, *58*, 1245. (d) Li, J.-H.; Jiang, H.-F.; Feng, A.-Q.; Jia, L.-Q. *J. Org. Chem.* **1999**, *64*, 5984. (e) Li, J.-H.; Jiang, H.-F.; Chen, M.-C. *J. Org. Chem.* **2001**, *66*, 3627. (f) Li, J.-H.; Liang, Y.; Xie, Y.-X. *J. Org. Chem.* **2004**, *69*, 8125. (g) Li, J.-H.; Tang, S.; Xie, Y.-X. *J. Org. Chem.* **2005**, *70*, 477. (h) Ma, S.; Wu, B.; Zhao, S. *Org. Lett.* **2003**, *5*, 4429. (i) Ma, S.; Wu, B.; Jiang, X.; Zhao, S. *J. Org. Chem.* **2005**, *70*, 2568. (j) Ma, S.; Wu, B.; Jiang, X. *J. Org. Chem.* **2005**, *70*, 2588. (k) Huang, J.; Zhou, L.; Jiang, H. *Angew. Chem., Int. Ed.* **2006**, *45*, 1945.

(5) For selected reviews and papers on Pd-catalyzed decarboxylative coupling reactions, see: (a) Goossen, L. J.; Goossen, K.; Rodríguez, N.; Blanchot, M.; Linder, C.; Zimmermann, B. *Pure Appl. Chem.* **2008**, *80*, 1725. (b) Goossen, L. J.; Rodríguez, N.; Goossen, K. *Angew. Chem., Int. Ed.* **2008**, *47*, 3100, and references cited therein. (c) Tatamidani, H.; Yokota, K.; Kakiuchi, F.; Chatani, N. *J. Org. Chem.* **2004**, *69*, 5615. (d) Goossen, L. J.; Knauber, T. *J. Org. Chem.* **2008**, *73*, 8631. (e) Goossen, L. J.; Zimmermann, B.; Knauber, T. *Angew. Chem., Int. Ed.* **2008**, *47*, 7103. (f) Goossen, L. J.; Rodríguez, N.; Linder, C. *J. Am. Chem. Soc.* **2008**, *130*, 15248. (g) Goossen, L. J.; Rudolph, F.; Oppel, C.; Rodríguez, N. *Angew. Chem., Int. Ed.* **2008**, *47*, 3043. (h) Maehara, A.; Tsurugi, H.; Satoh, T.; Miura, M. *Org. Lett.* **2008**, *10*, 1259. (i) Nakano, M.; Tsurugi, H.; Satoh, T.; Miura, M. *Org. Lett.* **2008**, *10*, 1851. (j) Forgiione, P.; Brochu, M.-C.; St-Onge, M.; Thesen, K. H.; Bailey, M. D.; Bilodeau, F. *J. Am. Chem. Soc.* **2006**, *128*, 11350.


(6) (a) Harvey, R. G. *Polycyclic Aromatic Hydrocarbons*; Wiley-VCH: New York, 1996. (b) Harvey, R. G. In *Handbook of Environmental Chemistry*; Hutzinger, D., Neilson, A., Eds.; Springer: Berlin, Heidelberg, 1997; Vol. 3, part 1, ch. 1. (c) Watson, M. D.; Fechtenkötter, A.; Mullen, K. *Chem. Rev.* **2001**, *101*, 1267. (d) Law, K. Y. *Chem. Rev.* **1993**, *93*, 449. (e) Jung, M. E. *Tetrahedron* **1976**, *32*, 3. (f) Molander, G. A. *Acc. Chem. Res.* **1998**, *31*, 603. (g) Tiwari, S. S.; Srivastava, S. C. *J. Med. Chem.* **1967**, *10*, 983.

Table 1. Screening Optimal Conditions^a

entry	R	CuX ₂ (equiv)	solvent	t (h)	yield (%) ^b
1	Et (1a)	CuCl ₂ (3)	benzene	16	69 (2a)
2	Et (1a)	CuCl ₂ (3)	toluene	23	70 (2a)
3	Et (1a)	CuCl ₂ (3)	ClCH ₂ CH ₂ Cl	6	60 (2a)
4	Et (1a)	CuCl ₂ (3)	CH ₃ CN	24	20 (2a)
5	Et (1a)	CuCl ₂ (3)	THF	16	24 (2a)
6	Et (1a)	CuCl ₂ (3)	cyclohexane	23	54 (2a)
7	Et (1a)	CuCl ₂ (1)	benzene	24	55 (2a)
8	Et (1a)	CuCl ₂ (5)	benzene	16	70 (2a)
9	Et (1a)	—	benzene	24	trace (2a)
10	Et (1a)	LiCl (3)	benzene	16	0 (3a)
11 ^c	Et (1a)	CuCl ₂ (3)	benzene	24	trace (2a)
12 ^d	Et (1a)	CuCl ₂ (3)	benzene	24	16 (2a)
13	<i>i</i> -Pr (1b)	CuCl ₂ (3)	benzene	16	59 (2a)
14	Bn (1c)	CuCl ₂ (3)	benzene	16	65 (2a)
15	Et (1a)	CuBr ₂ (3)	benzene	16	35 (3a)
16 ^e	Et (1a)	CuBr ₂ (3)	MeCN	16	59 (3a)


^a Reaction conditions: **1** (0.3 mmol), PdCl₂ (10 mol %), and CuCl₂ (3 equiv) in solvent (5 mL) at 100 °C. ^b Isolated yield. ^c Without PdCl₂. ^d At room temperature. ^e PdBr₂ (10 mol %) instead of PdCl₂.

nyl)phenyl)benzofuran-3-carboxylate (**1b**) or benzyl 2-(2-(2-phenylethynyl)phenyl)benzofuran-3-carboxylate (**1c**), were also suitable for the halopalladation/decarboxylation/carbon–carbon forming reaction with PdCl₂ and CuCl₂ in moderate yields (entries 13 and 14). To our delight, the reaction of substrate **1a** with PdBr₂ and CuBr₂ was conducted successfully in MeCN to afford the corresponding 5-bromo-6-phenylbenzo[*b*]naphtho[2,1-*d*]furan (**3a**) in 59% yield (entry 16).



As shown in Table 2, we examined the scope of the halopalladation/decarboxylation/carbon–carbon bond forming process with respect to ethyl carboxylate substrates under the standard conditions. Substituents at the terminal alkynic moiety of 2-(2-(ethynyl)phenyl)benzofuran-3-carboxylates were first evaluated in the presence of PdX₂ and CuX₂ (X = Cl, Br) (entries 1–13). While carboxylates **1d–1f** bearing electron-rich aryl groups were treated with PdX₂ and CuX₂ (X = Cl or Br) in good yields (entries 1–6), substrates **1g–1k**, having either electron-deficient aryl or alkyl groups, reduced the yields to some extent (entries 7–12). Ethyl 2-(2-(*o*-tolylethynyl)phenyl)benzofuran-3-carboxylate (**1e**), for instance, underwent the tandem reaction with PdCl₂ and

Table 2. Halopalladation/Decarboxylation/Carbon–Carbon Forming Reactions of **1** with PdX₂ and CuX₂^a




entry	substrate 1	<i>t</i> (h)	CuX ₂	yield (%) ^b
1		10	CuCl ₂	76 (2d)
2 ^c		10	CuCl ₂	76 (2d)
3		16	CuCl ₂	82 (2e)
4 ^d		4	CuBr ₂	89 (3e)
5		18	CuCl ₂	73 (2f)
6 ^d		5	CuBr ₂	79 (3f)
7		16	CuCl ₂	48 (2g)
8		7	CuCl ₂	47 (2h)
9		5.5	CuCl ₂	58 (2i)
10 ^c		5.5	CuCl ₂	64 (2i)
11		5.5	CuCl ₂	52 (2j)
12		4	CuCl ₂	15 (2k)
13		18	CuCl ₂	51 (2l)
14		30	CuCl ₂	69 (2m)
15		12	CuCl ₂	42 (2n)
16		7	CuCl ₂	45 (2o)
17		7	CuCl ₂	46 (2p)
18		12	CuCl ₂	70 (2q)
19		9	CuCl ₂	57 (2r)

^a Reaction conditions: **1** (0.3 mmol), PdCl₂ (10 mol %), CuCl₂ (3 equiv) in benzene (5 mL) at 100 °C. ^b Isolated yield. ^c In toluene. ^d PdBr₂ (10 mol %) and CuBr₂ (3 equiv) in MeCN (5 mL).

CuCl₂ in 82% yield (entry 3) and with PdBr₂ and CuBr₂ in 89% yield (entry 4). The reaction of substrate **1g**, bearing an electron-deficient NO₂ group, with PdCl₂ and CuCl₂ afforded the target product **2g** in only 48% yield (entry 7). The alkyl groups, including octyl, hexyl, and 2-cyclohexenyl at the alkyne moiety, also provided the corresponding products in moderate yields (entries 8–11), and the hy-


droxymethyl groups resulted in a low yield (entry 12). It is noted that the reactions of substrates **1d** or **1i** can be conducted efficiently in toluene (entries 2 and 10). Gratifyingly, the standard conditions were compatible with the thiophen-2-yl group, providing a moderate yield (entry 13). It was found that two groups, OMe and F, on the 2-aryl moiety were perfectly tolerated under the standard conditions (entries 14–17). Ethyl 2-(5-methoxy-2-(2-phenylethynyl)phenyl)benzofuran-3-carboxylate (**1m**), for example, underwent the reaction with PdCl₂ and CuCl₂ smoothly in 69% yield (entry 14). Remarkably, naphtho[2,1-*b*]furan-1-carboxylates **1q** and **1r** were suitable substrates providing good yields under the same conditions (entries 18 and 19).

The reaction of 2-(2-(2-phenylethynyl)phenyl)-benzofuran (**3**) was also tested under the standard conditions, and a mixture was observed by GC-MS analysis (eq 2). In the presence of PdCl₂ and CuCl₂, 30% of substrate **3** was consumed after 24 h to provide a mixture of the desired product **2a** and another hydroarylation product **4** in 16% total yield.



Two possible mechanisms as outlined in Scheme 2 were proposed for the present reaction on the basis of the previously reported mechanism.^{1–5} Complexation of PdX₂ with the carbon–carbon triple bond affords intermediate **B**, followed by *trans*-addition to give intermediate **C**, the σ -vinylpalladium intermediate.^{1–3} Intermediate **C** undergoes

Scheme 2. A Working Mechanism



anion exchange with Cu to produce intermediate **D**. Intermediate **E** is formed from decarboxylation of intermediate **D**.⁵ Subsequently, intramolecular transmetalation of intermediate **E** takes place to give intermediate **F**. Finally, reductive elimination of intermediate **F** yields both the target product and Pd(0) species. The Pd(0) species are oxidized by CuX₂ to regenerate the active PdX₂ species. Another possible pathway is a Heck-like mechanism. Insertion of the vinylpalladium species **C** into benzofuran via a Heck-like process forms intermediate **G**, which subsequently undergoes decarboxylative β -carbo-elimination to give the desired product.

We deduced that three roles of CuX₂ are played in the process: (1) as a halide source; (2) as a cocatalyst to react with carbonyls generating the Cu intermediate in the decarbonylation process; and (3) as a cocatalyst to cleave the C–Pd σ -bond and regenerate the active PdX₂ species in the halopalladation reaction. Study of the real function of CuX₂ is in progress.

In summary, we have developed a new and general protocol for the synthesis of polycyclic aromatic furans by a halopalladation/decarbonylation/carbon–carbon forming

domino process. In the presence of PdCl₂ and CuX₂, a variety of 2-(2-(2-substitutedethynyl)phenyl)benzofuran-3-carboxylates underwent the reaction to afford 5-halo-6-substituted benzo[*b*]naphtho[2,1-*d*]furans in moderate to good yields. It is noteworthy that these products with a halide (Cl or Br) provide an attractive and useful route to introduce new groups for the synthesis of new natural products. Work to apply the reaction in organic synthesis is currently underway.

Acknowledgment. We thank the Specialized Research Fund for the Doctoral Program of Higher Education (No. 20060542007), National Natural Science Foundation of China (No 20572020), New Century Excellent Talents in University (No. NCET-06-0711), and Fok Ying Tung Education Foundation (No. 101012) for financial support.

Supporting Information Available: Analytical data and spectra (¹H and ¹³C NMR) for all the products **2** and **3**; typical procedure. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL8029752