



Selective synthesis of 3-methyleneindolin-2-ones by one-pot multicatalytic processes

Ren-Jie Song, Yu Liu, Rong-Jiang Li, Jin-Heng Li *

Key Laboratory of Chemical Biology and Traditional Chinese Medicine Research (Ministry of Education), Hunan Normal University, Changsha 410081, China

ARTICLE INFO

Article history:

Received 2 March 2009

Revised 11 April 2009

Accepted 17 April 2009

Available online 22 April 2009

Keywords:

Copper

Palladium

Amination

C–H functionalization

Propiolamide

3-Methyleneindolin-2-one

ABSTRACT


A novel one-pot multicatalytic route for the synthesis of 3-methyleneindolin-2-ones has been developed involving sequential copper-catalyzed amination and palladium-catalyzed C–H functionalization processes. In the presence of CuI and Pd(OAc)₂, a variety of propiolamides underwent the reaction with iodides to afford the corresponding 3-methyleneindolin-2-ones in moderate yields.

© 2009 Elsevier Ltd. All rights reserved.

The 3-methyleneindolin-2-one moieties are frequently present in naturally occurring and biologically active compounds that display great potential utilizations in many major therapeutic areas, such as oncology, inflammation, CNS, immunology, and endocrinology.¹ Therefore, attention of organic chemists has been increasingly drawn to the development of efficient methods for their preparation.^{2–4} Among the numerous transformations, palladium-catalyzed cyclization of *N*-arylpropiolamide involving an arene C–H functionalization process displayed efficiency particularly for the synthesis of 3-methyleneindolin-2-ones.^{2–6} However, only few papers have been reported for this purpose.^{2,3} Zhu and co-workers, for example, have first developed the palladium-catalyzed domino carbopalladation/C–H activation/C–C bond-forming reaction that uses an anilide sp² C–H bond and an electrophile (aryl iodide) as the coupling partners (Scheme 1).² Subsequently, we have demonstrated some new protocols for constructing the 3-methyleneindolin-2-one skeleton by palladium-catalyzed oxidative C–H functionalization of an anilide sp² C–H bond with an nucleophile (based on amide, acid, and ArI(OAc)₂).³ Due to continuing interest in this area,^{3,4f} we report here a novel one-pot multicatalytic route⁷ for the synthesis of 3-methyleneindolin-2-ones involving sequential copper-catalyzed amination and palladium-catalyzed C–H functionalization processes (Scheme 1).^{8,9}

The reaction between *N*-(4-methoxyphenyl)-3-phenylpropiolamide (**1a**) and 1-iodo-4-methoxybenzene (**2a**) was investigated to


optimize the reaction conditions, and the results are summarized in Table 1. Initially, a series of ligands **L1–L7** were examined, and the results showed that **L1**, *N,N*-dimethylethane-1,2-diamine, was the most efficient (entries 1–7). After treatment of amide **1a** with iodide **2a**, CuI, and **L1** in DMF/MeCN for 12 h, the reaction was conducted in the same vessel for another 12 h after the addition of Pd(OAc)₂ to afford the target product **3** in a 57% yield (entry 1). However, the other nitrogen-containing ligands **L2–L6** were inferior to **L1** (entries 2–6). The reported effective Buchwald–Hartwig amination phosphine **L7** was also examined for the one-pot reaction, and only 34% yield of **3** was obtained (entry 7). Among the effects of solvents examined, we found that a mixture of DMF/MeCN (1:1) provided the best results (entries 1 and 8–14).



Scheme 1.

* Corresponding author. Tel.: +86 731 8872 576; fax: +86 731 8872 101.

E-mail address: jhli@hunnu.edu.cn (J.-H. Li).

Table 1
Screening optimal conditions^a

Entry	Ligand	[Pd]	Solvent	Isolated yield (%)
1	L1	Pd(OAc) ₂	DMF/MeCN (1:1)	57
2	L2	Pd(OAc) ₂	DMF/MeCN (1:1)	Trace
3	L3	Pd(OAc) ₂	DMF/MeCN (1:1)	17
4	L4	Pd(OAc) ₂	DMF/MeCN (1:1)	47
5	L5	Pd(OAc) ₂	DMF/MeCN (1:1)	Trace
6	L6	Pd(OAc) ₂	DMF/MeCN (1:1)	18
7	L7	Pd(OAc) ₂	DMF/MeCN (1:1)	34
8	L1	Pd(OAc) ₂	DMF	29
9	L1	Pd(OAc) ₂	MeCN	Trace
10	L1	Pd(OAc) ₂	Toluene	28
11	L1	Pd(OAc) ₂	Toluene/MeCN (1:1)	22
12	L1	Pd(OAc) ₂	DMSO/MeCN (1:1)	24
13	L1	Pd(OAc) ₂	DMF/MeCN (1:4)	20
14	L1	Pd(OAc) ₂	DMF/MeCN (4:1)	32
15	L1	Pd(CF ₃ CO ₂) ₂	DMF/MeCN (1:1)	37
16	L1	PdBr ₂	DMF/MeCN (1:1)	27
17	L1	PdCl ₂ (PPh ₃) ₂	DMF/MeCN (1:1)	32
18	L1	Pd ₂ (dba) ₃	DMF/MeCN (1:1)	55
19	L1	K ₂ PdCl ₄	DMF/MeCN (1:1)	56
20 ^b	L1	Pd(OAc) ₂	DMF/MeCN (1:1)	21
21 ^c	L7	Pd(OAc) ₂	DMF/MeCN (1:1)	20

Reaction conditions: **1a** (0.2 mmol), **2a** (2.2 equiv), Pd(OAc)₂ (10 mol %), **L7** (20 mol %), K₂CO₃ (2.2 equiv), and DMF/MeCN (1:1, 4 mL) at 100 °C for 12 h, then Pd(OAc)₂ (10 mol %) at 100 °C for 12 h.

^a Reaction conditions: **1a** (0.2 mmol), **2a** (2.2 equiv), CuI (10 mol %), ligand (20 mol %), K₂CO₃ (2.2 equiv), and solvent (4 mL) at 100 °C for 12 h, then [Pd] (10 mol %) at 100 °C for 12 h. Some side products were observed from decomposition of the C–N bonds by GC–MS analysis.

^b Both CuI (10 mol %) and Pd(OAc)₂ (10 mol %) were added at the same time for 24 h.

^c Reaction conditions: **1a** (0.2 mmol), **2a** (2.2 equiv), Pd(OAc)₂ (10 mol %), **L7** (20 mol %), K₂CO₃ (2.2 equiv), and DMF/MeCN (1:1, 4 mL) at 100 °C for 12 h, then Pd(OAc)₂ (10 mol %) at 100 °C for 12 h.

Finally, a variety of other Pd catalysts, including Pd(CF₃CO₂)₂, PdBr₂, PdCl₂(PPh₃)₂, Pd₂(dba)₃, and K₂PdCl₄, were evaluated (entries 15–19). The screening results demonstrated that the efficiency of Pd₂(dba)₃ or K₂PdCl₄ is the same as that of Pd(OAc)₂, but Pd(CF₃CO₂)₂, PdBr₂, and PdCl₂(PPh₃)₂ were less effective. It is noteworthy that both CuI and Pd(OAc)₂ were added to the mixture of amide **1a** and iodide **2a** at the same time for 24 h providing the product **3** in a low yield (entry 20). The results demonstrated that only a low yield was obtained without Cu catalysts by a Pd-catalyzed amination/C–H functionalization process (entry 21).^{9,8e}

With the optimal conditions in hand, the scopes of both propiolamides **1** and iodides **2** were examined (Table 2).¹⁰ We were pleased to find that the reactions of amide **1a** with various iodides **2b**, **2c**, and **2e** were still conducted smoothly in moderate yields (entries 1, 2, and 4), but the reaction of the electron-deficient aryl iodide **2d** gave a mixture of products (entry 3). Treatment of substrate **1a** with iodobenzene (**2b**), CuI, **L1**, and Pd(OAc)₂, for instance, selectively afforded two products **4** and **5** in 62% total yield (entry 1). The electron-rich aryl iodide **2c** was also suitable for the one-pot reaction under the standard conditions (entry 2). Interestingly, iodomethane (**2e**) could undergo the reaction with amide **1a** to afford the corresponding 3-methyleneindolin-2-one **8** in 47% yield (entry 4). Subsequently, we chose both propiolamides **1** and iodides **2** to control the selectivity (entries 5–9). 3-Phenyl-*N*-*p*-tolylpropiolamide (**1d**), for instance, was reacted with 1-iodo-4-methylbenzene (**2c**) to give the product **11** alone in 53% yield (entry 7). It is noteworthy that 21% yield can still be achieved from the reaction between the two bulk substrates, 3-

phenyl-*N*-*o*-tolylpropiolamide (**1f**) and 1-iodo-2-methylbenzene (**2g**) (entry 9). *N*-Alkyl phenylpropiolamides **1g** and **1h** were also investigated under the standard conditions (entries 10 and 11). The results demonstrated that both amides **1g** and **1h** could undergo the one-pot reaction selectively in 36% and 33% yields, respectively.

As shown in Scheme 2, a controlled reaction was conducted. Treatment of *N*-(4-methoxyphenyl)-3-phenylpropiolamide (**1a**) with 1-iodo-4-methoxybenzene (**2a**), CuI, and **L1** afforded the corresponding amination product **16** in 50% isolated yield. Subsequently, the reaction of **16** was carried out under the Zhu's conditions to give the target product **3** in 48% yield.

A possible mechanism was proposed as outlined in Scheme 3 on the basis of the previously reported mechanisms and the present results.^{2,3,8,9} Reaction of amide **1** with CuL₂ affords intermediate **A** with the aid of base. Intermediate **A** undergoes the reaction with iodide **2** to yield the product **B** and to regenerate the active Cu species.⁸ Subsequently, C–H functionalization of the product **B** occurs after the addition of Pd catalyst to give the target 3-methyleneindolin-2-ones by Zhu's process.² We deduced that Pd-catalyzed amination of substrate **1** with iodide **2** may take place to give the product **B** in the multicatalytic processes (entry 21 in Table 1).^{9,8e} Based on the present results, the stereoselectivity may mainly depend on the characters of both aryl group on the propiolamide moiety and substituent R¹ of iodides, such as electronic effect and steric effect. Study of the true mechanism is in progress.

In summary, we have developed a novel one-pot protocol for the synthesis of 3-methyleneindolin-2-ones by sequential copper-catalyzed amination and palladium-catalyzed C–H functional-

Table 2Selective synthesis of 3-methyleneindolin-2-ones by one-pot multicatalytic processes^a



























Entry	Amide 1	Iodide 2	Product/isolated yield ^b (%)
1	 1a	 2b	 4 (27)  5 (35)
2	 1a	 2c	 6 (37; <i>E/Z</i> = 2 : 1)  7 (27)
3	 1a	 2d	Mixture
4	 1a	Mel 2e	 8 (47)
5	 1b	 2f	 9 (32; <i>E/Z</i> = 1 : 1)
6	 1c	 2b	 10 (45)
7	 1d	 2c	 11 (53)
8	 1e	 2g	 12 (55; <i>E/Z</i> = 2 : 1)
9	 1f	 2h	 13 (21; <i>E/Z</i> = 3 : 1)

Table 2 (continued)


Entry	Amide 1	Iodide 2	Product/isolated yield ^b (%)
10			 14 (36)
11			 15 (33; <i>E/Z</i> = 3 : 1)

^a Reaction conditions: **1** (0.2 mmol), **2** (2.2 equiv), CuI (10 mol %), **L1** (20 mol %), and K₂CO₃ (2.2 equiv) in DMF/MeCN (1:1, 4 mL) at 100 °C for 12 h, then Pd(OAc)₂ (10 mol %) at 100 °C for 12 h. Some side products were observed from decomposition of the C–N bonds by GC–MS analysis.

^b Ratios of *Z/E* isomers were determined by ¹H NMR.



Scheme 2. A controlled reaction.



Scheme 3. A possible mechanism.

ization reactions. In the presence of CuI and Pd(OAc)₂, a variety of propiolamides underwent the multicatalytic processes with iodides to afford the corresponding 3-methyleneindolin-2-ones in moderate yields. Efforts to extend the application of the transformation in organic synthesis are underway in our laboratory.

Acknowledgments

The authors thank the New Century Excellent Talents in University (No. NCET-06-0711), Scientific Research Fund of Hunan Provincial Education Department (No. 08A037), and National Natural Science Foundation of China (No. 20572020) for financial support.

Supplementary data

Supplementary data associated with this Letter can be found, in the online version, at [doi:10.1016/j.tetlet.2009.04.066](https://doi.org/10.1016/j.tetlet.2009.04.066).

References and notes

- (a) Mohammadi, M.; McMahon, G.; Sun, L.; Tang, C.; Hirth, P.; Yeh, B. K.; Hubbard, S. R.; Schlessinger, J. *Science* **1997**, 276, 955; (b) Sun, L.; Tran, N.; Liang, C.; Tang, F.; Rice, A.; Schreck, R.; Waltz, K.; Shawver, L. K.; McMahon, G.; Tang, C. *J. Med. Chem.* **1999**, 42, 5120; (c) Hare, B. J.; Walters, W. P.; Caron, P. R.; Bemis, G. W. *J. Med. Chem.* **2004**, 47, 4731; (d) Noble, M. E. M.; Endicott, J. A.; Johnson, L. N. *Science* **2004**, 304, 1800; (e) Liao, J. J.-L. *J. Med. Chem.* **2007**, 50, 409; (f) *Drugs Future* **1990**, 15, 898; (g) Robinson, R. P.; Reiter, L. A.; Barth, W. E.; Campeta, A. M.; Cooper, K.; Cronin, B. J.; Destito, R.; Donahue, K. M.; Falkner, F. C.; Fiese, E. F.; Johnson, D. L.; Kuperman, A. V.; Liston, T. E.; Malloy, D.; Martin, J. J.; Mitchell, D. Y.; Rusek, F. W.; Shamblin, S. L.; Wright, C. F. *J. Med. Chem.* **1996**, 39, 10; (h) Andreani, A.; Burnelli, S.; Granaola, M.; Leoni, A.; Locatelli, A.; Morigi, R.; Rambaldi, M.; Varoli, L.; Kunkel, M. W. *J. Med. Chem.* **2006**, 49, 6922; (i) Graczyk, P. P. *J. Med. Chem.* **2007**, 50, 5773.
- (a) Pinto, A.; Neuville, L.; Retaillieu, P.; Zhu, J. *Org. Lett.* **2006**, 8, 4927; (b) Pinto, A.; Neuville, L.; Zhu, J. *Angew. Chem., Int. Ed.* **2007**, 46, 3291; (c) Pinto, A.; Neuville, L.; Zhu, J. *Tetrahedron Lett.* **2009**, 8, 9–89. doi:10.1016/j.tetlet.2009.03.086.
- (a) Tang, S.; Peng, P.; Pi, S.-F.; Liang, Y.; Wang, N.-X.; Li, J.-H. *Org. Lett.* **2008**, 10, 1179; (b) Tang, S.; Peng, P.; Wang, Z.-Q.; Tang, B.-X.; Deng, C.-L.; Li, J.-H.; Zhong, P.; Wang, N.-X. *Org. Lett.* **2008**, 10, 1875; (c) Tang, S.; Peng, P.; Zhong, P.; Li, J.-H. *J. Org. Chem.* **2008**, 73, 5476.
- For other papers on the synthesis of oxindoles using Pd catalysts, see: (a) Cheung, W. S.; Patch, R. J.; Player, M. R. *J. Org. Chem.* **2005**, 70, 3741; (b) Yanada, R.; Obika, S.; Inokuma, T.; Yanada, K.; Yamashita, M.; Ohta, S.; Takemoto, Y. *J. Org. Chem.* **2005**, 70, 6972; (c) Couty, S.; Liégault, B.; Meyer, C.; Cossy, J. *Org. Lett.* **2004**, 6, 2511; (d) D'Souza, D. M.; Rominger, F.; Müller, T. J. *J. Angew. Chem., Int. Ed.* **2005**, 44, 153; (e) Yanada, R.; Obika, S.; Oyama, M.; Takemoto, Y. *Org. Lett.* **2004**, 6, 2825; (f) Tang, S.; Yu, Q.-F.; Peng, P.; Li, J.-H.; Zhong, P.; Tang, R.-Y. *Org. Lett.* **2007**, 9, 3413.
- For papers on the synthesis of oxindoles using Rh as the catalysts and 2-alkynylaryl isocyanates as the substrates, see: (a) Shintani, R.; Yamagami, T.; Hayashi, T. *Org. Lett.* **2006**, 8, 4799; (b) Miura, T.; Takahashi, Y.; Murakami, M. *Org. Lett.* **2007**, 8, 5075; (c) Miura, T.; Takahashi, Y.; Murakami, M. *Org. Lett.* **2008**, 10, 1743.
- For papers on the other methods for the synthesis of oxindoles, see: (a) Mori, M.; Ban, Y. *Tetrahedron Lett.* **1979**, 20, 1133; (b) Sun, L.; Liang, C.; Shirazian, S.; Zhou, Y.; Miller, T.; Cui, J.; Fukuda, J. Y.; Chu, J.-Y.; Nematalla, A.; Wang, X.; Chen, H.; Sista, A.; Luu, T. C.; Tang, F.; Wei, J.; Tang, C. *J. Med. Chem.* **2003**, 46, 1116; (c) Wang, L.; Zhang, Y.; Hu, H.-Y.; Fun, H. K.; Xu, J.-H. *J. Org. Chem.* **2005**, 70, 3805; (d) Kalinski, C.; Umkehrer, M.; Schmidt, J.; Ross, G.; Kolb, J.; Burdack, C.; Hiller, W.; Hoffmann, S. D. *Tetrahedron Lett.* **2006**, 47, 4683; (e) Xing, X.; Wu, J.; Luo, J.; Dai, W.-M. *Synlett* **2006**, 2099; (f) Yang, T.-M.; Liu, G. J. *Comb. Chem.* **2007**, 9, 86.

7. For selected reviews on the transition metal-catalyzed multicatalytic processes, see: (a) Dragutan, V.; Dragutan, I. *J. Organomet. Chem.* **2006**, 691, 5129; (b) Schmidt, B. *Pure Appl. Chem.* **2006**, 78, 469; (c) De Meijere, A.; Von Zezschwitz, P.; Bräse, S. *Acc. Chem. Res.* **2005**, 38, 413; (d) Wasilke, J. C.; Obrey, S. J.; Baker, R. T.; Bazan, G. C. *Chem. Rev.* **2005**, 105, 1001; (e) Ajamian, A.; Gleason, J. L. *Angew. Chem., Int. Ed.* **2004**, 43, 3754; (f) Fogg, D. E.; dos Santos, E. N. *Coord. Chem. Rev.* **2004**, 248, 2365; (g) Prestat, G.; Poli, G. *Chemtracts* **2004**, 17, 97; (h) Lee, J. M.; Na, Y.; Han, H.; Chang, S. *Chem. Soc. Rev.* **2004**, 33, 302; (i) Wang, C.; Xi, Z. *Chem. Soc. Rev.* **2007**, 36, 1395.
8. For selected papers on Cu-catalyzed amination of amides with aryl halides, see: (a) Klapars, A.; Antilla, J. C.; Huang, X.; Buchwald, S. L. *J. Am. Chem. Soc.* **2001**, 123, 7727; (b) Klapars, A.; Huang, X.; Buchwald, S. L. *J. Am. Chem. Soc.* **2002**, 124, 7421; (c) Jiang, L.; Job, G. E.; Klapars, A.; Buchwald, S. L. *Org. Lett.* **2003**, 5, 3667; (d) Klapars, A.; Parris, S.; Anderson, K. W.; Buchwald, S. L. *J. Am. Chem. Soc.* **2004**, 126, 3529; (e) Cuny, G. J.; Bois-Choussy, M.; Zhu, J. *J. Am. Chem. Soc.* **2004**, 126, 14475; (f) Yuen, J.; Fang, Y.; Lautens, M. *Org. Lett.* **2006**, 8, 653; (g) Joues, C. P.; Anderson, K. W.; Buchwald, S. L. *J. Org. Chem.* **2007**, 72, 7968; (h) Zhang, N.; Buchwald, S. L. *Org. Lett.* **2007**, 9, 4749; (i) Martin, R.; Cuenca, A.; Buchwald, S. L. *Org. Lett.* **2007**, 9, 5521; (j) Mino, T.; Harada, Y.; Shindo, H.; Sakamoto, M.; Fujita, T. *Synlett* **2008**, 4, 614; (k) Woler, M.; Klapars, A.; Buchwald, S. L. *Org. Lett.* **2009**, 11, 947; (l) Ma, D.; Cai, Q. *Acc. Chem. Res.* **2008**, 41, 1450; (m) Deng, W.; Wang, Y.-F.; Zou, Y.; Liu, L.; Guo, Q.-X. *Tetrahedron Lett.* **2004**, 45, 2311; (n) Yang, T.; Lin, C.; Fu, H.; Jiang, Y.; Zhao, Y. *Org. Lett.* **2005**, 7, 4781; (o) Pan, X.; Cai, Q.; Ma, D. *Org. Lett.* **2004**, 6, 1809–1812; (p) Shen, R.; Porco, J. A. *Org. Lett.* **2000**, 2, 1333.
9. For selected papers on Pd-catalyzed amination of amides with aryl halides, see: (a) Yang, B. H.; Buchwald, S. L. *Org. Lett.* **1999**, 1, 35; (b) Yin, J.; Buchwald, S. L. *Org. Lett.* **2000**, 2, 1101; (c) Yin, J.; Buchwald, S. L. *J. Am. Chem. Soc.* **2002**, 124, 6043; (d) Huang, X.; Anderson, K. W.; Zim, D.; Jiang, L.; Klapars, A.; Buchwald, S. L. *J. Am. Chem. Soc.* **2003**, 125, 6653; (e) Manley, P. J.; Bilodeau, M. T. *Org. Lett.* **2004**, 6, 2433; (f) Willis, M. C.; Brace, G. M.; Holmes, L. P. *Synthesis* **2005**, 3229; (g) Ligthart, G. B. W. L.; Ohkawa, H.; Sijbesma, R. P.; Meijer, E. W. *J. Org. Chem.* **2006**, 71, 375; (h) Lkawa, T.; Barder, T. E.; Biscoe, M. R.; Buchwald, S. L. *J. Am. Chem. Soc.* **2007**, 129, 13001; (i) Kotecki, B. J.; Fernando, D. P.; Haight, A. R.; Lukin, K. A. *Org. Lett.* **2009**, 11, 947.
10. **Typical procedure:** A mixture of propiolamide **1** (0.2 mmol), iodide **2** (2.2 equiv), CuI (10 mol %), *N,N'*-dimethylethane-1,2-diamine **L1** (20 mol %), and K₂CO₃ (2.2 equiv) was stirred in DMF/MeCN (v/v 1:1, 4 mL) at 100 °C for 12 h, then Pd(OAc)₂ (10 mol %) was added to the mixture, and stirred at 100 °C for another 12 h until complete consumption of starting material which was monitored by TLC. After the reaction was complete, the mixture was washed with saturated NaCl, and was extracted with diethyl ether. The organic layers were dried with Na₂SO₄ and were evaporated under vacuum, the residue was purified by flash column chromatography (hexane/ethyl acetate) to afford the pure product.
(E)-5-Methoxy-1-(4-methoxyphenyl)-3-((4-methoxyphenyl)(phenyl)methylene)-indolin-2-one (**3**): Red oil; ¹H NMR (500 MHz) δ: 7.36–7.30 (m, 9H), 6.97 (d, *J* = 4.5 Hz, 1H), 7.28 (t, *J* = 4.5 Hz, 3H), 7.27–7.18 (m, 4H), 7.17–7.15 (m, 1H), 7.02 (t, *J* = 6.5 Hz, 1H), 6.99 (t, *J* = 6.5 Hz, 1H), 6.71 (d, *J* = 8.0 Hz, 1H), 4.90 (s, 2H), 2.37 (s, 3H), 2.29 (s, 3H); ¹³C NMR (125 MHz) δ: 166.5, 160.8, 158.7, 155.7, 154.9, 140.2, 137.5, 133.5, 131.6, 130.5, 129.2, 128.1, 127.8, 127.7, 124.4, 123.6, 114.6, 114.3, 114.1, 109.3, 109.2, 55.5 (2C), 55.4; IR (KBr, cm⁻¹): 1699, 1508, 1247; LRMS (EI, 70 eV) *m/z* (%): 464 (33), 463 (M⁺, 100), 448 (13), 420 (2), 386 (3), 340 (1), 232 (19), 195 (2); HRMS (EI) for C₃₀H₂₅NO₄ (M⁺): calcd 463.1784, found 463.1783.
5-Methoxy-3-((4-methoxyphenyl)(phenyl)methylene)-1-methylindolin-2-one (**15**): ²E/Z = 3:1; Yellow oil, ¹H NMR (500 MHz) δ: 7.49 (s, 1H), 7.42–7.28 (m, 6H), 7.10 (d, *J* = 7.0 Hz, 2.3H), 7.06 (d, *J* = 7.0 Hz, 0.7H), 6.97–6.87 (m, 1H), 6.75–6.64 (m, 1H), 3.86 (s, 3H), 3.83 (s, 1H), 3.54 (s, 3H), 3.44 (s, 1H), 3.20 (s, 1H), 3.17 (s, 3H); ¹³C NMR (125 MHz) δ: 166.9, 160.7, 154.9, 154.7, 140.3, 137.2, 137.0, 133.3, 132.5, 131.6, 130.3, 129.7, 129.2, 129.1, 128.9, 127.7, 124.4, 123.9, 114.1, 114.0, 113.8, 113.0, 109.5, 109.2, 107.8; IR (KBr, cm⁻¹): 1684, 1507, 1490; LRMS (EI, 70 eV) *m/z* (%): 372 (21), 371 (M⁺, 100), 356 (18), 341 (4), 328 (3), 294 (5), 264 (3), 178 (8).