

Article

# Enediyne-comprising amino aldehydes in the Passerini reaction

Mladena Glavaš, Matija Gredi#ak, and Ivanka Jeric

ACS Comb. Sci., Just Accepted Manuscript • DOI: 10.1021/acscombsci.7b00167 • Publication Date (Web): 17 Jan 2018

Downloaded from http://pubs.acs.org on January 17, 2018

## **Just Accepted**

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Combinatorial Science is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

# Enediyne-comprising amino aldehydes in the Passerini reaction

Mladena Glavaš, Matija Gredičak, and Ivanka Jerić\*

Division of Organic Chemistry and Biochemistry, Ruđer Bošković Institute, Bijenička cesta 54, 10000 Zagreb, Croatia

KEYWORDS: multicomponent reactions, enediynes, Passerini reaction, receptors.

ABSTRACT: Multicomponent reactions represent a highly efficient approach to a broad spectrum of structurally diverse compounds starting from simple and affordable compounds. A focused library of tweezers-like compounds is prepared by employing the multicomponent Passerini reaction comprising enediyne-derived amino aldehydes. The reaction proceeds under mild conditions yielding Passerini products in good to excellent yields. Post-condensation modifications of Passerini products are demonstrated through a simple deprotection/coupling approach comprising amino functionality, furnishing enediyne cores with highly decorated arms.

# Introduction

Isolation of biologically active acetylene-containing compounds from natural sources in 1980s boosted the research efforts in different areas of organic chemistry, biochemistry, natural product synthesis, medicinal chemistry, material science, and others. Enediyne anticancer antibiotics isolated from marine and terrestrial plant sources represent a class of antitumor/antimicrobial agents with high cytotoxic activity.<sup>1</sup> The remarkable activity and a striking mode of action made enediyne compounds an attractive target molecules. Motivated by their biological importance, many cyclic and acyclic enediyne derivatives were prepared and their biological activities evaluated.<sup>1-5</sup> Apart from their role in biologically relevant processes, enediynes are versatile building blocks for the construction of complex aromatic systems through thermal or photochemical activation.<sup>6-11</sup> Furthermore, rigid, stable and spatially well-directed enediyne motif can be exploited as an inducer of turn-like conformation. In our previous studies, we showed that enedivne-based small molecule tweezers are efficient Cu(II) atom binders.<sup>12</sup> Comparing stability of studied complexes with those reported for histidine-rich peptide fragments with high affinity for copper(II), revealed that enediyne tweezers effectively mimic properties of copper(II) binding peptides. It is noteworthy that in many proteins, the proline residue is found in the vicinity of Cu(II) binding site, mostly preceding histidine residue. We therefore, concluded that structural rigidity of enediyne tweezers significantly enhances coordination properties of the receptor, thus conforming importance of the turn-like peptide conformation for the copper(II) binding. Furthermore, we demonstrated that amino acid-derived enediyne ligands account for high enantioselectivities observed in a rhodium-catalyzed asymmetric hydrogenation reaction.<sup>13</sup> In both cases, enedivne unit acts as structural element. while function is determined by groups attached at the terminal positions. We are therefore

interested to explore further the potential of enediyne compounds primarily as receptors. In that sense, conjugation of enediynes with peptide-like structures is particularly interesting, since the presence of different functional groups on the periphery of the synthetic receptors, especially chiral ones, is often the key factor for the recognition and/or discrimination of different ligands.<sup>14-18</sup> A nice example is recent study of Delsuc et al. where two peptide segments attached to a rigid template are designed to hybridize with a guest peptide molecule into a collagen-like triple helix.<sup>17</sup> Also, Faggi et al. described a family of pseudopeptidic cage receptors with a remarkable selectivity for N-protected dipeptides with an aromatic amino acid at the C-terminus.<sup>18</sup> Although cyclic and macrocyclic structures hold a prominent position as receptors for chiral molecules<sup>15</sup> and probes for protein-protein interactions,<sup>19</sup> tweezers-like compounds with cavities of flexible size proved to be effective synthetic receptors as well.<sup>14,20</sup> The main advantage of tweezers-like receptors is adaptability of the receptor geometry to the ligand topology. While size and shape of the receptor cavity is influenced by the template, selectivity of the receptor depends on the properties of its arms.

As a part of our ongoing research in enediyne chemistry, we explored methodologies for the easy construction of structurally diverse enediyne cores for the binding of different ligands. Multicomponent reactions (MCRs) offer rapid, efficient and reliable approach to libraries of structurally diverse compounds starting from simple and affordable compounds. Isocyanide-based MCRs, like Passerini and Ugi reaction<sup>21</sup> are exceptionally useful for the assembly of chimeras composed of enediynes and peptide-like structures. Among different approaches, we opted for bifunctional enediyne cores, particularly amino aldehydes **1** and **2** (Figure 1) equipped with two different functional groups that can be used in step-by-step diversification process. Aldehyde group takes part in the first MCR, while amine group is available, (after deprotection)

for the second MRC, or standard peptide coupling reaction. Our main motivation was to build a reliable and robust strategy toward library of receptors for different ligands. In this work, we explored the reactivity of aldehyde group in enediynes **1** and **2** in the Passerini reaction comprising various acids and isocyanides Thus obtained Passerini products can be further functionalized to give tweezers with highly decorated arms available for even further modifications.



Figure 1. Structures of amino aldehydes 1 and 2.

# **Results and Discussion**

Boc protected amino aldehydes **1** and **2** were prepared by a straightforward procedure comprising two consecutive Sonogashira couplings<sup>22</sup> followed by an oxidation (Scheme 1). Cross-coupling of Boc-protected propargyl amine with 1,2-diiodobenzene gave Sonogashira product **3** in 50 % yield. Next, coupling of **3** with either pent-4-yn-1-ol or prop-2-yn-1-ol gave amino alcohols **4** and **5** in excellent yields (87 and 98 %, respectively). Finally, oxidation with Dess Martin periodane proceeded smoothly providing enediyne-bridged Boc-protected amino aldehydes **1** (80 % yield) and **2** (87 % yield) (Scheme 1).





a) *tert*-butyl prop-2-ynylcarbamate, CuI (10 mol %), Pd(PPh<sub>3</sub>)<sub>4</sub> (3 mol %), in TEA, RT, under Ar, 24 h, 50 % yield; b) pent-4-yn-1-ol or prop-2-yn-1-ol (2 equiv.), CuI (3 mol %), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (3 mol %), in TEA, RT, under Ar, 24, 87 % and 98 % yield, respectively; c) DMP (2 equiv) in DCM, RT, 3 h.

In our initial experiments, Passerini reaction was performed with equimolar amounts of amino aldehyde **1**, benzoic acid and cyclohexyl isocyanide, and after 3 days in different solvents (DCM, ethyl acetate, THF) at room or elevated temperature, still showed presence of starting material. Reactions performed with excess of acid and isocyano components proceeded faster; reaction was completed in 24 h and the Passerini product **6** was obtained in 63 % yield (Chart 1). Therefore, optimized reaction conditions for the Passerini reaction include 3.1 equivalent of acid and isocyanide and reaction were carried out in DCM at room temperature until the consumption of the aldehyde component, typically 24-48h. Three additional isocyanides were used as nucleophiles; with *tert*-butyl isocyanide Passerini product **7** was gained in a 56 % yield, while a highly branched 1,1,3,3-tetramethylbutyl isocyanide gave Passerini products **8** in 81 % yield. Finally, methyl isocyanoacetate was used as a representative of amino acid-derived isocyanides, and the corresponding Passerini product was obtained in a 76 % yield. A robust methodology for the synthesis of libraries should tolerate a broad range of substrates with different electronic and steric features. Therefore, benzoic acids with different electronic properties were tested in the

Passerini reaction. As seen in Chart 1, benzoic acid with strong electron-withdrawing group at the *para*-position gave product 10 in the same yield (65 %) as benzoic acid with strong electrondonating group at the same position (product 11, 67 %). Therefore, electronic properties of the acid component do not affect its reactivity in the Passerini reaction. When halogen atoms were introduced at the *para* position of the benzoic acid aromatic ring, isolated yields were slightly lower (29-53 % for products12-15), regardless of isocyanide used (Chart 1). Next, aliphatic carboxylic acids were investigated. Acetic and isobutyric acid reacted with 1,1,3,3tetramethylbutyl isocyanide and gave rise to Passerini products 16 and 17 in 51 % and 72 % yield, respectively. Amino acids are preferable building blocks in multicomponent reactions, since they provide possibility for multiple post-condensation modifications, mainly through deprotection/cyclization protocols. Moreover, as discussed previously, amino acids and peptides are often key elements important for recognition of different ligands. Three Boc-protected amino acids were used as acid components: cyclic amino acid proline, aromatic phenylalanine and aliphatic lysine. Cyclohexyl isocyanide was used in all three reactions and Passerini products 18-20 were isolated in 63-92 % yield as a 1:1 mixture of two diastereoisomers. It is known that stereoselectivity of the Passerini reaction performed with chiral components is mainly influenced by the aldehyde component, while contribution of acid and isocyanide is generally negligible.<sup>21</sup> Therefore, the absence of stereoselectivity in the reaction performed with achiral amino aldehyde 1 is expected.

Next, we examined the reactivity of amino aldehyde **2** in the Passerini reaction with selected carboxylic acid and isocyanides. All reactions proceeded smoothly and Passerini products were isolated in 54-91 % yield (Chart 2). The influence of scale-up on the efficiency of the presented

#### **ACS Combinatorial Science**

methodology was also examined. Reaction performed with 1 mmol of aldehyde **2**, 3.1 mmol of Boc-protected phenylalanine and cyclohexyl isocyanide gave Passerini product **25** in 87 % yield. An interesting feature of Passerini products **21-26** is presence of triple bond attached to the sp<sup>3</sup> carbon atom. Recently, synthesis of structurally analogous propargylic alcohol derivatives by a copper-catalyzed oxy-alkynylation of diazo compounds with hypervalent iodine reagents was reported.<sup>23</sup> Here, presented Passerini reaction comprising easy accessible aldehyde **2** offers an alternative, metal-free approach to sp-sp<sup>3</sup> carbon bonds in very good yields and under mild reaction conditions.





<sup>a</sup>Reactions were performed with  $\sim 0.05$  mmol of 1, 3.1 equiv. of acid and 3.1 equiv. of isocyanide component. Isolated yields are given in parentheses.





<sup>a</sup> Reactions were performed with ~ 0.05 mmol of **2**, 3.1 equiv. of acid and 3.1 equiv. of isocyanide component. Isolated yields are given in parentheses.

Increasing the structural complexity of receptor is often needed to increase the binding selectivity, especially in competitive solvents, and can be achieved by adding new "arm" through a straightforward approach.<sup>18</sup> Multicomponent approach coupled with post-condensation modification(s) provides efficient strategy for the synthesis of structurally and functionally different receptors, where binding site can be crafted by careful selection of starting compounds.

Thus, multiple-arm receptors carrying hydrophobic or hydrophilic peripheral structures can be designed to probe binding of different ligands. To demonstrate the susceptibility of prepared Passerini products toward further modifications, we subjected compound **24** to Boc deprotection under acid conditions followed by coupling with *N*-terminally protected phenylalanine, and compound **27** was obtained in 50 % yield over two steps. Thus, presented Passerini reaction/post-Passerini coupling of enediyne-related amino aldehydes can be used as synthetic tool to produce tweezers-like molecules with tailored binding site. Synthesis of multiple-arm derivatives through here presented strategy, as well as a thorough complexation study of prepared Passerini compounds with a range of metal ions is currently underway, and will be published in due course.

Scheme 2. Post-Passerini modification



a) TFA/H<sub>2</sub>O (9/1) RT, 30 min; b) Boc-Phe-OH (1.2 equiv), BOP (1.5 equiv.), HOBt (1.5 equiv.), NMM (2 equiv.), in DMF, RT, overnight

## Conclusions

In summary, we successfully introduced Boc-protected enediyne-related amino aldehydes as carbonyl components in the Passerini reaction. Reactions were performed under mild reaction

conditions affording Passerini products in good to excellent yields. The reaction was tolerant of aromatic and aliphatic carboxylic acids, including amino acids, as well as of various isocyanides. Furthermore, we demonstrated that Passerini products can be additionally modified by a deprotection/coupling approach through their amino functionality.

# **Experimental procedures**

General procedure for the Passerini reaction (Charts 1 and 2) Compound 1 or 2 (~ 0.05 mmol) were dissolved in dichloromethane under argon. Carboxylic acid (3.1 equiv.) and isocyanide (3.1 equiv.) were added and the reaction mixture was stirred at room temperature until the consumption of the aldehyde component, typically 24-48h. The reaction was quenched with saturated NaHCO<sub>3</sub> and the product was extracted with dichloromethane. Organic layer was washed with brine and water, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated under reduced pressure. The product was obtained by flash column chromatography (petrol/ethyl acetate (v/v = 2/1)) to afford corresponding products 6 - 26.

#### **Representative example of the Passerini product (Chart 1)**

6-(2-(3-(*tert*-butoxycarbonylamino)prop-1-ynyl)phenyl)-1-(cyclohexylamino)-1-oxohex-5yn-2-yl benzoate (6) Compound 6 was obtained from *tert*-butyl 3-(2-(5-hydroxypent-1ynyl)phenyl)prop-2-ynylcarbamate (1) (23 mg), benzoic acid (28 mg) and cyclohexyl isocyanide (28 μL) according to the general procedure in 63% yield (25 mg, 0.046 mmol) as a yellow oil; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>):  $\delta$  = 8.08 (d, *J* = 7.6 Hz, 2H), 7.58 (t, *J* = 7.4 Hz, 1H), 7.44 (t, *J* = 7.7 Hz, 2H), 7.38 (s, 1H), 7.32 – 7.30 (m, 1H), 7.20 – 7.18 (m, 2H), 6.12 (s, 1H), 5.57 (d, *J* = 30.9 Hz, 2H), 4.21 (s, 2H), 3.84 – 3.79 (m, 1H), 2.66 (t, *J* = 6.8 Hz, 2H), 2.41 – 2.28 (m, 2H), 1.90 (t, *J* = 11.5 Hz, 2H), 1.68 – 1.65 (m, 3H), 1.45 (s, 9H), 1.35 – 1.33 (m, 2H), 1.16 – 1.10 ppm

(m, 3H); <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>):  $\delta = 167.9$ , 165.1, 133.1, 131.7, 131.3, 129.4, 128.6, 128.1, 127.4, 126.9, 125.4, 124.7, 91.6, 89.1, 81.0, 80.0, 79.1, 73.1, 59.9, 47.7, 32.4, 32.3, 30.3, 27.9, 24.9, 24.2, 24.1, 15.3 ppm; HRMS (M + Na<sup>+</sup>) calculated 565.2678, found 565.2660; IR:  $\tilde{v}_{max} = 3307, 2930, 1714, 1407, 1249, 814, 710 \text{ cm}^{-1}$ .

#### **Representative example of the Passerini product (Chart 2)**

4-(2-(3-(tert-butoxycarbonylamino)prop-1-ynyl)phenyl)-1-oxo-1-(2,4,4-trimethylpentan-

**2-ylamino)but-3-yn-2-yl benzoate (21)** Compound **21** was obtained from *tert*-butyl 3-(2-(3-oxoprop-1-ynyl)phenyl)prop-2-ynylcarbamate (**2**) (32 mg), benzoic acid (30 mg) and 1,1,3,3-tetramethylbutyl isocyanide (42 µL) according to the general procedure in 54 % yield (33 mg, 0.061 mmol) as a yellow oil; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta = 8.15 - 8.12$  (m, 2H), 7.65 - 7.60 (m, 1H), 7.51 - 7.40 (m, 4H), 7.32 - 7.22 (m, 2H), 6.36 (s, 1H), 6.15 (s, 1H), 5.78 (s, 1H), 4.20 (s, 2H), 1.77 (d, J = 4.9 Hz, 2H), 1.49 - 1.47 (m, 15H), 1.01 ppm (s, 9H); <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>):  $\delta = 164.3$ , 162.8, 133.3, 131.2, 129.5, 128.5, 128.3, 128.1, 127.4, 125.8, 123.8, 90.0, 85.6, 85.4, 80.6, 65.0, 55.4, 51.6, 30.9, 30.8, 28.5, 28.3, 27.9 ppm; HRMS (M + Na<sup>+</sup>) calculated 567.2835, found 567.2831; IR:  $\tilde{v}_{max} = 3324$ , 3127, 2953, 1681, 1409, 1366, 1246, 1303, 1026, 898, 710 cm<sup>-1</sup>.

#### ASSOCIATED CONTENT

**Supporting Information**. General methods, synthetic procedures, characterization data and copies of <sup>1</sup>H, <sup>13</sup>C and HRMS spectra are included in the SI file (PDF).

3
4
5
6
7
, 0
ð
9
10
11
12
13
14
15
16
10
17
18
19
20
21
22
23
24
24
25
26
27
28
29
30
31
32
22
22
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
<u>4</u> 0
77 50
50
51
52
53
54
55
56
57
50
20
59

60

## AUTHOR INFORMATION

**Corresponding Author** 

\*E-mail address: <u>ijeric@irb.hr</u>

## **Author Contributions**

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

#### ACKNOWLEDGMENT

We thank the Croatian Science Foundation, Grant number 3102 for financial support.

#### REFERENCES

- M. C. Joshi, D. S. Rawat. Recent development in enediyne chemistry *Chem. Biodivers.* 2012, 9, 459–498.
- (2) D. Song, Y. Tian, S. Huang, B. Li, Y. Yuan, A. Hu. An acyclic enediyne with a furyl tethering group for efficient inhibition of tumor cell viability *J. Mater. Chem. B* 2015, *3*, 8584–8588.
- D. Song. S. Sun, Y. Tian, S. Huang, Y. Ding, Y. Yuan, A. Hu. Maleimide-based acyclic enediyne for efficient DNA-cleavage and tumor cell suppression. *J. Mater. Chem. B* 2015, *3*, 3195-3200

- (4) S. E. Lindahl, H. Park, M. Pink, J. M. Zaleski. Utilizing Redox-Mediated Bergman Cyclization toward the Development of Dual-Action Metalloenediyne Therapeutics. J. Am. Chem. Soc. 2013, 135, 3826–3833.
- J. Das, S. Roy, S. Halnor, A. K. Das, A. Basak. Enediyne-based protein capture agents: demonstration of an enediyne moiety acting as a photoaffinity label. *Org. Biomol. Chem.*, 2017, *15*, 1122–1129.
- (6) C. Raviola, S. Protti, D. Ravelli, M. Fagnoni. (Hetero)aromatics from dienynes, enediynes and enyne–allenes. *Chem. Soc. Rev.* **2016**, *45*, 4364-4390
- K. Pati, G. dos Passos Gomes, I. V. Alabugin, Combining Traceless Directing Groups with Hybridization Control of Radical Reactivity: From Skipped Enynes to Defect-Free Hexagonal Frameworks. *Angew. Chem. Int. Ed.* 2016, 55. 11633-11637.
- K. Pati, G. dos Passos Gomes, T. Harris, A. Hughes, H. Phan, T. Banerjee, K. Hanson, I.
  V. Alabugin. Traceless Directing Groups in Radical Cascades: From Oligoalkynes to Fused Helicenes without Tethered Initiators. J. Am. Chem. Soc. 2015, 137, 1165–1180.
- D. Wang, F. Ling, X. Liu, Z. Li, C. Ma. Aerobic Dimerization of Enediyne Compounds: Construction of Naphthalene Frameworks. *Chem. Eur. J.* 2016, *22*, 124 – 128.
- S. Chen, Q. Li, S. Sun, Y. Ding, A. Hu. A Novel Approach toward Polyfulvene: Cationic Polymerization of Enediynes, *Macromolecules*, 2017, *50*, 534–541.
- P. M. Byers, J. I. Rashid, R. K. Mohamed, I. V. Alabugin. Polyaromatic Ribbon/Benzofuran Fusion via Consecutive Endo Cyclizations of Enediynes. *Org. Lett.* 2012, 14, 6032-6035.

- (12) M. Gredičak, N. Bregović, D. Carić, I. Jerić. Amino acid-based tweezers: The role of turn-like conformation in the binding of copper(II) *J. Inorg. Biochem.* 2012, *116*, 45–52.
- (13) Z. Kokan, Z. Glasovac, M. Majerić Elenkov, M. Gredičak, I. Jerić, S. I. Kirin. "Backdoor Induction" of Chirality: Asymmetric Hydrogenation with Rhodium(I) Complexes of Triphenylphosphane-Substituted β-Turn Mimetics. *Organometallics*, **2014**, *33*, 4005– 4015.
- (14) F. G. Klärner, B. Kahlert. Molecular Tweezers and Clips as Synthetic Receptors. Molecular Recognition and Dynamics in Receptor–Substrate Complexes *Acc. Chem. Res.* 2003, *36*, 919–932.
- (15) T. Ema. Synthetic macrocyclic receptors in chiral analysis and separation J. Incl.
  Phenom. Macrocycl. Chem. 2012, 74, 41–55.
- (16) S. Matile, A. Vargas Jentzsch, J. Montenegro, A. Fin. Recent synthetic transport systems *Chem. Soc. Rev.* 2011, 40, 2453–2474.
- N. Delsuc, S. Uchinomiya, A. Ojidab, I. Hamachi, A Host-Guest System Based on Collagen-Like Triple-Helix Hybridization. *Chem. Commun.*, 2017, 53, 6856-6859.
- (18) E. Faggi, A. Moure, M. Bolte, C. Vicent, S. V. Luis, I. Alfonso. Pseudopeptidic Cages as Receptors for N-Protected Dipeptides. J. Org. Chem. 2014, 79, 4590–4601.
- (19) T. A. F. Cardote, A. Ciulli. Cyclic and Macrocyclic Peptides as Chemical Tools To Recognise Protein Surfaces and Probe Protein–Protein Interactions *ChemMedChem*, 2016, 11, 787–794.

 H. Y. Kuchelmeister, C. Schmuck. Nucleotide Recognition in Water by a Guanidinium-Based Artificial Tweezer Receptor. *Chem. Eur. J.* 2011, *17*, 5311–5318.

- (21) C. de Graaff, E. Ruijter and R. V. A. Orru. Recent developments in asymmetric multicomponent reactions. *Chem. Soc. Rev.* 2012, *41*, 3969–4009.
- (22) R. Chinchilla, C. Najera. The Sonogashira Reaction: A Booming Methodology in Synthetic Organic Chemistry *Chem. Rev.* 2007, 107, 874–922.
- (23) D. P. Hari, J. Waser. Copper-Catalyzed Oxy-Alkynylation of Diazo Compounds with Hypervalent Iodine Reagents. J. Am. Chem. Soc. 2016, 138, 2190–2193.



R<sub>2</sub>-NC

21 examples yields up to 92 %

ő

NHBoc

0

0

HN-R2