


This article was downloaded by: [Nanyang Technological University]

On: 28 April 2015, At: 16:54

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Incn20>

One-Pot Synthesis of Antiviral Acyclovir and Other Nucleosides Derivatives Using Doped Natural Phosphate as Lewis Acid Catalyst

Hassan B. Lazrek ^a, Laila Baddi ^a, Michael Smietana ^b, Jean-Jacques Vasseur ^b, Saïd Sebti ^c & Mohamed Zahouily ^d

^a Unité de Chimie Biomoléculaire et Médicinale, Faculty of Science Semlalia, Marrakech, Morocco

^b Institut des Biomolécules Max Mousseron, UMR 5247 CNRS-UMI-UM II, Université de Montpellier II, Montpellier, France

^c Laboratoire de Chimie Organique Catalyse et Environnement, Faculté des Sciences Ben M'Sik, Casablanca, Morocco

^d Laboratoire de Catalyse, Chimiométrie et Environnement, Faculté des Sciences et Techniques, Mohammedia, Morocco

Published online: 11 Sep 2008.

To cite this article: Hassan B. Lazrek, Laila Baddi, Michael Smietana, Jean-Jacques Vasseur, Saïd Sebti & Mohamed Zahouily (2008) One-Pot Synthesis of Antiviral Acyclovir and Other Nucleosides Derivatives Using Doped Natural Phosphate as Lewis Acid Catalyst, *Nucleosides, Nucleotides and Nucleic Acids*, 27:10-11, 1107-1112, DOI: [10.1080/15257770802341285](https://doi.org/10.1080/15257770802341285)

To link to this article: <http://dx.doi.org/10.1080/15257770802341285>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

ONE-POT SYNTHESIS OF ANTIVIRAL ACYCLOVIR AND OTHER NUCLEOSIDES DERIVATIVES USING DOPED NATURAL PHOSPHATE AS LEWIS ACID CATALYST

Hassan B. Lazrek,¹ Laila Baddi,¹ Michael Smietana,² Jean-Jacques Vasseur,² Saïd Sebti,³ and Mohamed Zahouily⁴

¹*Unité de Chimie Biomoléculaire et Médicinale, Faculty of Science Semlalia, Marrakech, Morocco*

²*Institut des Biomolécules Max Mousseron, UMR 5247 CNRS-UMI-UM II, Université de Montpellier II, Montpellier, France*

³*Laboratoire de Chimie Organique Catalyse et Environnement, Faculté des Sciences Ben M'Sik, Casablanca, Morocco*

⁴*Laboratoire de Catalyse, Chimiométrie et Environnement, Faculté des Sciences et Techniques, Mohammedia, Morocco*

- *Natural phosphate doped with iodine or potassium iodide is an active catalyst for the one-pot synthesis of acyclonucleosides. To demonstrate the utility of the new catalyst system, the highly important antiviral drug acyclovir was directly and regioselectively obtained from NAcG with no byproducts.*

Keywords Acyclovir natural phosphate; Lewis Acid catalyst

INTRODUCTION

Reactions facilitated on various solid inorganic surfaces are receiving increasing attention.^[1] The advantages of these methods over conventional homogeneous reactions is that they provide greater selectivity, enhanced rates, clean products, and ease of handling. In an effort to develop new

Received 11 February 2008; accepted 6 June 2008.

Part of this work was first presented on the IS3NA XVI International Roundtable, Minneapolis, Minnesota, USA, September 12–16, 2004.

This project was supported by the Comité Mixte Interuniversitaire Franco-Marocain Programme (A.I. no. MA/06/143). The authors are indebted to Dr. C. Kwong from the Southern Research Institute Birmingham, Alabama, USA, for his help discussion and correcting the manuscript.

Address correspondence to H. B. Lazrek, Unité de Chimie Biomoléculaire et Médicinale, Faculty of Science Semlalia, Marrakech, Morocco. E-mail: hblazrek50@gmail.com; hblazrek@ucam.ac.ma




FIGURE 1 Acyclic nucleoside analogues of acyclovir.

practical and economic catalysts, the use of natural phosphate (NP) and doped NP in various chemical transformations has recently been investigated.^[2] These types of catalysts represent an important environmentally friendly alternative to reactions otherwise toxic and expensive, and many efforts are being pursued to promote NP.^[3] Acyclovir **1**, ganciclovir **2**, and penciclovir **3** (Figure 1) are widely used drugs for the treatment of opportunistic herpes virus infections in immunocompromised patients. Although commercialized for more than 25 years, an inexpensive production of acyclovir is still challenging, especially in the field of generic drugs.


As a continuation of our ongoing program toward the development of new and economical syntheses of antiviral drugs, we report here a clean and efficient preparation of acyclovir and its analogues using NP as catalyst.

RESULTS AND DISCUSSION


Several methods for the synthesis of these antiviral agents have been reported. A review covering these approaches was recently published.^[4] The major problems associated with the synthesis of acyclovir and its analogues were 1) the reagents employed were quite expensive; 2) the amination of the 2-chloro group was tedious; 3) the yield of acyclovir was usually not satisfactory; and 4) the accompanying formation of the *N*⁷-isomer, which can be very difficult to separate from the expected *N*⁹-isomer.^[5] According to the patent literature^[6] for the synthesis of acyclovir, the drug has been prepared in 24% yield by debenzoylation of the product obtained by the reaction of trimethylsilylated guanine and 2-benzyloxyethoxymethyl chloride.

Considering the high value of acyclovir and its analogues, we decided to undertake a study to identify the best catalyst for a simple, inexpensive, and general synthesis of acyclonucleosides. Herein, we describe the virtues of NP as an efficient heterogeneous catalyst for use in this emerging area. Pyrimidines or purines were first condensed with hexamethyldisilazane (HMDS) and directly treated with 1.1 equivalent of (2-acethoxyethoxy)methylacetate without distilling to remove excess HMDS. Indeed, upon refluxing uracil

with ammonium sulfate in HMDS for 1 hour, the addition of the alkylating agent led to a small amount of desired acyclonucleoside (Table 1, entry 1). Therefore, in connection with our previous work, we studied the influence of various additives as doping partners and increase the catalytic activity of NP. With this aim, NP was impregnated with various amount of either I₂ [2e] or KI [2d] (Scheme 1). These additives were assumed to aid in both the N¹ regioselective isomer (pyrimidine) formation as well as activating the alkylation reagent, thereby significantly increasing rates of acyclonucleosides formation. Control experiments indicated that an impregnation of NP with these additives is essential for this one-pot reaction to be efficient. Whereas addition of 0.2 equivalent of TMSI showed no catalytic effect whatsoever (entry 2) we found that 0.2 equivalent of I₂/NP allowed a 20% yield increase, as compared to 0.2 equivalent of I₂ alone (compare entries 3 and 6). With these conditions in hand, other acyclonucleosides derivatives were obtained in yields ranging from 34% to 50% (entries 7–10). In all cases, the reaction was found to be regioselective, allowing only the formation of the N¹ isomer for pyrimidines and the N⁹ isomer for purines, with no traces of other isomers (N⁷).




BH= U=uracil, T= thymine, C= cytosine, A= adenine, Nac-G= N-acetyl guanine



SCHEME 1 Conditions for synthesis of acyclonucleoside analogues to acyclovir.

The mechanism of the above glycosylation can be depicted as follows (Scheme 2): Silylated uracil reacts with KI/NP to give (CH₃)₃Si-I. The latter reacts with the alkylating agent R-O-CH₂-OAc to afford R-O-CH₂-I. Further, the complex [Heterocycle]-/[NP-K]⁺ then reacts with R-O-CH₂-I to conduct to the target acyclonucleoside.^[7] To further examine the possibility of NP catalysis with non volatile additives, NP was impregnated with potassium iodide. As expected, KI alone gave only traces of product (entry 11) whereas activated NP with 0.8 equivalent of KI gave modest-to-good amounts of the acyclonucleosides, depending on the starting bases, with no loss of regioselectivity (entries 15–18). Finally, acyclovir was quantitatively obtained from its diacetate precursor after treatment with a saturated solution of



SCHEME 2 Mechanism of the glycosylation.

ammonia in methanol followed by evaporation of the solvent, with 30% overall yield from N_{Ac}G.^[8,9]

The used and recovered catalysts were then shown to be reusable after drying at 150°C in vacuum, and more efficiently after washing with acetone followed by calcinations at 900°C.^[2g]

In conclusion, a general one-pot synthesis of acyclonucleosides starting from unprotected nucleobases is presented. Although the yields are modest, the use of an inexpensive NP coated with I₂ or KI as catalyst make this reaction particularly attractive. Easy to set up and totally regioselective, the reaction is notably facile and does not require fastidious work-up. Because of the high potential of acyclovir and its analogues, we believe that this reaction will be of broad interest to scientists involved in the synthesis of antiviral drugs.

EXPERIMENTAL SECTION

Typical Procedure for Catalyst Preparation

- Natural Phosphate Coated with Iodine (NP/I₂):* To a solution of iodine (759 mg) in CH₂Cl₂ (5 ml) was added natural phosphate (3 g) and stirred for 15 minutes and evaporated to dryness.

TABLE 1 Alkylation of natural bases with (2-acetoxyethoxy)methylacetate (1.1 equivalent) under various catalytic conditions

Entry	Base	Catalyst (Eq.)	Yield (%) ^a
1	U	NP	5
2	U	TMSI (0.2)	3
3	U	I ₂ (0.2)	30
4	U	NP/I ₂ (0.8)	35
5	U	NP/I ₂ (0.5)	40
6	U	NP/I ₂ (0.2)	50
7	T	NP/I ₂ (0.2)	50
8	C	NP/I ₂ (0.2)	35
9	A	NP/I ₂ (0.2)	48
10	N _{Ac} G	NP/I ₂ (0.2)	34
11	U	KI (0.8)	8
12	U	NP/KI (0.5)	30
13	U	NP/KI (0.8)	50
14	U	NP/KI (1)	44
15	T	NP/KI (0.8)	45
16	C	NP/KI (0.8)	30
17	A	NP/KI (0.8)	50
18	N _{Ac} G	NP/KI (0.8)	24

^aIsolated yield.

- b) *Natural Phosphate Coated with Potassium Iodide (NP/KI)*: To a solution of potassium iodide (1 g) in water (5 ml) was added natural phosphate (3 g). The mixture was stirred for 15 minutes and evaporated to dryness.

Typical Procedure for One-Pot Synthesis

To uracil (112 mg, 1 mmol) was added hexamethyldisilazane (4 mL) and ammonium sulphate (10 mg). The mixture was refluxed for 2 hours. To the obtained clear solution was added 2-acetoxyethyl acetoxymethyl ether (200 mg, 1.1 mmol), the catalyst (473 mg, 0.2 mmol of I₂, or 473 mg, 0.8 mmol of KI) and acetonitrile (5 mL). After being refluxed overnight, the mixture was filtered and the solvent evaporated. The crude product was purified by column chromatography. The desired product was obtained with 50% yield.

REFERENCES

- Clark, J.H. Solid acids for green chemistry. *Acc. Chem. Res.* **2002**, 35, 791–797.
- a) Zahouily, M.; Mezdar, A.; Elmakssoudi, A.; Mounir, B.; Rayadh, A.; Sebti, S.; Lazrek, H.B. Comparison of different Lewis acids supported on natural phosphate as new catalysts for chemoselective dithioacetalization of carbonyl compounds under solvent-free conditions. *ARKIVOC* **2006**, 31–40; b) Zahouily, M.; Bahlaouan, B.; Rayadha, A.; Sebti, S. Natural phosphate and potassium fluoride doped natural phosphate: efficient catalysts for the construction of a carbon-nitrogen bond. *Tetrahedron Lett.* **2004**, 45, 4135–4138; c) Alahiane, A.; Rochdi, A.; Taourirte, M.; Redwane, N.; Sebti, S.; Engels, J.W.; Lazrek, H.B. Building blocks for polyamide nucleic acids: Facile synthesis using potassium fluoride doped natural phosphate as basic catalyst. *Nucleosides Nucleotides Nucleic Acids*

- 2003, 22, 109–114; d) Lazrek, H.B.; Rochdi, A.; Redwane, N.; Ouzebila, D.; Vasseur J.J. A one pot synthesis of D-ribonucleosides using Natural phosphate doped with KI in HMDS. *Lett. Org. Chem.* **2006**, 3, 313–314; e) Lazrek, H.B.; Ouzebila, D.; Baddi, L.; Vasseur, J.J. *Nucleosides Nucleotides Nucleic Acids* **2007** 26, 1095; f) Sebti, S.; Smahi, A.; Solhy, A. Natural phosphate doped with potassium fluoride and modified with sodium nitrate: efficient catalysts for the Knoevenagel condensation. *Tetrahedron Lett.* **2002**, 43, 1813–1815.; g) Zahouily, M.; Mezdar, A.; Rakik, J.; Elmakssoudi, A.; Rayadh, A.; Sebti, S.J. A mild and efficient method for the protection of carbonyl compounds as dithioacetals, dithiolanes and dithianes catalysed by iodine supported on natural phosphate. *Mol. Catal. A* **2005**, 233, 43–47.
- NP comes from an ore extracted in the region of Khouribga (available in raw form or treated form from CERPHOS Casablanca, Morocco). Prior to use this material requires initial treatments such as crushing and washing. For use in organic synthesis, the NP is treated by techniques involving attrition, sifting, calcinations (900°C), washing, and recalcination. These treatments lead to a fraction between 100 and 400 μm , which is rich in phosphate. The structure of NP is similar to that of fluorapatite $[\text{Ca}_{10}(\text{PO}_4)_6\text{F}_2]$, as shown by x-ray diffraction and chemical analysis. The surface area of NP was measured at $\mu\text{m}^2 \text{g}^{-1}$ (nitrogen adsorption) and the total pore volume was $0.005 \text{ cm}^3 \text{ g}^{-1}$.
 - Gao, H.; Mitra, A.K. Synthesis of acyclovir, ganciclovir, and their prodrugs. *Synthesis* **2000**, 329–351 and references cited therein.
 - Matsumoto, H.; Kaneko, C.; Yamada, K.; Takeuchi, T.; Mori, T.; Mizuno, Y. A convenient synthesis of 9-(2-Hydroxyethoxymethyl)guanine (Acyclovir) and related compounds. *Chem. Pharm. Bull.* **1988**, 36, 1153–1157.
 - Schaeffer, H.J. Ger. Pat. No. 2539963, **1974**.
 - Ubasawa, M.; Takashima, H.; Sekiya, K. A convenient one-pot synthesis of acyclonucleosides. *Chem. Pharm. Bull.* **1995**, 43, 142–143.
 - Lazrek, H. B.; Taouririte, M.; Barascut, J.L.; Imbach, J.L. Solid-liquid phase transfer catalysis II: A convenient approach to the synthesis of ACV, HBV and related compounds. *Bull. Soc. Chim. Belg.* **1996**, 105, 391–395 and references cited therein.
 - Robins, H. J.; Hatfield, P.W. Nucleic acid related compounds.37. Convenient and high-yield synthesis of N-[(2-hydroxyethoxy)methyl] heterocycles as acyclic nucleoside analogues *Can. J. Chem.* **1982**, 60, 547–553.