Use of InCl₃ as a Cocatalyst and a Cl₂Pd(DPEphos)–P(2-Furyl)₃ Catalyst System for One-Pot Hydrometalation-Cross-Coupling and Carbometalation-Cross-Coupling **Tandem Processes**

Mingxing Qian, Zhihong Huang, and Ei-ichi Negishi*

Herbert C. Brown Laboratories of Chemistry, Purdue University, 560 Oval Drive, West Lafayette, Indiana 47907-2084

negishi@purdue.edu

Received February 17, 2004

ORGANIC LETTERS

2004Vol. 6, No. 10 1531 - 1534

ABSTRACT

R¹C ≡CZ
$$\xrightarrow{\mathbb{R}^{2}\mathsf{ML}_{n}} \begin{bmatrix} z \\ \mathbb{R}^{1} \\ \mathbb{R}^{2} \\ \mathbb{R}$$

One-pot conversion of alkynes to regio- and stereodefined alkenylmetals containing AI and Zr via hydrometalation or carbometalation followed by their Pd-catalyzed cross-coupling with (E)-ICH=CHBr or (E)-ICH=CHCI proceeds cleanly and selectively to give the corresponding 1-halo-1,3-dienes in excellent yields using a catalyst system consisting of Cl₂Pd(DPEphos), DIBAL-H, and TFP (catalyst A) with InCl₃ as a cocatalyst.

The one-pot synthesis of conjugated dienes via hydrometalation involving Al1 and Zr2 hydrides followed by Pdcatalyzed cross-coupling, as well as via Zr-catalyzed carboalumination and Pd-catalyzed cross-coupling,³ was originally reported during between 1976 and 1978. In many cases, however, it was not only desirable but necessary to use either cocatalysts or stoichiometric promoters, and none have been superior to ZnCl₂ or ZnBr₂ as a cocatalyst for these processes.^{3,4} Even so, these one-pot processes have still been inferior, in some highly demanding cases,^{4f,5} to the process involving (i) hydrometalation or carbometalation, (ii) iodinolysis followed by isolation and purification of alkenyl iodides, (iii) lithiation with 2 equiv of t-BuLi in ether, (iv)

transmetalation with dry ZnCl₂ or ZnBr₂ to generate preformed alkenylzinc derivatives, and (v) Pd-catalyzed crosscoupling.

In our recent attempts to improve the synthetic usefulness of the above-mentioned tandem processes for the syntheses of dienes and polyenes further, Pd-catalyzed cross-coupling reactions of (E)-ICH=CHBr^{4d,f,g,6,7} and (E)-ICH=CHCl⁷ have

⁽¹⁾ Baba, S.; Negishi, E. J. Am. Chem. Soc. 1976, 98, 6729. See also: Negishi, E.; Baba, S. J. Chem. Soc., Chem. Commun. 1976, 596.

⁽²⁾ Okukado, N.; Van Horn, D. E.; Klima, W. L.; Negishi, E. Tetrahedron Lett. 1978, 1027. See also: Negishi, E.; Van Horn, D. E. J. Am. Chem. Soc. 1977, 99, 3168.

⁽³⁾ Negishi, E.; Okukado, N.; King, A. O.; Van Horn, D. E.; Spiegel, B. I. J. Am. Chem. Soc. 1978, 100, 2254.

⁽⁴⁾ For representative examples of the use of Pd-Zn double-metal catalysis, see: (a) Panek, J. S.; Hu, T. J. Org. Chem. **1997**, 62, 4912; 4914. (b) Hu, T.; Panek, J. S. J. Org. Chem. 1999, 64, 3000; J. Am. Chem. Soc. 2002, 124, 11368. (c) Drouet, K. E.; Theodorakis, E. A. J. Am. Chem. Soc. 1999, 121, 456; Chem. Eur. J. 2000, 1987. (d) Negishi, E.; Alimardanov, A.; Xu, C. Org. Lett. 2000, 2, 65. (e) Thompson, C. F.; Jamison, T. F.; Jacobson, E. N. J. Am. Chem. Soc. 2000, 122, 10482; J. Am. Chem. Soc. 2001, 123, 9974. (f) Zeng, F.; Negishi, E. Org. Lett. 2001, 3, 719. (g) Zeng, F.; Negishi, E. Org. Lett. 2002, 4, 703. (5) Negishi, E.; Owczarczyk, Z. Tetrahedron Lett. 1991, 32, 6682.

⁽⁶⁾ Available from Aldrich Chemical Co.

^{(7) (}a) Negishi, E.; Okukado, N.; Lovich, S. F.; Luo, F. T. J. Org. Chem. 1984, 49, 2629. (b) Negishi, E.; Hata, M.; Xu, C. Org. Lett. 2000, 2, 3687. (c) Negishi, E.; Qian, M.; Zeng, F.; Anastasia, L.; Babinski, D. Org. Lett. 2003, 5, 1597.

Scheme 1												
,Br												
ⁿ Hex ⁱ Bu ₂ AlH ⁿ Hex / Al ⁱ Bu ₂	Cat. A,	additive	▶ ″⊦	lex I Br + n	Hex 2 + "	Hex (1)						
	additive mol%	^o C	time h	I (%)	II (%)	III (%)						
	None	23	10	5	9	75						
	ZnBr ₂ (10	0) 23	4	34	35	24						
	InCl ₃ (10)	0	4	82	2	7						
Br												
ⁿ Hex <u>HZrCp₂Cl</u> ⁿ Hex ZrCp ₂ d	CI Cat. A,	additive	•	"Hex Br.	+ ⁿ Hex 2	+ "Hex (2)						
	additive mol%	^o C	time h	I (%)	II (%)	III (%)						
	None	23	10	15	14	46						
	ZnBr ₂ (10	0) 23	12	31	28	30						
	InCl ₃ (34)	0	4	77	1	12						
Me₃Al Me	ı~	_Br		Ме	Me	Me						
ⁿ Hex — cat. Cl ₂ ZrCp ₂ n _{Hex} AlMe	e ₂ Cat. A,	additive	•	ⁿ Hex IV Br +	ⁿ Hex 2 ⁺	ⁿ Hex (3)						
	additive mol%	^o C	time h	IV (%)	V (%)	VI (%)						
	None	0	10	4	trace	81						
	ZnBr ₂ (10	0) 0	4	26	42	24						
	InCl ₃ (34)	0	4	91	2	6						
1. ${}^{i}Bu_{2}AIH$ 2. I_{2} 3. ${}^{t}BuLi$ (2 equiv) 4. MX ₀ \longrightarrow MX = 4	I Cat. A, a	3r dditive	_	ⁿ Hex Br +	ⁿ Hex 2+	ⁿ Hex (4)						
Hex	THF, 0 °	C, 4 h		I		III (1)						
	MXn			I (%)	II (%)	III (%)						
Cat. A = 1% Cl ₂ Pd(DPEphos)	ZnBr ₂ (1	0 equiv)		75	4	10						
2% DIBAL-H, 2% TFP	InCl ₃ (0.3	34 equiv)	67	16	12						

been investigated. While their Pd-catalyzed cross-coupling with alkynylzincs^{4d,7} has been shown to be widely applicable and generally satisfactory, their Pd-catalyzed reactions with alkenyl,- aryl,- and alkylmetals containing Al, Zr, and others have proved to be much more demanding, and satisfactory results have, in the past, been obtained only in those cases where β , β -dialkyl-substituted alkenylalanes generated by Zrcatalyzed carboalumination were employed.4f Extensive screening and optimization of Pd catalysts and cocatalysts, as well as other parameters such as solvents, have therefore been conducted, and a system consisting of Cl₂Pd(DPEphos)⁸ [typically 1 mol %, DPEphos = bis(2-diphenylphosphinophenyl)ether], tris(2-furyl)phosphine⁹ (TFP) (typically 2 mol %), and InCl₃, but not ZnCl₂ or ZnBr₂, as a cocatalyst has proved to be uniquely satisfactory and distinctly superior to any of the previously developed protocols and all of the other procedures that have been examined in this study. Both the use of InCl₃¹⁰ as a cocatalyst¹¹ and the demonstration of synthetically beneficial effects of double-phosphine catalyst systems appear to be unprecedented. Although InBr₃ was as effective as InCl₃, the latter was chosen, as it is less expensive, especially on a molar basis.

Representative experimental results are summarized in Scheme 1. For comparison, the results observed in the

⁽⁸⁾ Kranenburg, M.; Kamer, P. C. J.; Van Leeuwen, P. W. N. M. Eur. J. Inorg. Chem. **1998**, 155.

⁽⁹⁾ Farina, V.; Krishnan, B. J. Am. Chem. Soc. 1991, 113, 9585.

⁽¹⁰⁾ For the stoichiometric use of InCl₃ in the Pd-catalyzed crosscoupling, see: (a) Pérez, I.; Sestelo, J. P.; Sarandeses, L. A. Org. lett. 1999, 121, 1267. J. Am. Chem. Soc. 2001, 123, 4155. (b) Pena, M. A.; Pérez, I.; Sestelo, J. P.; Sarandeses, L. A. Chem. Commun. 2002, 2246. (c) Rodriguez, D.; Sestelo, J. P.; Saradeses, L. A. J. Org. Chem. 2003, 68, 2518. (d) Gelman, D.; Schumann, H.; Blum, J. *Tetrahedron Lett.* **2000**, *41*, 7555. (e) Shenglof, M.; Gelman, D.; Heymer, B.; Schumann, H.; Molander, G. A.; Blum, J. Synthesis 2003, 302. (f) Jaber, N.; Schumann, H.; Blum, J. J. Heterocycl. Chem. 2003, 40, 565. (g) Takami, K.; Yorimitsu, H.; Shinokubo, H.; Matsubara, S.; Oshima, K. *Org. Lett.* **2001**, *3*, 1997. (h) Takami, K.; Yorimitsu, H.; Oshima, K. *Org. Lett.* **2002**, *4*, 2993. (i) Nakamura, T.; Kinoshita, H.; Shinokubo, H.; Oshima, K. Org. Lett. 2002, 4, 3165. (j) Takami, K.; Mikami, S.; Yorimitsu, H.; Shinokubo, H.; Oshima, K. J. Org. Chem. 2003, 68, 6627. (k) Lee, P. H.; Sung, S.; Lee, K. Org. Lett. 2001, 3, 3201. (1) Lee, K.; Lee, J.; Lee, P. H. J. Org. Chem. 2002, 67, 8265. (m) Lee, K.; Seomoon, D.; Lee, P. H.; Angew. Chem., Int. Ed. 2002, 41, 3901. (n) Lee, P. H.; Lee, S. W.; Lee, K. Org. Lett. 2003, 5, 1103. (o) Lee, P. H.; Lee, S. W.; Seomoon, D. Org. Lett. 2003, 5, 4963. (p) Legros, J. Y.; Promault, G.; Fiaud, J. C. Tetrahedron 2001, 57, 2507. (q) Lehmann, U.; Awasthi, S.; Minehan, T. Org. Lett. 2003, 5, 2405.

 Table 1.
 Pd- and In-Cocatalyzed Hydrometalation-Cross-Coupling and Carbometalation-Cross-Coupling Tandem Processes for

 One-Pot Conversion of Alkynes into Stereo- and Regiodefined 1-Halo-1,3-dienes



Cat. A = 1% Cl₂Pd(DPEphos), 2% DIBAL-H, 2% TFP

						yield, ^a %		
entry	R ¹ C≡CZ	R^2ML_n	ICH=CHX	mol % InCl ₃	time, h	product	dimer	monomer
1	ⁿ HexC≡CH	HAl ⁱ Bu ₂	ICH=CHBr	34	4	85 (81 ^b)	<1	5
2	ⁿ HexC≡CH	HAl ^{<i>i</i>} Bu ₂	ICH=CHBr	10	8	82	2	7
3	ⁿ HexC≡CH	HAl ⁱ Bu ₂	ICH=CHBr	7	10	76	4	15
4	ⁿ HexC≡CH	HAl ^{<i>i</i>} Bu ₂	ICH=CHCl	34	4	87 (85 ^b)	<1	4
5	ⁿ HexC≡CH	HAl ^{<i>i</i>} Bu ₂	ICH=CHCl	10	8	80	<1	11
6	ⁿ HexC≡CH	HZrCp ₂ Cl	ICH=CHBr	34	4	77	1	12
7	ⁿ HexC≡CH	HZrCp ₂ Cl	ICH=CHBr	10	8	76	3	13
8	ⁿ HexC≡CH	HZrCp ₂ Cl	ICH=CHBr	7	10	70	6	19
9	PhC≡CH	HZrCp ₂ Cl	ICH=CHBr	34	4	89 (84 ^b)	<1	5
10	EtC≡CEt	HZrCp ₂ Cl	ICH=CHBr	34	4	86 (82 ^b)	<1	с
11	ⁿ HexC≡CH	Me ₃ Al-Cl ₂ ZrCp ₂	ICH=CHBr	34	4	91 (87 ^b)	2	6
12	ⁿ HexC≡CH	Me ₃ Al-Cl ₂ ZrCp ₂	ICH=CHBr	10	8	85	3	10
13	ⁿ HexC≡CH	Me ₃ Al-Cl ₂ ZrCp ₂	ICH=CHBr	7	10	77	8	12
14	ⁿ HexC≡CH	Me ₃ Al-Cl ₂ ZrCp ₂	ICH=CHCl	34	4	95 (93 ^b)	<1	<1
15	$HOCH_2CH_2C \equiv CH$	Me ₃ Al-Cl ₂ ZrCp ₂	ICH=CHBr	34	10	80 (70 ^b)	<1	с
^a Yield	by GLC. ^b Isolated yield.	Not readily detected by	GLC.					

corresponding reactions of preformed (E)-1-octenylzinc bromide and tris[(E)-1-octenyl]indium are also shown in eq 4 in Scheme 1. These results indicate the following: (1) In none of the tandem reactions shown in eqs 1-3 were the previously developed Pd-Zn double-metal-catalyzed procedures satisfactory, even in cases where the optimized catalyst system (catalyst A) consisting of 1 mol % Cl₂Pd(DPEphos), 2 mol % TFP, and 2 mol % DIBAL-H for reduction of Cl₂Pd(DPEphos) (vide infra) was used. (2) In contrast, the Pd-In double-metal-catalyzed process employing catalyst A has been highly and uniquely satisfactory. (3) And yet, In is not superior to Zn in the corresponding stoichiometric reactions of alkenylmetals containing Zn and In. Furthermore, the Pd-In cocatalysis is superior to either of the reactions represented by eq 4 in Scheme 1, pointing to favorable synergistic effects associated with Al-In and Zr-In combinations. Irrespective of the precise mechanistic interpretation, the Pd-In double-metal-catalyzed protocol presented in eqs 1-3 of Scheme 1 has provided a one-pot tandem procedure that is far superior to any previously developed ones for the indicated transformations.

As indicated earlier, one of our recent papers^{4f} described a satisfactory one-pot conversion of terminal alkynes into (E,E)-1-bromo-4-methyl-1,3-dienes via (i) carboalumination with Me₃Al-Cl₂ZrCp₂ and (ii) cross-coupling in the presence of 5 mol % Pd(PPh₃)₄, 1 equiv of ZnBr₂, and DMF used as a cosolvent for the synthesis of unsymmetrical carotenoids. The use of THF as a solvent led to very low product yields. It was also necessary to evaporate CH₂Cl₂ used for Zr-catalyzed methylalumination before the addition of DMF and/ or THF. More critically, even this procedure was found to be rather unsatisfactory in cases where β -monosubstituted alkenylmetals containing Al or Zr were used. The procedure reported herein has solved all of the problems mentioned above and appears to be widely applicable and generally dependable, as indicated by the results summarized in Table 1. All of the isolated products were of \geq 98% isomeric purity by ¹H and ¹³C NMR spectroscopy and GLC analysis.

It may be worth describing in some detail how uniquely advantageous catalyst A consisting of Cl₂Pd(DPEPhos) and TFP is, even though no persuasive rationalization can be presented at this time. In view of multistage nature of Pdcatalyzed cross-coupling, it was envisioned that optimal Pd catalysts for two or more microsteps may be different from one another. On this basis, various two-phosphine catalyst systems were screened initially for the reaction of preformed (E)-1-octenylzinc bromide with (E)-ICH=CHBr (1.2 equiv) at 23 °C in THF, and the combination of Cl₂Pd(DPEphos) and TFP, i.e., catalyst A, has been found to be distinctly superior to any others. Specifically, the use of one-phosphine Pd catalyst systems (5 mol % in Pd) such as Pd(PPh₃)₄ (18%), Pd(*t*-Bu₃P)₂ (12%), Pd₂(dba)₃-2TFP (24\%), Cl₂Pd(dppf)-2DIBAL-H (40%), and Cl₂Pd(DPEphos)-2DIBAL-H (43%) all led to the disappointingly low yields indicated in parentheses. Various binary combinations of the

⁽¹¹⁾ For the use of InBr₃ as a cocatalyst in the Pd-catalyzed crosscoupling of terminal alkynes with aryl iodides, see: Sakai, N.; Annaka, K.; Konakahara, T. *Org. Lett.* **2004**, *6*, 1527. We thank Prof. N. Sakai for sharing their results with us before publication.

above-mentioned phosphines were then screened, and catalyst A was found to be by far the most satisfactory combination, as shown in eq 4 in Scheme 1. On the other hand, all of the others, including Cl₂Pd(DPEphos)-dppf (31%), Cl₂Pd-(DPEphos)-2PPh₃ (39%), Cl₂Pd(dppf)-2TFP (51%), $Cl_2Pd(dppf)-2PPh_3$ (44%), $Cl_2Pd(dppf)-DPEphos$ (38%), $Cl_2Pd(PPh_3)_2$ -dppf (42%), $Pd(t-Bu_3P)_2$ -dppf (38%), Pd(*t*-Bu₃P)₂-2TFP (32%), Pd(*t*-Bu₃P)₂-DPEphos (17%), and $Pd(t-Bu_3P)_2-2PPh_3$ (21%), led to low product yields accompanied by extensive formation of byproducts. In cases where Pd(II) complexes were used, DIBAL-H was used as an external reducing agent. Mysteriously, even Cl₂Pd(TFP)₂-DPEphos (36%) and Pd₂(dba)₃-2DPEphos-4TFP (37%) were distinctly and unmistakably inferior to catalyst A. At present, the unique superiority of catalyst A over all of the others tested remains as a mechanistic puzzle to be clarified.

The superior catalytic activity of catalyst A extends beyond the 1-halo-1,3-diene synthesis described above. Thus, it has substantially improved the reaction of preformed arylzinc derivatives with (E)-ICH=CHBr, as indicated in Scheme 2.

It has become increasingly clear from this and other recent works reported by us and others that selective Pd- or Nicatalyzed cross-coupling reactions involving $1,1^{-12}$ and $1,2^{-12}$ dihaloalkenes^{7c} tend to be considerably more capricious and demanding but that satisfactory procedures may nevertheless be developed through optimization of (i) catalysts, especially ligands, (ii) cocatalysts, and (iii) solvents. In this paper, the use of InCl₃ as a cocatalyst and/or a double-phosphine



^{*a*} Yield by GLC. ^{*b*}Isolated yield. ^{*c*}(p-MeOC₆H₄)₃In used in place of p-MeOC₆H₄ZnBr.

catalyst system consisting of $Cl_2Pd(DPEphos)$ and TFP has been shown to be effective in selective monosubstitution of 1,2-dihaloethylenes.

Acknowledgment. We thank the National Science Foundation (CHE-0309613), the National Institutes of Health (GM 36792), and Purdue University for support of this work. We also thank Boulder Scientific Co. and FMC Corp. for their assistance in the procurement of Zr and P reagents, respectively.

Supporting Information Available: Detailed experimental procedures and compound characterization data. This material is available free of charge via the Internet at http://pubs.acs.org.

OL049716F

^{(12) (}a) Shi, J.; Zeng, X.; Negishi, E. Org. Lett. 2003, 5, 1825. (b) Zeng, X.; Hu, Q.; Qian, M.; Negishi, E. J. Am. Chem. Soc. 2003, 125, 13636. (c) Shi, J.; Negishi, E. J. Organomet. Chem. 2003, 687, 518. (d) Zeng, X.; Qian, M.; Hu, Q.; Negishi, E. Angew. Chem., Int. Ed. 2004, in press.