1685

Mono- and Bi-dentate Phosphaalkene Ligands: Structures of their Copper(I) Chloride Complexes[†]

Abdelaziz Jouaiti,[#] Michel Geoffroy^{*,#} and Gérald Bernardinelli^b

^a Department of Physical Chemistry, University of Geneva, 30 Quai Ernest Ansermet, 1211 Geneva 4, Switzerland

^b Laboratory of Crystallography, University of Geneva, 24 Quai Ernest Ansermet, 1211 Geneva 4, Switzerland

While the reaction of the monophosphaalkene $[(2,4,6-tri-tert-butylphenyl)phosphanediylmethyl]-benzene (L²) with copper(1) chloride led to a dimeric complex <math>[Cu_2Cl_2L_2^2(NCMe)_2]$ 1, use of the diphosphaalkene 1,3-bis $[(2,4,6-tri-tert-butylphenyl)phosphanediylmethyl]benzene (L¹) gave rise to a tetrameric complex <math>[Cu_4Cl_4L_2^1]$ ·3EtOH 2. The crystal structures of the complexes have been determined and indicate that 1 is a centrosymmetric dimer with tetrahedrally co-ordinated copper(1) centres, whereas 2 adopts a 'cubane-like' structure with pairs of copper atoms bridged by different L¹ units.

Since the first synthesis¹ of a stable phosphaalkene in 1978, compounds containing a trivalent dico-ordinated phosphorus atom have been intensively investigated. These molecules exhibit particularly interesting chemical reactivities² and are known to form σ and η^2 complexes 3 with a large variety of transition metals. In this context, we have previously shown that 1,3-bis[(2,4,6-tri-tert-butylphenyl)phosphanediylmethyl]benzene,⁴ L^1 , is a specially interesting compound which can act as a terdentate chelate when reacting with palladium(II) and platinum(II) ions. However, the type of complexes that it can form with metals that are not likely to give rise to orthometallation (e.g. Cu¹) remains unknown. Surprisingly, whereas the literature on the complexation of Cu⁺ ions by phosphines is particularly abundant,⁵ to our knowledge no phosphaalkene complex of Cu^I has ever been reported. We will first show that such a σ bond can indeed be formed between Cu^I and the monodentate [(2,4,6-tri-tert-butylphenyl)phosphane-diylmethyl]benzene,⁶ L^2 , and will then describe the 1:2 L^{1} : copper(I) chloride complex, which, to our knowledge, is the first Cu₄X₄L₂ 'cubane-like' structure with bridging ligands.

Results and Discussion

The binuclear complex 1 was obtained as a yellow solid by treating stoichiometric quantities of L^2 with CuCl in CH₂Cl₂–MeCN. It is air-stable for many weeks when stored at 5 °C. This complex is a centrosymmetric dimer with copper(1) centres tetrahedrally co-ordinated to a phosphorus atom, two chlorine atoms and a solvent molecule (Fig. 1). The Cu₂Cl₂ core has an irregular diamond shape with two slightly different Cu–Cl bonds (Table 1). A similar chloride-bridged dimeric structure has recently been reported for a phosphine complex⁷ [(Ph₃P)(py)CuCl₂Cu(py)(PPh₃)] (py = pyridine); a notable difference between these two structures lies in the Cu₂Cl₂ unit in which the bond angles are closer to 90° for the phosphaalkene complex (Table 1) than for the phosphine complex ⁷ (Cl–Cu–Cl 105.05, Cu–Cl–Cu 74.4 and 75.5°).

The reaction of compound L^1 with CuCl or CuCl₂ in CH₂Cl₂ at 40 °C leads to the yellow crystalline complex 2. The crystal structure indicates that, as for the 1:1 triphenylphosphine: copper(1) chloride complex,⁸ a cubane-like structure is formed (Fig. 2). In 2, however, two faces of the Cu₄Cl₄ core contain two



copper atoms which are σ bonded to the two phosphaalkene groups of the same L¹ ligand. These 'bridged faces' (Cu₂Cl₂) are planar (maximum deviations = 0.047 and 0.048 Å) and parallel (0.6°). Moreover, the two phosphaalkene bonds of each bridging ligand are coplanar and the corresponding planes of the two L¹ entities are perpendicular to each other (90.1°) and almost perpendicular to the 'bridged face' (89.4 and 86.2° respectively). Although the Cu₄Cl₄ core of the molecule is largely distorted from an idealized cube, all geometrical

[†] Supplementary data available: see Instructions for Authors, J. Chem. Soc., Dalton Trans., 1994, Issue 1, pp. xxiii-xxviii.

Table 1 Selected bond lengths (Å) and angles (°) for complexes 1 and 2

		1	2 <i>ª</i>
Cu • • • Cu	b	3.1659(9)	3.058(3)
	с		3.218(11)
$Cl \cdots Cl$	b	3.553(2)	3.836(6)
	с		3.530(21)
Cu-Cl	b	2.355(2), 2.404(2)	2.453(16)
	с		2.354(18)
Cu-N		2.041(6)	` `
Cu-P		2.205(1)	2.167(4)
P-Calkene		1.663(5)	1.66(1)
P-C _{aryl}		1.840(5)	1.85(1)
ClCuCl	b	96.59(5)	102.9(8)
	с		94.5(4)
Cu-Cl-Cu	b	83.41(6)	77.1(3)
	с		84.0(6)
ClCu-P	b	121.56(7), 112.52(6)	114(2)
	с		131.2(6)
Cl-Cu-N		105.2(2), 104.4(2)	``
PCuN		114.1(1)	_
Cu-P-Calkene		130.8(2)	126.8(5)
Cu-P-C		127 4(2)	131(1)



Fig. 1 Molecular structure of complex 1

parameters (Table 1) show excellent correlations if we distinguish the two 'bridged faces' from the four others. The presence of the bridging ligand acts as an ordering factor of the Cu_4Cl_4 core, contrary to the soft structure observed in the analogous triphenylphosphine complex.⁸ Owing to the presence of the bridging ligand, the Cu···Cu distances of the 'bridged faces' [3.060(3) and 3.056(4) Å] are smaller than in the other faces [mean 3.218(11) Å]. Consequently, the Cu-Cl-Cu angles decrease and the Cl-Cu-Cl angles increase significantly.

It should be noted that, even if no stacking interaction were observed, the benzene rings of the two ligands are not similarly oriented. Thus, one L^1 unit is almost planar (angles between the benzene ring and the plane of each P=C-C moiety are 6°), whereas for the second L^1 the corresponding angles are equal to 22°.

 Table 2
 Summary of crystal data, intensity measurements and structure refinements for complexes 1 and 2*

	1	2
Formula	$C_{54}H_{76}Cl_2Cu_2N_2P_2$	$C_{88}H_{128}Cl_4Cu_4P_4$.
М	1013.2	1844.1
a/Å	9.238(1)	15.420(2)
b/Å	11.463(2)	19.101(3)
c/Å	14.116(4)	19.932(5)
a/°	110.38(1)	108.93(1)
β/°	101.94(1)	96.74(1)
ν/°	94.73(1)	111.76(1)
Ú/Å ³	1351.3(5)	4965(2)
Z	1	2
F(000)	536	1956
$D_{-}/g \mathrm{cm}^{-3}$	1.25	1.23
$\mu(Mo-K\alpha)/mm^{-1}$	0.981	1.062
$\Gamma(\sin\theta)/\lambda 1_{max}/A^{-1}$	0.55	0.49
No. measured reflections	3763	9222
No. observed reflections	3277	5635
$[F_{o} > 4\sigma(F_{o})]$		
No. parameters	281	954
Weighting scheme, w	$1/\sigma^2(F_o)$	1
Maximum and minimum $\Delta \rho/e \ \text{Å}^{-3}$	0.78, -0.68	1.21, -0.91
S	3.61	4.97
R, R'	0.060, 0.049	0.076, 0.076
		• · · · ·

* Details in common: crystal size $0.2 \times 0.2 \times 0.2$ mm; triclinic, space group $P\overline{1}$; full-matrix least-squares refinement on F.

Such 'cubane-like' $Cu_4X_4L_2$ edifices are quite unusual. As recently reported for the 1,2-bis(diphenylphosphino)methane ligand,^{9,10} in the presence of two bidentate ligands, the Cu_4X_4 core adopts a 'step' structure; the present cubane-like geometry is likely to be due to the phosphaalkene moieties which increase the rigidity of the ligands.

As shown by ¹H and ³¹P NMR spectroscopy, in solution the two L¹ ligands in complex 2 are equivalent; it is therefore probable that the difference observed in the crystal is due to a matrix effect. A noticeable difference between the spectra obtained for L¹ and 2 concerns the chemical shift of the aromatic proton located between the two phosphaalkene groups. This chemical shift passes from δ 7.52 (L¹) to 9.68 (2) (for L², δ 7.6; for 1, δ 7.76); the chemical shifts of the protons *ortho* to the C=P bonds remain similar to that measured for L¹. The variation probably reflects the fact that, in 2, in contrast to the other three molecules, there is no free rotation around the C(phosphaalkene)–C(phenyl) bond; the two magnetic susceptibility tensors associated with the phosphaalkene double bonds are therefore parallel and fixed with respect to the *o*-proton.

Experimental

All experiments were performed under a nitrogen atomosphere in carefully dried glassware. Reagents were purchased from Fluka and used as received. Solvents were dried, where necessary, by standard methods before use. NMR spectra were recorded at room temperature using a Bruker AC-F200 spectrometer (¹H, 200 MHz; ³¹P, 81 MHz).

Syntheses.— $[Cu_2Cl_2L^2_2(NCMe)_2]$ 1. A solution containing CuCl (0.027 g, 0.2 mmol) in MeCN (2 cm³) was added to a solution of compound L² (0.1 g, 0.2 mmol) in CH₂Cl₂ (5 cm³) and refluxed for 30 min. After filtration of unreacted CuCl, yellow crystalline 1 was obtained by slow evaporation of the solution, m.p. 174 °C (decomp.). NMR: ³¹P-{¹H} (CDCl₃, external reference H₃PO₄), δ 205.9; ¹H(CD₂Cl₂, internal reference SiMe₄), δ 1.4 (s, 18 H), 1.6 (s, 36 H), 1.97 (s, 6 H), 7.28

 Table 3
 Non-hydrogen positional parameters

168	I
-----	---

able 3	Non-nyarogen positio	onal parameters					
Atom	X/a	Y/b	Z/c	Atom	X/a	Y/b	Z/c
Complex	1						
Cu	0.364 93(8)	0.389 00(6)	0.429 86(5)	C(14)	0.255 1(6)	-0.015 0(5)	0.359 9(5)
Cl	0.423 3(2)	0.593 7(1)	0.431 5(1)	C(15)	0.296(1)	0.092 2(7)	0.461 9(6)
Р	0.342 4(2)	0.219 6(1)	0.288 4(1)	C(16)	0.227(1)	-0.129 5(7)	0.383 6(8)
C(1)	0.240 6(6)	0.180 8(5)	0.166 7(4)	C(17)	$0.103\ 3(8)$	-0.0002(9)	0.299 5(6)
C(2)	0.1401(6)	0.254 0(5)	0.126 1(4) 0.033 2(4)	C(18)	0.0008(7)	-0.2759(5)	0.178 8(4)
C(3)	-0.0300(7)	$0.194 \ 3(3)$ 0.260 \ 3(7)	-0.003.5(4)	C(20)	0.000(1)	-0.287 3(0) -0.382 0(6)	0.0970(0) 0.1360(7)
C(4)	-0.059.3(8)	0.3897(7)	0.049 3(6)	C(21)	0.6975(9)	-0.2930(7)	0.270 9(6)
C(6)	0.049 6(8)	0.449 6(6)	0.138 0(5)	C(22)	0.655 6(7)	0.181 1(5)	0.217 3(5)
C(7)	0.151 2(7)	0.382 7(5)	0.177 7(4)	C(23)	0.816 4(8)	0.168 1(7)	0.230 1(7)
C(8)	0.438 4(5)	0.082 6(4)	0.276 9(4)	C(24)	0.652 2(9)	0.313 4(6)	0.285 4(7)
C(9)	0.376 7(5)	-0.019 7(5)	0.300 4(4)	C(25)	0.605 5(9)	0.171 9(8)	0.105 2(7)
C(10)	0.433 8(6)	-0.1325(5)	0.267 8(4)	N(01)	0.1860(5)	0.393 9(4)	0.494 3(4)
C(11)	0.5471(6) 0.6118(6)	-0.1471(3) -0.0437(5)	0.2100(4) 0.2025(4)	C(01)	-0.092.2(7)	0.389 4(3)	0.334 0(3)
C(12)	0.563 3(6)	-0.0437(3) 0.0729(5)	0.2336(4)	C(02)	-0.025 0(7)	0.370 3(1)	0.571 0(0)
Complex	2	0.0120 (0)	0.200 0(1)				
Cu(1)	0.731 6(1)	0.330 6(1)	0.181 2(1)	C(02)	0.378 3(9)	0.236 5(7)	0.289 9(7)
Cu(2)	0.745 7(1)	0.476 9(1)	0.312 2(1)	C(03)	0.381 3(9)	0.213 4(8)	0.349 3(7)
Cu(3)	0.534 6(1)	0.346 5(1)	0.201 1(1)	C(04)	0.310(1)	0.211(1)	0.385 8(9)
Cu(4)	0.631 8(1)	0.294 2(1)	0.305 9(1)	C(05)	0.240(1)	0.231(1)	0.365(1)
Cl(1)	0.673 0(3)	0.434 0(2)	0.180 5(2)	C(06)	0.233(1)	0.250(1)	0.303 9(9)
Cl(2)	0.795 7(3)	0.368 4(2)	0.311 6(2)	C(07)	0.296 2(9)	0.277 3(8)	0.203 8(8)
C(3)	0.5782(3) 0.5980(3)	0.2290(2) 0.4145(2)	0.1719(2) 0.3360(2)	C(08)	0.318(1) 0.279(1)	0.332 8(9)	0.0919(7)
P(1)	0.832.8(3)	0.3238(2)	0.1150(2)	C(010)	0.218(1)	0.2820(9)	-0.0297(8)
P(2)	$0.861 \ 1(3)$	0.6007(2)	0.3657(2)	C(011)	0.197(1)	0.348 1(9)	-0.0143(9)
C(1)	0.981 0(9)	0.486 1(8)	0.175 1(7)	C(012)	0.243(1)	0.408 7(9)	0.055 9(9)
C(2)	0.945 1(9)	0.518 2(8)	0.232 9(7)	C(013)	0.304(1)	0.405 1(9)	0.109 8(8)
C(3)	0.991(1)	0.601 6(8)	0.277 3(7)	C(014)	0.458(1)	0.192 9(8)	0.377 2(7)
C(4)	1.072(1)	0.651 6(9)	0.263 1(9)	C(015)	0.620(1)	0.176 2(9)	0.418 5(7)
C(5)	1.108(1)	0.618(1) 0.526(1)	0.206 2(9)	C(016)	0.626(1)	0.100(3(9))	0.382 8(8)
C(0)	1.002(1) 0.935 4(9)	0.330(1)	$0.102 \ 5(8)$ $0.123 \ 4(7)$	C(017)	0.643(1)	0.0027(3)	0.428 2(9)
C(8)	0.826(1)	0.235 0(8)	0.0365(8)	C(019)	0.667(1)	0.057(1) 0.173(1)	0.534 8(8)
C(9)	0.776(1)	0.217 9(9)	-0.036 0(9)	C(020)	0.649(1)	0.216 4(9)	0.495 9(9)
C(10)	0.784(1)	0.156 8(9)	-0.094 1(8)	C(021)	0.294(2)	0.194(1)	-0.009(1)
C(11)	0.833(1)	0.113 2(9)	-0.083 1(8)	C(022)	0.398(2)	0.209(2)	0.013(2)
C(12)	0.875(1)	0.128 5(9)	-0.0122(8)	C(023)	0.258(2)	0.148(1)	-0.091(1)
C(13)	0.8/3(1)	0.1880(9)	0.049 4(7)	C(024)	0.239(3)	0.131(1) 0.258(1)	0.019(1)
C(14)	0.939(1) 0.881(1)	0.640 5(8)	0.338 7(8)	C(023)	0.135(1)	0.336(1) 0.293(2)	-0.139(2)
C(16)	0.848(1)	0.748 3(9)	0.4401(7)	C(027)	0.176(2)	0.422(3)	-0.091(3)
C(17)	0.891(1)	0.825(1)	0.495 8(9)	C(028)	0.056(3)	0.362(4)	-0.050(2)
C(18)	0.962(1)	0.849 3(8)	0.559 1(8)	C(029)	0.344(1)	0.478(1)	0.186(1)
C(19)	0.982(1)	0.786 7(9)	0.568 8(7)	C(030)	0.446(1)	0.501(1)	0.224(1)
C(20)	0.940(1)	0.707 2(9)	0.516 2(8)	C(031)	0.282(2)	0.463(1)	0.237(1)
C(21)	0.717(1) 0.780(2)	0.262(1) 0.249(1)	-0.0572(8)	C(032)	0.346(2)	0.55/(1)	0.17/(1)
C(22)	0.780(2) 0.645(2)	0.348(1) 0.264(2)	-0.013(1)	C(034)	0.009(1)	-0.022(1)	0.300 3(8)
C(24)	0.658(1)	0.201(2)	-0.138(1)	C(035)	0.508(1)	0.022(1)	0.253 3(8)
C(25)	0.835(2)	0.046(1)	-0.149(1)	C(036)	0.690(1)	0.118(1)	0.277 5(9)
C(26)	0.818(3)	0.054(2)	-0.218(1)	C(037)	0.690(1)	0.055(1)	0.552(1)
C(27)	0.917(2)	0.028(2)	-0.136(1)	C(038)	0.796(2)	0.092(1)	0.586(1)
C(28)	0.753(3)	-0.030(2)	-0.162(2)	C(039)	0.663(2)	-0.035(1)	0.508(1)
C(29)	0.922(1)	0.19/(1) 0.124(1)	0.125 5(8)	C(040)	0.636(2)	0.056(2)	0.609(2)
C(30)	0.944(2) 0.856(2)	0.124(1) 0.198(1)	0.119(1) 0.176(1)	C(041)	0.002(1) 0.744(2)	0.305 7(9) 0.342(1)	0.544 4(8)
C(32)	1.020(1)	0.269(1)	0.160(1)	C(042)	0.690(1)	0.364 4(9)	0.506.8(9)
C(33)	0.769(1)	0.733(1)	0.376(1)	C(044)	0.569(1)	0.300(1)	0.569(1)
C(34)	0.813(2)	0.749(2)	0.317(1)	C(101)	0.399(2)	-0.024(1)	0.035(1)
C(35)	0.720(1)	0.789(1)	0.396 8(9)	C(102)	0.400(2)	-0.025(2)	-0.032(2)
C(36)	0.696(2)	0.651(2)	0.341(2)	C(103)	0.435(2)	0.012(1)	-0.067(1)
C(37)	1.016(1)	0.937 3(8)	U.019 9(9)	C(104)	0.522(3)	0.036(2)	-0.046(2)
C(30)	0.957(1) 1.042(2)	0.984(1)	0.020(1) 0.696.6(9)	C(201)	0.435(2)	0.410(2) 0.441(2)	0.522(1)
C(40)	1.107(1)	0.940(1)	0.602(1)	C(202)	0.420(2)	0.441(2) 0.470(2)	0.400(2)
C(41)	0.963(1)	0.643(1)	0.537 0(9)	C(204)	0.484(4)	0.535(3)	0.463(3)
C(42)	0.983(2)	0.667(1)	0.619(1)	C(301)	0.538(2)	0.468(2)	0.036(2)
C(43)	1.054(2)	0.640(2)	0.517(1)	C(302)	0.545(2)	0.425(2)	0.013(1)
C(44)	0.887(2)	0.558(1)	0.501(1)	C(303)	0.557(5)	0.518(5)	0.072(3)
P(01)	0.382 4(3)	0.316 2(2)	0.165 0(2)	C(304)	0.460(5)	0.433(3)	-0.024(4)
P(02) C(01)	0.308 3(3)	0.219 /(2)	0.364 8(2)	O(01)	0.578(2)	0.647(1)	0.120(1)
C(01)	0.505(1)	0.234 3(0)	0.203 0(7)				



Fig. 2 Molecular structure of complex 2

(m, 6 H), 7.52 (d, 4 H, 2 Hz), 7.76 (m, 4 H) and 7.83 (d, 2 H, 27 Hz).

[Cu₄Cl₄L¹₂]·3EtOH **2**. A solution containing CuCl (0.03 g, 0.3 mmol) in MeCN (2 cm³) was added to a solution of compound L¹ (0.1 g, 0.15 mmol) in CH₂Cl₂ (5 cm³) and refluxed for 30 min. The solvent of the resulting pale yellow solution was removed under reduced pressure. Stable yellow crystals of complex **2** were obtained by slow evaporation of a CHCl₃–EtOH (7:3) solution, m.p. 242–244 °C (decomp.). NMR: ³¹P-{¹H} (CDCl₃), δ 199.35; ¹H(CDCl₃), δ 1.37 (s, 36 H), 1.62 (s, 72 H), 6.84 (m, 4 H), 7.1 (t, 2 H, 8 Hz), 7.5 (d, 8 H, 1.8 Hz), 7.56 (d, 4 H, 26 Hz) and 9.68 (t, 2 H, 5 Hz).

Crystallography.-The cell parameters and diffracted intensities were measured at room temperature on Nonius CAD4 (1) and Philips PW1100 (2) diffractometers with graphitemonochromated Mo-K_{α} radiation ($\lambda = 0.71069$ Å). Two reference reflections showed variations of less than $3.2\sigma(I)$ for complex 1 and about 16% for 2; all intensities were corrected for this drift. Data were corrected for Lorentz and polarization effects but not for absorption. The structures were solved by direct methods using MULTAN 87,11 all other calculations used the XTAL¹² system and ORTEP.¹³ Atomic scattering factors and anomalous dispersion terms were taken from ref. 14. All hydrogen-atom coordinates were calculated. The three molecules of EtOH observed in the crystal structure of 2 are disordered. A summary of the crystal data, intensity measurements and structure refinements is given in Table 2, final atomic coordinates in Table 3.

Additional material available from the Cambridge Crystallographic Data Centre comprises H-atom coordinates, thermal parameters and remaining bond lengths and angles.

Acknowledgements

We thank the Swiss National Research Fund for support of this work and Dr. Françoise Barbalat-Rey for obtaining the NMR spectra.

References

- 1 Th. C. Klebach, R. Lourens and F. Bickelhaupt, J. Am. Chem. Soc., 1978, 100, 4886.
- 2 R. Appel, F. Knoll and I. Ruppert, Angew. Chem., Int. Ed. Engl., 1981, 20, 731.
- 3 S. I. Al-Resayes, S. I. Klein, H. W. Kroto, M. F. Meidine and J. Nixon, J. Chem. Soc., Chem. Commun., 1983, 930; H. Estiagh-Hosseini, H. W. Kroto, J. F. Nixon, M. J. Maah and M. J. Taylor, J. Chem. Soc., Chem. Commun., 1981, 199; T. A. van der Knaap, F. Bickelhaupt, J. G. Kraaykamp, G. van Koten, J. P. C. Bernards, H. T. Edzes, W. S. Veeman, E. de Boer and E. J. Baerends, Organometallics, 1984, 3, 1804.
- 4 A. Jouaiti, M. Geoffroy, G. Terron and G. Bernardinelli, J. Chem. Soc., Chem. Commun., 1992, 155.
- 5 B. J. Hathaway, Comprehensive Co-ordination Chemistry, Pergamon, Oxford, 1987, vol. 5.
- 6 M. Yoshifuji, K. Toyota and N. Inamoto, *Tetrahedron Lett.*, 1993, 26, 1727.
- 7 L. M. Engelhardt, P. C. Healy, J. D. Kildea and A. H. White, *Aust. J. Chem.*, 1989, **42**, 913.
- 8 M. R. Churchill, B. G. de Boer and D. J. Donovan, *Inorg. Chem.*, 1975, 14, 617.
- 9 S. Ramaprabu, N. Amstutz, E. A. C. Lucken and G. Bernardinelli, J. Chem. Soc., Dalton Trans., 1993, 871.
- 10 N. Marsich, G. Nardin and L. Randaccio, J. Am. Chem. Soc., 1977, 95, 4053.
- 11 P. Main, S. J. Fiske, S. E. Hull, L. Lessinger, G. Germain, J.-P. Declercq and M. M. Woolfson, MULTAN 87, A System of Computer Programs for the Automatic Solution of Crystal Structures from X-Ray Diffraction Data, Universities of York and Louvain-la-Neuve, 1987.
- 12 S. R. Hall and J. M. Stewart (Editors), XTAL 3.0 User's Manual, Universities of Western Australia and Maryland, 1992.
- 13 C. K. Johnson, ORTEP II, Report ORNL-5138, Oak Ridge National Laboratory, Oak Ridge, TN, 1976.
- 14 International Tables for X-Ray Crystallography, Kynoch Press, Birmingham, 1974, vol. 4.

Received 30th December 1993; Paper 3/07633D