


Preparation of the $[(\text{DippNP})_2(\text{P}_4)_2]^{2+}$ -Dication by the Reaction of $[\text{DippNPCl}_2]_2$ and a Lewis Acid with P_4

Michael H. Holthausen and Jan J. Weigand*

Institut für Anorganische und Analytische Chemie, Westfälische Wilhelms-Universität Münster,
Corrensstrasse 30, 48149 Münster, Germany

Received August 21, 2009; E-mail: jweigand@uni-muenster.de

The activation and functionalization of white phosphorus by *N*-heterocyclic carbenes¹ and carbene-like main group element fragments² is of considerable current interest. Carbene-analogous phosphonium cations display a pronounced ambiphilic nature which renders their behavior both Lewis acidic and Lewis basic. Their electrophilic character comes to the fore in element–element bond insertion reactions,³ making them interesting species for the systematic investigation of P_4 functionalization. Only recently, the first structurally characterized inorganic cationic P_5 cluster $[\text{P}_5\text{Br}_2]^+$ was obtained *via* phosphonium insertion by Krossing and co-workers.⁴ Our solvent-free approach to consecutively insert the phosphonium cation $[\text{Ph}_2\text{P}]^+$ into $\text{P}-\text{P}^{\text{a,b}}$ bonds of P_4 resulted in the formation of unprecedented phosphorus-rich cationic clusters $[\text{Ph}_2\text{P}_5]^+$, $[\text{Ph}_4\text{P}_6]^{2+}$, and $[\text{Ph}_6\text{P}_7]^{3+,5c}$. In this communication, we report on the functionalization of P_4 formally through the cationic bifunctional Lewis acid $[\text{DippNP}]^{2+}$ obtained from *cyclo*-1,3-diphospha-2,4-diazane $[\text{DippNPCl}_2]_2$ (**1**). This has enabled the targeted preparation of novel mono- and dicationic phosphorus-rich clusters **3** $[\text{GaCl}_4] \cdot \text{C}_6\text{H}_5\text{F}$ and **4** $[\text{Ga}_2\text{Cl}_7]_2$ (Scheme 1).

Scheme 1. Reaction of **1** with GaCl_3 and P_4 ^a

^a (i) 1 eq. GaCl_3 , $\text{C}_6\text{H}_5\text{F}$, rt, 10 min, (ii) 1 eq. P_4 , $\text{C}_6\text{H}_5\text{F}$, rt, 2 h; (iii) 3 eq. GaCl_3 , P_4 , $\text{C}_6\text{H}_5\text{F}$, rt, 6 h.

Cyclic phosphonium cation **2** can be generated from **1** in the presence of the Lewis acid GaCl_3 .⁶ The addition of one eq. GaCl_3 to a solution of **1** in $\text{C}_6\text{H}_5\text{F}$ afforded an instant color change of the initially colorless solution to deep red. $^{31}\text{P}\{^1\text{H}\}$ NMR investigation of the reaction mixture showed a new broad signal (C_6D_6 capillary, rt, $\delta = 242.3$ ppm, $\Delta\nu_{1/2} = 104$ Hz) shifted downfield compared to the sharp signal of **1** (*cis* isomer, $\delta = 210.5$ ppm, $\Delta\nu_{1/2} = 10$ Hz),⁷ indicating the formation of cation **2**.^{8a} The subsequent addition of one eq. P_4 gave rise to a clear, pale orange solution within 2 h (Scheme 1). The ^{31}P NMR spectrum of the reaction mixture revealed the exclusive formation of monocation **3**, which showed an A_2MVXZ spin system ($\delta_{\text{A}} = -346.7$ ppm, $\delta_{\text{M}} = 85.1$ ppm, $\delta_{\text{V}} = 152.7$ ppm, $\delta_{\text{X}} = 168.2$ ppm, $\delta_{\text{Z}} = 197.5$ ppm; $^1J_{\text{AV}} = -139.7$ Hz, $^1J_{\text{AX}} = -139.9$ Hz, $^1J_{\text{MV}} = -282.5$ Hz, $^1J_{\text{MX}} = -293.4$ Hz, $^2J_{\text{AM}} = 16.4$ Hz, $^2J_{\text{VX}} = 57.0$ Hz, Figure 1).^{8b,9} The resonances for this spin system are in an approximate ratio of 2:1:1:1:1. Similar to other N_2P_2 systems,^{7,10} the two ring phosphorus atoms of **3** do not couple, resulting in the observation of a singlet for the tri-coordinate phosphorus atom (s, $\delta_{\text{Z}} = 197.5$ ppm). The very air- and moisture-sensitive material was isolated as the $[\text{GaCl}_4]^-$ salt as fluorobenzene solvate.




Figure 1. ^{31}P NMR spectrum of cation **3** (in $\text{C}_6\text{H}_5\text{F}$, C_6D_6 -capillary, 25 °C; 161.94 MHz). Full spectrum (bottom) and expansions (inset) showing the experimental (up) and fitted (down) spectra;⁹ a very small amount of an unidentified side-product is indicated by an asterisk.^{8b}

A single-crystal X-ray study of **3** $[\text{GaCl}_4] \cdot \text{C}_6\text{H}_5\text{F}$ (Figure 2) confirmed the insertion of **2** into one of the $\text{P}-\text{P}$ bonds of P_4 . The structural features of the P_5 core are comparable to $[\text{Ph}_2\text{P}_5]^+$.^{5c} The $\text{P}-\text{P}$ bonds involving the four-coordinate phosphorus center P1 and the $\text{P4}-\text{P5}$ bond (2.1462(7) – 2.164(1) Å) are significantly shorter than the remaining $\text{P}-\text{P}$ bonds (2.2461(8) – 2.2535(8) Å). Comparably short $\text{P}-\text{P}$ distances have been observed in other compounds with cationic four-coordinate phosphorus centers and strained phosphorus cages.^{2i,11} The $\text{P}-\text{N}$ bonds between the tri-coordinate phosphorus atom P6 and N1 or N2 (1.733(2), 1.732(2) Å) are typical for neutral diphosphadiazanes.¹² In contrast, the $\text{P}-\text{N}$ bonds involving the phosphonium center P1 are significantly shorter (1.662(2), 1.669(2) Å).




Figure 2. ORTEP plot of the molecular structure of the cation **3** in **3** $[\text{GaCl}_4] \cdot \text{C}_6\text{H}_5\text{F}$. Thermal ellipsoids at 50% probability (hydrogen atoms, counteranion and $\text{C}_6\text{H}_5\text{F}$ omitted for clarity). Selected bond lengths (Å): P6–C11 2.0788(7), P6–N1 1.733(2), P6–N2 1.732(2), P1–N1 1.669(2), P1–N2 1.662(2), P1–P2 2.1518(7), P1–P3 2.1462(7), P2–P4 2.2535(8), P2–P5 2.2461(8), P3–P4 2.2484(8), P3–P5 2.2472(8), P4–P5 2.164(1).




Figure 3. $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of $3[\text{GaCl}_4]$ and $4[\text{Ga}_2\text{Cl}_7]_2$ (in $\text{C}_6\text{H}_5\text{F}$, C_6D_6 -capillary, 25°C ; 161.94 MHz); very small amounts of unidentified side-products are indicated by an asterisk.^{8b}

The reaction of two equiv. of P_4 with cation **2** and an excess of Lewis acid (**2**, GaCl_3 1:4) in $\text{C}_6\text{H}_5\text{F}$ resulted in the formation of a pale-yellow solution with small amounts of orange-yellow precipitate (Scheme 1). The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of the filtrate is depicted in Figure 3. Beside the resonances of monocation **3**, signals of a new A_2MX_2 spin system ($\delta_{\text{A}} = -341.8$ ppm, $\delta_{\text{M}} = 67.4$ ppm, $\delta_{\text{X}} = 142.3$ ppm; $^1J_{\text{AX}} = -320.3$ Hz, $^1J_{\text{MX}} = -135.2$ Hz, $^2J_{\text{AM}} = 21.9$ Hz) in a ratio of 2:1:2 indicate the formation of a new species in approximately 60% yield.^{8b} The resonances for the A_2MX_2 spin system are consistent with two C_{2v} -symmetric P_5 cages bridged by two imido groups, suggesting the formation of dication **4** (Figure 3). $4[\text{Ga}_2\text{Cl}_7]$ crystallized as a conglomerate with $3[\text{GaCl}_4]$. The postulated structure in solution was confirmed by X-ray diffraction (Figure 4). To our knowledge, dication **4** represents the first structurally characterized example of two homoatomic P_5 cages fused *via* an imido bridge. In the solid state, dication **4** is centrosymmetric, consistent with the A_2MX_2 pattern observed in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum. The bond lengths and angles in the P_5 cores of dication **4** follow a similar trend as observed for monocation **3**. The N_2P_2 ring is planar with a short $\text{P1}-\text{N1}$ bond (1.684(2) Å). The pronounced short character of the $\text{P}-\text{N}$ bonds in the P_2N_2 core might account for an increased reactivity. In solution, dication **4** is not very stable and readily decomposes to an insoluble orange material at room temperature. The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of the reactions mixture after ~ 12 h shows an additional complex set of signals. This indicates the formation of a further oligomer which we believe to be cation $[(\text{DippNP})_3(\text{P}_4)\text{Cl}_2]^+$, presumably resulting from a condensation of cations **2** and **4**. This unusual process is currently being investigated. Details will be reported in a subsequent full paper.

In summary, the synthesis of the unique phosphorus-rich organophosphorus cation **4** has been achieved by stepwise insertion




Figure 4. ORTEP plot of the molecular structure of the cation **4** in $4[\text{Ga}_2\text{Cl}_7]_2$. Thermal ellipsoids at 50% probability (hydrogen atoms and counteranions omitted for clarity). Selected bond lengths (Å): $\text{N1}-\text{P1}$ 1.684(2), $\text{P1}-\text{P2}$ 2.1388(7), $\text{P1}-\text{P3}$ 2.1380(7), $\text{P2}-\text{P4}$ 2.2603(8), $\text{P2}-\text{P5}$ 2.2450(8), $\text{P3}-\text{P4}$ 2.2598(8), $\text{P3}-\text{P5}$ 2.2425(8), $\text{P4}-\text{P5}$ 2.1659(9); [symmetry code: (i) $-x, -y+1, -z$].

of the disguised bifunctional Lewis acid $[\text{DippNP}]_2^{2+}$ into the $\text{P}-\text{P}$ bonds of two P_4 tetrahedra. The utilization of bifunctional phosphonium cations¹³ represents a rational and potentially versatile synthetic method for the assembly of large clusters using P_4 as a building block. Due to the cationic charge such cluster may be amenable to a host of subsequent transformations. Studies directed at the synthesis of further cationic clusters and the reactivity of **3** and **4** are in progress.

Acknowledgment. We gratefully acknowledge the Alexander von Humboldt-Foundation (Feodor Lynen Return Fellowship for J.J.W.), the FCI (Liebig fellowship for J.J.W.), the European Phosphorus Science Network (PhoSciNet CM0802), and the International Research Training Group (IRTG 1444). J.J.W. thanks Prof. F. Ekkehardt Hahn (WWU Münster) for his generous support and advice. We thank Dr. Robert Wolf for helpful discussions.

Supporting Information Available: Full experimental and spectroscopic data for compounds $2[\text{Ga}_2\text{Cl}_7]$ and $3[\text{GaCl}_4] \cdot \text{C}_6\text{H}_5\text{F}$ and selected data for $4[\text{Ga}_2\text{Cl}_7]_2$ (including $^{31}\text{P}-^{31}\text{P}$ DQF COSY), and X-ray crystallographic data for $3[\text{GaCl}_4]$ and $4[\text{Ga}_2\text{Cl}_7]_2$ (CCDC numbers 743848, 743849). This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) (a) Masuda, J. D.; Schoeller, W. W.; Donnadiu, B.; Bertrand, G. *Angew. Chem., Int. Ed.* **2007**, *46*, 7052. (b) Dyker, C. A.; Bertrand, G. *Science* **2008**, *321*, 1050. (c) Masuda, J. D.; Schoeller, W. W.; Donnadiu, B.; Bertrand, G. *J. Am. Chem. Soc.* **2007**, *129*, 14180. (d) Back, O.; Kuchenbeiser, G.; Donnadiu, B.; Bertrand, G. *Angew. Chem., Int. Ed.* **2009**, *48*, 5530.
- (2) (a) Dohmeier, C.; Schnöckel, H.; Robl, C.; Schneider, U.; Ahlrichs, R. *Angew. Chem., Int. Ed. Engl.* **1994**, *33*, 199. (b) Peng, Y.; Fan, H.; Zhu, H.; Roesky, H. W.; Magull, J.; Hughes, C. E. *Angew. Chem., Int. Ed.* **2004**, *43*, 3443. (c) Power, M. B.; Barron, A. R. *Angew. Chem., Int. Ed. Engl.* **1991**, *30*, 1353. (d) Uhl, W.; Benter, M. *Chem. Commun.* **1999**, 771. (e) Fox, A. R.; Wright, R. J.; Rivard, E.; Power, P. P. *117*, 7907; *Angew. Chem., Int. Ed.* **2005**, *44*, 7729. (f) Wiberg, N.; Wörner, A.; Karaghiosoff, K.; Fenske, D. *Chem. Ber.* **1997**, *130*, 135. (g) Lerner, H.-W.; Bolte, M.; Karaghiosoff, K.; Wagner, M. *Organometallics* **2004**, *23*, 6073. (h) Chan, W. T. K.; García, F.; Hopkins, A. D.; Martin, L. C.; McPartlin, M.; Wright, D. S. *Angew. Chem., Int. Ed.* **2007**, *46*, 3084. (i) Xiong, Y.; Yao, S.; Brym, M.; Driess, M. *Angew. Chem., Int. Ed.* **2007**, *46*, 4511.
- (3) Gudat, D. *Eur. J. Inorg. Chem.* **1998**, 1087.
- (4) (a) Krossing, I.; Raabe, I. *Angew. Chem., Int. Ed.* **2001**, *40*, 4406. (b) Gonsior, M.; Krossing, I.; Müller, L.; Raabe, I.; Jansen, M.; van Wuellem, L. *Chem.-Eur. J.* **2002**, *8*, 4475.
- (5) (a) Weigand, J. J.; Burford, N.; Lumsden, M. D.; Decken, A. *Angew. Chem., Int. Ed.* **2006**, *45*, 6733. (b) Weigand, J. J.; Burford, N.; Decken, A. *Eur. J. Inorg. Chem.* **2008**, 4343. (c) Weigand, J. J.; Holthausen, M.; Fröhlich, R. *Angew. Chem., Int. Ed.* **2009**, *48*, 295.
- (6) (a) Michalik, D.; Schulz, A.; Villinger, A.; Weding, N. *Angew. Chem., Int. Ed.* **2008**, *47*, 6465. (b) David, G.; Niecke, E.; Nieger, M.; von der Gönna, V.; Schoeller, W. W. *Chem. Ber.* **1993**, *126*, 1513. (c) Burford, N.; Landry, J. C.; Ferguson, M. J.; McDonald, R. *Inorg. Chem.* **2005**, *44*, 5897. (d) Burford, N.; Conroy, K. D.; Landry, J. C.; Ragana, P. J.; Ferguson, M. J.; McDonald, R. *Inorg. Chem.* **2004**, *43*, 8245.
- (7) Burford, N.; Cameron, T. S.; Conroy, K. D.; Ellis, B.; MacDonald, C. L. B.; Ovars, R.; Phillips, A. D.; Ragogna, P. J.; Walsh, D. *Can. J. Chem.* **2002**, *80*, 1404.
- (8) (a) Only a broad signal for **2** is observed in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum due to fluxional behaviour. The observed chemical shift strongly depends on the GaCl_3 concentration. Cation **2** was isolated as $2[\text{Ga}_2\text{Cl}_7]$ when more than 2 eq. of GaCl_3 are used. (b) For experimental details and spectroscopic data see Supporting Information.
- (9) Budzelaar, P. H. M. *gNMR for Windows*, version 5.0.6.0; NMR Simulation Program; IvorySoft: Budzelaar, 2006.
- (10) Burford, N.; Clyburne, J. A. C.; Chan, M. S. W. *Inorg. Chem.* **1997**, *36*, 3204.
- (11) (a) Niecke, E.; Rüger, R.; Krebs, B. *Angew. Chem., Int. Ed. Engl.* **1982**, *21*, 544. (b) Schoeller, W. W.; Staemmler, V.; Rademacher, P.; Niecke, E. *Inorg. Chem.* **1986**, *25*, 4382. (c) Dyker, C. A.; Burford, N. *Chem. Asian J.* **2008**, *3*, 28, and references therein.
- (12) (a) Vijjilatha, M.; Kumara Swamy, K. C.; Vittal, J. J.; Koh, L. L. *Polyhedron* **1999**, *18*, 2249. (b) Thompson, M. L.; Tarassoli, A.; Haltiwanger, R. C.; Norman, A. D. *J. Am. Chem. Soc.* **1981**, *103*, 6770.
- (13) Davidson, R. J.; Weigand, J. J.; Burford, N.; Cameron, T. S.; Decken, A.; Werner-Zwanziger, U. *Chem. Commun.* **2007**, 4671.

JA906878Q