Tetrahedron Letters 50 (2009) 5834-5837

Contents lists available at ScienceDirect

Tetrahedron Letters

journal homepage: www.elsevier.com/locate/tetlet



Palladium complexes with abnormal *N*-heterocyclic carbene ligands derived from 1,2,3-triazolium ions and their application in Suzuki coupling

Thandavamurthy Karthikeyan, Sethuraman Sankararaman*

Department of Chemistry, Indian Institute of Technology Madras, Chennai 600 036, India

ARTICLE INFO	ABSTRACT
Article history: Received 15 July 2009 Revised 30 July 2009 Accepted 5 August 2009 Available online 9 August 2009	Chiral and achiral pincer type palladium complexes bearing abnormal <i>N</i> -heterocyclic carbene ligands derived from 1,2,3-triazole are reported. Both complexes are effective as catalysts for non-asymmetric Suzuki coupling reaction for the synthesis of biphenyl derivatives. However, both the complexes failed to catalyze the formation of binaphthyl derivatives, by the asymmetric Suzuki coupling in case of the chi- ral complex and the non-asymmetric version in case of the achiral pincer complex. Instead deboronation of arylboronic acid occurred very efficiently leaving the aryl bromide intact.

In 1968 Wanzlick and Ofele almost simultaneously and independently reported the potential of imidazolium-derived *N*-heterocyclic carbenes (NHC) as ligands for transition metals.¹ Although the ability of NHCs to act as efficient ligands by coordination through C2 carbon of the imidazolidene has been well established, isolation and structural characterization of free NHCs were reported only in the early 90s with the discovery of the Arduengo carbenes.² Recently, NHCs derived from deprotonation of imidazolium ions at

the C4 position have been reported.³ Unlike the Wanzlick–Ofele– Arduengo carbenes in this class of NHCs, the carbene center (C4) is not flanked by heteroatoms on both sides and their structures can only be represented as zwitter ions and not in the neutral canonical form (Scheme 1). Such carbenes are termed as abnormal NHCs. Metallation via C4-H activation results in the formation of C4-bound metal carbene complexes (Scheme 1).

A handful of abnormal NHCs from other heterocycles such as pyrazolylidene, isothiozolylidene, pyridylidene, and 1,2,3-triazolylidene have been reported.⁴ Recently, we have reported pyrrolidine-1,2,3-triazole conjugates as organocatalysts for the asymmetric Michael addition reaction with excellent diastereo and enantioselectivities.⁵ During our attempts to prepare the corresponding chiral quaternary ammonium hydroxide of **1**, we stumbled upon formation of the silver complex of abnormal NHC derived from **2** (Scheme 2). During the course of our investigation Albrecht⁶ has reported the synthesis and structural characterization of late transition metal complexes with abnormal NHC ligands derived from 1,2,3-triazolylidenes. Herein we report the first chiral palladium complex bearing abnormal NHC ligand and the first palladium pincer complex bearing abnormal bis-NHC chelating ligand, both based on 1,2,3-triazolylidene, and their utility in Suzuki coupling reactions.

Reaction of chiral triazole derivative 1^5 with methyl iodide yielded the corresponding quaternary ammonium iodide 2 in quantitative yield (Scheme 2). In an attempt to synthesize the corresponding chiral quaternary ammonium hydroxide (3), quaternary ammonium iodide 2 was treated with silver oxide. ¹H NMR spectrum of the crude product revealed the absence of triazole ring proton that appeared at 9.47 ppm for 2. The crude product was identified as the silver complex (4) of the abnormal NHC formed from 2 (Scheme 2). This observation is consistent with the report of Albrecht.⁶ The silver carbene complex (4) was unstable under ambient conditions and also decomposed upon exposure to light. Nevertheless it was characterized by ¹H NMR, ESI-MS, and HRMS. The molecular ion corresponding to the cation part of 4 showed



Scheme 1. Metal complexes derived from normal (*C*2) and abnormal (*C*4) NHCs from imidazolium ion.



^{*} Corresponding author. Tel.: +91 4422574210; fax: +91 4422570545. *E-mail address:* sanka@iitm.ac.in (S. Sankararaman).

^{0040-4039/\$ -} see front matter \odot 2009 Elsevier Ltd. All rights reserved. doi:10.1016/j.tetlet.2009.08.002



Scheme 2. Synthesis of silver complex of abnormal NHC derived from 2.

the isotope peaks that matched with the calculated spectrum. In view of the instability of the silver complex, it was decided to prepare the palladium complex of the abnormal carbene derived from **2**. The NHC complexes of palladium have been shown to be useful in cross-coupling reactions.⁷ Attempted synthesis of the corresponding palladium complex by treatment of **2** with $Pd(OAc)_2$ in DMSO, following the procedure reported by Albrecht,⁶ yielded a mixture of intractable material. Therefore transmetallation of in situ-generated silver carbene complex **4** was carried out.

Treatment of **2** with silver oxide followed by $[Pd(Cl)_2(CH_3CN)_2]$ in dichloromethane afforded the corresponding palladium complex **5** (Scheme 3) in 67% yield as an air stable pale yellow solid. It was purified by chromatography over silica gel and characterized by IR, ¹H and ¹³C NMR, ESI-MS, and HRMS data.

Attempts to grow single crystals of 5 yielded only a microcrystalline solid that was unsuitable for X-ray crystallographic analysis. Palladium catalysts bearing NHC ligands have been used for crosscoupling reactions, in particular Suzuki and Heck coupling reactions.⁷ NHC ligands are good alternative to the conventional phosphine ligands, and catalysts bearing NHC ligands react with a high degree of efficiency in C-C bond forming reactions compared to ligandless catalysts. Although complex 5 is chiral it was first tested for its catalytic activity in simple non-asymmetric Suzuki reactions for the synthesis of substituted biphenyls. Various substituted aryl bromides were treated with phenylboronic acid and 4-methoxyphenylboronic acid in THF in the presence of catalytic amounts of 5. The catalyst loading varied from 2 to 20 mol % depending upon the substrates. The coupling products were obtained in good vields and the results are summarized in Table 1. In Table 1, entries 3, 7, and 8 are noteworthy because methoxy-substituted derivatives are generally less reactive in Suzuki coupling and using the catalyst 5, high yields of the coupling products were obtained. To rule out the possibility of the reaction being catalyzed by leached



Scheme 3. Synthesis of palladium complex 5 from 2 via in situ-generated silver complex.

out Pd(0), the reaction of phenylboronic acid with 4-cyanobromobenzene was carried out in the presence of mercury. Though the reaction took longer time for completion (9 vs 14 h), the coupling product was obtained in 98% yield, indicating that the reaction was catalyzed by **5**. Attempted coupling of 1,4-phenylenediboronic acid with 4-methoxybromobenzene using **5** did not proceed and only starting materials were recovered under the conditions given in Table 1. The more reactive 2-bromo-5-*tert*-butylisophthalaldehyde (Table 1, entry 4, ArBr) also failed to react with 1,4-phenylenediboronic acid.

As stated earlier, complex 5 is chiral and it is potentially useful for the asymmetric C-C cross-coupling reactions. We focused our attention on the asymmetric Suzuki coupling for the synthesis of chiral binaphthalene derivatives. Although palladium catalysts bearing other NHC ligands have been successfully used in nonasymmetric Suzuki coupling,^{8,9} to this date there has been no report on the asymmetric version of Suzuki coupling using NHC ligand based catalysts, especially of the sterically hindered substrates. Therefore, we wanted to study the utility of complex 5 for the asymmetric synthesis of chiral binaphthalene derivatives. Accordingly, 1-bromo-2-methoxynaphthalene (6) was treated with 2-methoxy-1-naphthylboronic acid (7) in the presence of K_2CO_3 and 10 mol % of 5 in THF. To our dismay only deboronation of 7 to yield 2-methoxynaphthalene (8) was observed and 6 remained intact. The same deboronation was observed when 2-benzyloxy-1-bromonaphthalene (9) was reacted with 2-benzyloxy-1-naphthylboronic acid (10) to yield 2-benzyloxynaphthalene (11) and 9 was recovered. Also reaction of 1-iodo-2-naphthol with 7 led to only deboronation, and 2-methoxynaphthalene (8) and 1-iodo-2naphthol were recovered from the reaction mixture. Attempted coupling of 1-naphthylboronic acid with 6 yielded 1,1-binaphthylene as the product and 6 was recovered back from the reaction. It is well documented that sterically demanding boronic acids with two ortho substituents are prone to deboronation during Suzuki coupling reaction.¹⁰ Nevertheless, some successful coupling of sterically hindered boronic acids has been reported under rigorously anhydrous conditions. Our attempts to carry out asymmetric Suzuki coupling using rigorously dried THF and DME under various conditions failed to yield the desired binaphthalene derivatives. Only deboronated products were obtained in nearly quantitative yields in all the reactions (Table 2). Deboronation in Suzuki coupling of sterically hindered substrates is explained on the basis of formation of trans-diarylpalladium(0) intermediate in which C-C coupling is disfavored compared to *cis*-diarylpalladium(0) intermediate. As a result of the formation of trans isomer, the catalytic cycle is stagnated and the reaction does not proceed further to the reductive elimination step to complete the catalytic cycle. Under these conditions, the boronic acid undergoes deboronation in the presence of the base and often due to the presence of adventitious water in the reaction. This problem can be addressed by using a chelating ligand for the palladium complex which prevents the conversion of the *cis*-diarylpalladium(0) intermediate to the trans isomer. Alternatively, the transmetallation of the boronic acid could be a slow step, especially in sterically demanding substrates, leading to deboronation reaction. To rule out the formation of the trans-diarylpalladium(0) intermediate, we proceeded to synthesize an achiral version of a pincer type palladium complex. Methylation of the bis-triazole derivative 12 (obtained by the addition of benzyl azide to 1.3-diethynylbenzene) gave the bis-quaternary ammonium iodide 13 in 95% yield. It was further treated with silver oxide followed by PdCl₂(CH₃CN)₂ to afford complex 15 (Scheme 4). The intermediate silver carbene complex 14 was unstable for isolation and complete characterization. Nevertheless, it was characterized by ESI-MS, and HRMS data which corresponded to C₂₆H₂₄N₆Ag, the cationic part of the silver carbene complex 14. Complex 15 was characterized by IR, NMR, ESI-MS and

Table 1

Synthesis of biphenyls by Suzuki coupling using complex 5^a

ArBr + Ar'B(OH)₂ $\frac{5 (2 \text{ to } 20 \text{ mol}\%)}{K_2 \text{CO}_3, 70-75 \text{ °C}}$ Ar-Ar'

Entry	ArBr	Ar'B(OH) ₂	Ar-Ar'	5 (mol %)	Time (h)	Yield (%)
1 ^b	1-Bromonaphthalene	PhB(OH) ₂	1-Phenylnaphthalene	20	16	73
2	1,2-Dibromobenzene	PhB(OH) ₂	o-Terphenyl	20	18	60
3	1-Bromo-2-methoxynaphthalene	PhB(OH) ₂	2-Methoxy-1-phenylnaphthalene	10	12	93
4	OHC Bu-t	PhB(OH) ₂	OHC Bu-t	5	7	84
5	1-Bromo-4-nitrobenzene	PhB(OH) ₂	4-Nitrobiphenyl	5	8	90
6	4-Bromobenzonitrile	PhB(OH) ₂	4-Cyanobiphenyl	5	9	90
7 ^c	1-Bromo-4-methoxybenzene	4-Methoxy-phenylboronic acid	4,4'-Dimethoxybiphenyl	2	40	79
8 ^c	1-Bromo-4-methylbenzene	4-Methoxy-phenylboronic acid	4-Methoxy-4'-methylbiphenyl	2	40	96

^a Entries 1–6, 0.35 mmol scale, 1 equiv ArBr, 2 equiv PhB(OH)₂, 3 equiv K₂CO₃, entries 7–8, 0.88 mmol scale, 1 equiv ArBr, 1.2 equiv 4-methoxyphenylboronic acid, 3 equiv K₂CO₃. ^b CsF as base.

^c Yield based on 37% conversion for entry 7 and 30% conversion for entry 8.

Table 2

Attempted asymmetric Suzuki coupling for the synthesis of binaphthalene derivatives^a



Entry	ArBr	$ArB(OH)_2$	5 (mol %)	Base/solvent	Product
1	6	7	10	K ₂ CO ₃ /THF	8
2	9	10	10	K ₂ CO ₃ /THF	11
3	6	7	20	KOBu-t/DME	8
4	9	10	20	KOBu-t/DME	11
5	6	7	10	Cs ₂ CO ₃ /DME	8
6 ^b	6	1-NpB(OH) ₂	10	K ₂ CO ₃ /THF	Np-Np

1 equiv of ArBr, 2 equiv of ArB(OH)₂ and 3 equiv of base.

^b 1-NpB(OH)₂ = 1-naphthylboronic acid and Np–Np = 1,1-binaphthylene.

HRMS. The absence of triazolium ring proton resonance at 9.47 ppm in the ¹H NMR spectrum indicated that both the rings had undergone deprotonation resulting in the formation of the corresponding biscarbene leading to the formation of 15 via 14. The molecular ion peak in MS corresponded to C₂₆H₂₅N₆Cl₂Pd, [M+1] ion, and showed the isotope distribution pattern expected for this formula, based on the calculated spectrum. Complex 15 was tested for its catalytic activity in Suzuki coupling. Coupling of 4-bromonitrobenzene and 4-bromobenzonitrile with phenylboronic acid vielded 4-nitro- and 4-cyanobiphenyl, respectively, in >90% yields. However, attempted coupling of 1-bromo-2-methoxynaphthalene (6) and 2-methoxynaphthalen-1-ylboronic acid (7) using 15 and Cs₂CO₃ in DME resulted only in deboronation of 7 to yield 2-methoxynaphthylene (8) and 6 remained intact in the reaction. To our disappointment, even the use of the pincer complex **15** as catalyst in the Suzuki reaction did not result in the formation of the desired binaphthyl derivative. We speculate that the failure of the Suzuki coupling to yield binaphthyl derivatives using 5 and 15 as pre-cat-



Scheme 4. Synthesis of pincer palladium complex 15.

alysts might be due to slow transmetallation of the sterically demanding boronic acids, leading to deboronation reaction.

In conclusion, we report the first example of a chiral palladium complex (5) bearing an abnormal NHC ligand based on 1,2,3-triazolylidene precursor and the first example of the corresponding pincer complex (15) derived from bis-1,2,3-triazolylidene precursor. We have demonstrated that both the complexes are useful as catalysts for the non-asymmetric Suzuki coupling reaction for the synthesis of biphenyl derivatives. However, both the complexes failed to catalyze the formation of binaphthyl derivatives, the asymmetric version of Suzuki coupling in the case of 5 and the ordinary version in the case of 15. Under the reaction conditions deboronation of arylboronic acid occurred very efficiently leaving the aryl bromide intact in the reaction.

Acknowledgments

We thank DST, New Delhi, for financial support, Department of Chemistry and SAIF, IIT Madras for spectral data and IIT Madras for a fellowship to T.K.

Supplementary data

Supplementary data (detailed experimental procedure, copies of ¹H NMR, ¹³C NMR, mass spectrum of palladium complexes, characterization data for the Suzuki products are available in supplementary material) associated with this article can be found, in the online version, at doi:10.1016/j.tetlet.2009.08.002.

References and notes

- Wanzlick, H. W.; Schonherr, H. J. Angew. Chem., Int. Ed. Engl. 1968, 7, 141–142; Ofele, K. J. Organomet. Chem. 1968, 12, P42–P43.
- Arduengo, A. J., III; Harlow, R. L.; Kline, M. J. Am. Chem. Soc. 1991, 113, 361–363.
 Grundemann, S.; Kovacevic, A.; Albrecht, M.; Faller, J. W.; Crabtree, R. H. Chem.
- Commun. 2001, 2274–2275; Chianese, A. R.; Kovacevic, A.; Zeglis, B. M.; Faller, J.

W.; Crabtree, R. H. Organometallics **2004**, 23, 2461–2468; Albrecht, M. Chem. Commun. **2008**, 3601–3610; Lebel, H.; Janes, M. K.; Charette, A. B.; Nolan, S. P. J. Am. Chem. Soc. **2004**, 126, 5046–5047; Heckenroth, M.; Kluser, E.; Neels, A.; Albrecht, M. Dalton Trans. **2008**, 6242–6249.

- Arnold, P. L.; Pearson, S. Coord. Chem. Rev. 2007, 251, 596–609; Schuster, O.; Yang, L.; Raubenheimer, H. G.; Albrecht, M. Chem. Rev., doi:10.1021/cr8005087.
- Karthikeyan, T.; Sankararaman, S. Tetrahedron: Asymmetry 2008, 19, 2741– 2745.
- Mathew, P.; Neels, A.; Albrecht, M. J. Am. Chem. Soc. 2008, 130, 13534– 13535.
- Marion, N.; Nolan, S. P. Acc. Chem. Res. 2008, 41, 1440–1449; Kantchev, E. A. B.; O'Brien, C. J.; Organ, M. G. Angew. Chem., Int. Ed. 2007, 46, 2768–2813.
- Wurtz, S.; Glorius, F. Acc. Chem. Res. 2008, 41, 1523–1533; Marion, N.; Nolan, S. P. Acc. Chem. Res. 2008, 41, 1440–1449.
- Wallace, T. W. Org. Biomol. Chem. 2006, 4, 3197–3210; Zhang, T.; Wang, W.; Gu, X.; Shi, M. Organometallics 2008, 27, 753–757.
- For mechanistic scheme explaining deboronation reaction see: Cammidge, A. N.; Crpy, K. V. L. J. Org. Chem. 2003, 68, 6832–6835.