This article was downloaded by: [Umeå University Library] On: 17 August 2014, At: 15:09 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



# Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

### An Unusual Reaction of Triphenylphosphine with Dichlorodinitrobenzofuroxan

Irina V. Galkina<sup>a</sup>, Elena V. Tudriy<sup>a</sup>, Olga N. Kataeva<sup>ab</sup>, Luiza M. Yusupova<sup>a</sup>, Heinrich Luftmann<sup>c</sup> & Vladimir I. Galkin<sup>a</sup>

<sup>a</sup> A. M. Butlerov Chemistry Institute of the Kazan State University , Kazan, Russia

<sup>b</sup> A. E. Arbuzov Institute of Organic and Physical Chemistry of the Russian Academy of Sciences , Kazan, Russia

<sup>c</sup> Institute of Organic Chemistry, University of Münster , Münster, Germany Published online: 26 Mar 2009.

To cite this article: Irina V. Galkina , Elena V. Tudriy , Olga N. Kataeva , Luiza M. Yusupova , Heinrich Luftmann & Vladimir I. Galkin (2009) An Unusual Reaction of Triphenylphosphine with Dichlorodinitrobenzofuroxan, Phosphorus, Sulfur, and Silicon and the Related Elements, 184:4, 987-991, DOI: <u>10.1080/10426500902719636</u>

To link to this article: <u>http://dx.doi.org/10.1080/10426500902719636</u>

#### PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views

expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <u>http://www.tandfonline.com/page/terms-and-conditions</u>



## An Unusual Reaction of Triphenylphosphine with Dichlorodinitrobenzofuroxan

Irina V. Galkina,<sup>1</sup> Elena V. Tudriy,<sup>1</sup> Olga N. Kataeva,<sup>1,2</sup> Luiza M. Yusupova,<sup>1</sup> Heinrich Luftmann,<sup>3</sup> and Vladimir I. Galkin<sup>1</sup>

<sup>1</sup>A. M. Butlerov Chemistry Institute of the Kazan State University, Kazan, Russia

<sup>2</sup>A. E. Arbuzov Institute of Organic and Physical Chemistry of the Russian Academy of Sciences, Kazan, Russia

<sup>3</sup>Institute of Organic Chemistry, University of Münster, Münster, Germany

A novel phosphorylation reaction of dichlorodinitrobenzofuroxan by triphenylphosphine has been studied. An unusual course of the reaction with the participation of benzofuroxan nitro groups as reactive centers has been established. It is shown that as final product of the reaction, a stable diphenolate diphosphabetaine with strong participation of the mesomeric ketoylide structure is formed. The structure of the product has been confirmed by X-ray diffraction.

Keywords Dichlorodinitrobenzofuroxan; NO generation; phosphorylation; triphenylphosphine

#### INTRODUCTION

5,7-Dichloro-4,6-dinitrobenzofuroxan (1) is a highly reactive system containing the electrophilic benzofuroxan cycle, which is prone to nucleophilic substitution, owing to the presence of electron-acceptor substituents and easy leaving groups. In particular, the reactions of chlorine substitution by various amines have been frequently investigated.<sup>1</sup> It is essential that amination products as well as initial furoxane **1** 

Received 28 December 2007; accepted 31 January 2008.

Dedicated to Professor Marian Mikołajczyk from the CBMiM PAN in Łódź, Poland, on the occasion of his 70th birthday.

The authors are thankful to Dr. Roland Fröhlich for providing the opportunity to carry out a single crystal X-ray diffraction experiment at the University of Münster.

Address correspondence to Vladimir I. Galkin, A. M. Butlerov Chemistry Institute of the Kazan State University, Kremlevskaya 18, Kazan 420008, Russia. E-mail: vig54@mail.ru possess pronounced antibacterial and fungicidal activity.<sup>2,3</sup> Introduction of biogenic organophosphorus substituents to the benzofuroxan ring might even increase the activity of these compounds; however, no data on the phosphorylation of benzofuroxan is available in the literature until now.

#### **RESULTS AND DISCUSSION**

In this article, we report the interaction of benzofuroxan 1 with triphenylphosphine. The reaction proceeds readily at room temperature to give the bisphosphorylation product, the ratio of 1 to triphenylphosphine being 1:2.5. The reaction most probably is not a simple nucleophilic substitution, as it does not stop upon the formation of 5,7-(bis-triphenylphosphonio)-4,6-dinitrobenzofuroxanyldichloride (2), but rather continues with the elimination of two nitrogen oxide and chlorine molecules, as is shown in Scheme 1. The final product 3 can be described by a series of mesomeric structures.



SCHEME 1 Phosphorylation of benzofuroxan 1.

The structure of **3** has been determined by X-ray single-crystal diffraction (Figure 1) and mass spectrometry.

The distribution of the bond lengths shows that the molecular structure of **3** in crystal is closer to the ylide mesomeric form. The average value for the ylide P=C bond (1.72 Å) derived from the Cambridge



SCHAKAL

**FIGURE 1** Molecular structure of **3** in the crystal. Selected bond lengths: C1A-C5A 1.47(1), C1A-C5B 1.447(8), C5A-C6A 1.431(6), C5B-C6B 1.432(6), C6A-C7 1.457(5), C6B-C7 1.441(6), C7=O7 1.259(4), P9-C6A 1.70(1), P-Ph(average) 1.809(3) Å.

Crystallographic Database<sup>4</sup> is the same within experimental errors as in compound **3**. The C=O bonds are slightly elongated, while the C-C bonds in the benzo-ring are shorter than typical C-C bonds. The molecule lies on a special position on a symmetry plane *m*, implying the symmetrical positions of the triphenylphosphine groups and the disorder of the furoxan cycle. The latter causes low accuracy of the geometrical parameters. The cavity between the triphenylphosphine substituents is filled with disordered water molecules which are bound to the C=O and N→O bonds of the neighboring molecules via hydrogen bonds.

In the mass spectrum (ESI-MS positive), the observed peak at m/z 689 corresponds to the protonated molecule  $C_{42}H_{31}N_2O_4P_2$  [M + H<sup>+</sup>].

#### **EXPERIMENTAL**

#### **Compound 3**

To a solution of  $Ph_3P$  (1.64 g, 6.25 mmol) in 10 mL of 1:1 (in volume) mixture of EtOH and  $Et_2O$ , a solution of 5,7-dichloro-4,

6-dinitrobenzofuroxan (0.73 g, 2.5 mmol) was added dropwise in 20 mL of the same solvent mixture. The reaction mixture was stirred under argon atmosphere for 2.5 h at r.t. After 2 months, the precipitate was collected by filtration, washed with the same solvent mixture (2 × 30 mL), and dried in vacuo, affording compound **3** as a white solid; 1.28 g (75%). DSC/TG: mp 235°C (dec.). <sup>31</sup>P NMR (DMSO-*d*<sub>6</sub>):  $\delta = 23.7$  (s). ESI-MS: Mass spectra were measured with a Quattro LCZ (Waters-Micromass, Manchester, UK) with nanospray inlet. ESI-MS positive: The observed peak at *m*/*z* 689 corresponds to the protonated molecule C<sub>42</sub>H<sub>31</sub>N<sub>2</sub>O<sub>4</sub>P<sub>2</sub> [M + H<sup>+</sup>]. Anal. Calcd. for C<sub>42</sub>H<sub>30</sub>N<sub>2</sub>O<sub>4</sub>P<sub>2</sub>: C, 73.26; H, 4.36; N, 4.07; P, 9.01. Found: C, 73.33; H, 4.27; N, 4.21; P, 9.36.

#### X-Ray Structure Determination

Data were collected with Nonius Kappa CCD diffractometer equipped with a rotating anode generator Nonius FR591. Programs used: data collection COLLECT (Nonius B.V., 1998), data reduction Denzo-SMN,<sup>5</sup> absorption correction Denzo,<sup>6</sup> structure solution SHELXS-97,<sup>7</sup> structure refinement by full-matrix least-squares against  $F^2$  using SHELXL-97,<sup>8</sup> and the absolute configuration was assessed by Flack's enantiopole parameter. CCDC 652720 contains the supplementary crystallographic data for this article. These data can be obtained free of charge at www.ccdc.cam.ac.uk/conts/retrieving.html [or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44(1223)336033, E-mail: deposit@ccdc.cam.ac. uk].

#### X-Ray Crystal Data for 3

Formula  $C_{42}H_{30}N_2O_4P_2$  H<sub>2</sub>O, M = 706.64, colorless crystal 0.35 × 0.30 × 0.06 mm, a = 20.5066(5), b = 9.9288(2), c = 9.1246(1) Å, V = 1857.82(6) Å<sup>3</sup>,  $\rho_{calc} = 1.263$  g cm<sup>-3</sup>,  $\mu = 0.164$  mm<sup>-1</sup>, empirical absorption correction (0.945  $\leq$  T  $\leq$  0.990), Z = 2 (molecule in a special position on plane *m*), orthorhombic, space group Pmn2<sub>1</sub> (No. 31),  $\lambda = 0.71073$  Å, T = 198 K,  $\omega$  and  $\varphi$  scans, 24053 reflections collected ( $\pm$  h,  $\pm$  k,  $\pm$  l), [(sin $\theta$ )/ $\lambda$ ] = 0.66 Å<sup>-1</sup>, 4357 independent (R<sub>int</sub> = 0.073) and 3316 observed reflections [I  $\geq$  2  $\sigma$ (I)], 268 refined parameters, R = 0.055, wR2 = 0.134, max. residual electron density 0.75 (-0.31) e Å<sup>3</sup>, hydrogen atoms calculated and refined as riding atoms.

#### REFERENCES

- [1] W. R. Roger and W. P. Norris, Aust. J. Chem., 38, 435 (1985).
- [2] G. P. Sharnin, R. I. Mukharlyamov, and V. V. Golovin, Zh. Org. Chem., 19, 2358 (1983).
- [3] L. M. Yusupova, I. F. Falyakhov, L. V. Spatlova, and T. V. Garipova, Pat. 2255935 (Russia) (2005).
- [4] Cambridge Structural Database System. Version 5.27. Cambridge Crystallographic Data Centre, Cambridge, UK (2006).
- [5] Z. Otwinowski and W. Minor, Methods Enzymol., 276, 307 (1997).
- [6] Z. Otwinowski, D. Borek, W. Majewski, and W. Minor, Acta Crystallogr., Sect. A: Found. Crystallogr., A59, 228 (2003).
- [7] G. M. Sheldrick, Acta Crystallogr., Sect. A: Found. Crystallogr., A46, 467 (1990).
- [8] G. M. Sheldrick, Universität Göttingen, 1997.