Ene Reaction of Arylallyl Alkenes with C₆₀. A Mechanistic Approach

Nikos Chronakis and Michael Orfanopoulos*

Department of Chemistry, University of Crete, Iraklion 71409, Crete, Greece orfanop@chemistry.uch.gr

Received August 2, 1999

ABSTRACT



The ene reaction of arylallyl alkenes with C_{60} occurs either by a concerted mechanism or by the reversible formation of a charged or a dipolar intermediate, followed by the C–H(D) breakage in a rate-limiting step.

The isolation of C_{60} in large quantities¹ triggered a large number of investigations, over the past 10 years, on its chemical reactivity and physical properties. Several research groups have reported the preparation and isolation of well characterized [4 + 2],² [3 + 2],^{3,4} and $[2 + 2]^{5-8}$ monoadducts of C_{60} . However, the ene functionalization of C_{60}

(3) (a) Suzuki, T.; Li, Q.; Khemani, K. C.; Wudl, F.; Almarsson, Ö. Science 1991, 254, 1186. (b) Prato, M; Suzuki, T.; Wudl, F.; Lucchini, V.; Maggini, M. J. Am. Chem. Soc. 1993, 115, 7876. (c) Prato, M.; Bianco, A.; Maggini, M.; Scorrano, G.; Toniolo, C.; Wudl, F. J. Org. Chem. 1993, 58, 5578. (d) Maggini, M.; Scorrano, G.; Bianco, A.; Toniolo, C.; Sijbesma, R. P.; Wudl, F. J. Chem. Soc., Chem. Commun. 1994, 305. (e) Prato, M.; Li, Q.; Wudl, F.; Lucchini, V. J. Am. Chem. Soc. 1993, 115, 1148. (f) Maggini, M.; Scorrano, G.; Prato, M. J. Am. Chem. Soc. 1993, 115, 9798.

(4) (a) Isaacs, L.; Diederich, F. Helv. Chem. Acta **1993**, 76, 2454. (b) Diederich, F.; Isaacs, L.; Philp, D. J. Chem. Soc., Perkin. Trans. 1 **1994**, 391. (c) Isaacs, L.; Wehrsig, A.; Diederich, F. Helv. Chim. Acta **1993**, 76, 1231. (d) Vasella, A.; Uhlmann, P.; Waldraff, C. A.; Diederich, F.; Thilgen, C. Angew. Chem., Int. Ed. Engl. **1992**, 31, 1388. (e) Anderson, H. L.; Faust, R.; Rubin, Y.; Diederich, F. Angew. Chem., Int. Ed. Engl. **1994**, 33, 1366. (f) An, Y. Z.; Rubin, Y.; Schaller, C.; McEivany, S. W. J. Org. Chem. **1994**, 59, 2927.

(5) Hoke, S. H., II; Molstad, J.; Dilattato, D.; Jay, M. J.; Carlson, D.; Kahr, B.; Cooks, R. G. J. Org. Chem. **1992**, *57*, 5069.

10.1021/ol9902040 CCC: \$18.00 © 1999 American Chemical Society Published on Web 11/13/1999

has received less attention and, to our knowledge, only three reports describe the isolation and characterization of ene monoadducts. For example, Wu and Komatsu reported⁹ independently the first examples of ene addition of 1-heptene, 4-propenylanisole, and 3,5-di-*tert*-butyl-4-(trimethylsiloxy)allylbenzene to C₆₀. Later, Miles and co-workers¹⁰ reported an additional example of the ene reaction of C₆₀ with 3-methylene-2,3-dihydrofuran.

Our previous mechanistic studies of the ene reaction of several enophiles with alkenes,¹¹ as well as the recent reports of [2 + 2] cycloadditions⁸ of aryl alkenes and dienes to C₆₀, prompted us to investigate for the first time the mechanism

ORGANIC LETTERS 1999 Vol. 1, No. 12 1909–1912

^{(1) (}a) Kratschmer, W.; Fostiropoulos, K.; Huffman, D. R. *Chem. Phys. Lett.* **1990**, *170*, 167. (b) Scrivens, W. A.; Bedworth, P. V.; Tour, J. M. J. Am. Chem. Soc. **1992**, *114*, 7917.

^{(2) (}a) Hirsch, A. Synthesis 1995, 895. (b) Kräutler, B.; Puchberger, M. Helv. Chim. Acta 1993, 76, 1626. (b) Rubin, Y.; Khan, S.; Freedberg, D. I.; Yeretzian, C. J. Am. Chem. Soc. 1993, 115, 344. (c) Linssen, T. G.; Dürr, K.; Hirsch, A.; Hanack, M. J. Chem. Soc., Chem. Commun. 1995, 103. (d) Khan, S. I.; Oliver, A. M.; Paddon-Row: M. N.; Rubin, Y. J. Am. Chem. Soc. 1993, 115, 4919. (e) Gügel, A.; Kraus, A.; Spikermann, J.; Belik, P.; Müllen, K. Angew. Chem., Int. Ed. Engl. 1994, 33, 559. (f) Iyoda, M.; Sultana, F.; Sasaki, S.; Yoshida, M. J. Chem. Soc., Chem. Commun. 1994, 1929. (g) Zhang, X.; Foote, C. S. J. Org. Chem. 1994, 59, 5235.

^{(6) (}a) Zhang, X.; Romero, A.; Foote, C. S. J. Am. Chem. Soc. **1993**, 115, 11024. (b) Zhang, X.; Romero, A.; Foote, C. S. J. Am. Chem. Soc. **1995**, 117, 4271. (c) Zhang, X.; Fan, A.; Foote, C. S. J. Org. Chem. **1996**, 61, 5456.

^{(7) (}a) Wilson, S. R.; Kaprinidis, N.; Wu, Y.; Schuster, D. I. J. Am. Chem. Soc. **1993**, 115, 8495. (b) Wilson, S. R.; Wu, Y.; Kaprinidis, N.; Schuster, D. I. J. Org. Chem. **1993**, 58, 6548.

^{(8) (}a) Vassilikogiannakis, G.; Orfanopoulos, M. J. Am. Chem. Soc. **1997**, 119, 7394. (b) Vassilikogiannakis, G.; Orfanopoulos, M. Tetrahedron Lett. **1997**, 38, 4323. (c) Vassilikogiannakis, G.; Chronakis, N.; Orfanopoulos, M. J. Am. Chem. Soc. **1998**, 120, 9911.

^{(9) (}a) Wu, S.; Shu, L.; Fan, K. *Tetrahedron Lett.* **1994**, *35*, 919. (b) Komatsu, K.; Murata, Y.; Sugita, N.; S. M. Wan, T. *Chem. Lett.* **1994**, 635.

⁽¹⁰⁾ Miles, W. H.; Smiley, P. M. J. Org. Chem. 1996, 61, 2559.

⁽¹¹⁾ Stephenson, L. M.; Grdina, M. J.; Orfanopoulos, M. Acc. Chem. Res. **1980**, 13, 419. (b) Orfanopoulos, M.; Smonou, I.; Foote. C. S. J. Am. Chem. Soc. **1990**, 112, 3607. (c). Orfanopoulos, M.; Elemes, Y.; Stratakis, M.; Jensen F. J. Am. Chem. Soc. **1991**, 113, 3801. (d) Stratakis, M.; Orfanopoulos, M.; Foote, C. S. J. Org. Chem. **1998**, 63, 1315. (e) Stratakis, M.; Vassilikogiannakis, G.; Orfanopoulos, M. Tetrahedron Lett. **1998**, 39, 2393.



Figure 1. Intramolecular isotope effects in the ene reaction of $1-d_1$ and $2-d_1$ with C_{60} .

of C_{60} -ene functionalization. For this purpose we have studied and report here both intramolecular (product) and intermolecular (kinetic) isotope effects of the ene reaction of C_{60} with arylallyl alkenes.

It is well established that isotope effect measurements are a powerful tool for distinguishing between concerted and stepwise reaction pathways.^{11,12} To determine the intramolecular isotope effect, a mixture of C_{60} and a 100-fold excess of 3-phenyl-1-propene-3-d₁, **1-d₁**,¹³ were dissolved in deoxygenated toluene and heated at 200 °C in a thick glass tube equipped with a rotaflo valve and magnetic stirrer. The reaction was interrupted at various conversions and analyzed by HPLC equipped with a Separon $C_{18}\xspace$ reversed phase column.



A 24% yield of the ene monoadduct was obtained after 40 h, based on the recovered C₆₀. The adduct was purified by flash column chromatography (toluene/hexane 2/1) and characterized by ¹H NMR and FAB-MS [m/z (M + 1) 840.92].

A substantial intramolecular (product) isotope effect $k_{\rm H}/k_{\rm D} = 1.80$ (average) was measured by careful integration of the methylenic hydrogen absorptions at 4.46 ppm and the hydrogen on the fullerene carbon at 6.69 ppm, Figure 1.

The intermolecular (kinetic) isotope effect $k_{\rm H}k_{\rm D}$ was obtained from the equimolar competition of $1-d_0$ vs $1-d_2^{13}$ with C₆₀. The experimental conditions were similar to those described previously for the intramolecular competition. Again, a substantial kinetic isotope effect (average $k_{\rm H}k_{\rm D} =$ 2.0) was found. The results are shown in Table 1. These effects (substantial intra- and intermolecular isotope effects) are comparable to previously reported isotope effects for other ene reactions which were interpreted as supporting a concerted reaction mechanism.¹² It should be noted that the determination of the primary isotope effects was possible only from the integration of the methylenic absorption at 4.46 ppm and the C₆₀-H absorption at 6.69 ppm (Figure

^{(12) (}a) Song, Z.; Chrisope, D. R.; Beak, P. J. J. Org. Chem. 1987, 52, 3940 and references therein. (b) Snider, B. B.; Ron, E. J. Am. Chem. Soc. 1985, 107, 8160. (c) Stephenson, L. M.; Orfanopoulos, M. J. Org. Chem. 1981, 46, 2200.

⁽¹³⁾ Alkenes 1-d₁, 1-d₂, 2-d₁, and 2-d₂ were prepared as follows: reduction of benzaldehyde with LiAlD₄ followed by chlorination with thionyl chloride gave the corresponding benzyl chloride- d_1 . The Grignard reagent of benzyl chloride- d_1 coupled with vinyl bromide in the presence of NiCl₂/Ph₃P¹⁴ catalyst, to give $1-d_1$, in 70% yield. MS [m/z 118 (M - 1, 100)]. Reduction of methyl benzoate with $LiAlD_4$ gave benzyl chloride- d_2 , and then following the procedure described for the preparation of 1-d₁, **1-d**, was prepared in reasonable yield. MS $[m/z \ 119 \ (M - 1, \ 100)]$. NBS monobromination of methyl p-methoxyphenylacetate followed by LiAlD4 reduction gave 2-(p-methoxyphenyl)ethanol-1,1,2-d3 which was then converted to the corresponding bromide by using Ph3P·Br2 as the brominating reagent. Reaction of 2-(p-methoxyphenyl)ethyl-1,1,2-d3 bromide with Ph3P gave the corresponding Wittig salt. The α -deuterium atoms of this salt were exchanged for hydrogens by treating the Wittig salt with NaOH/CH3OH.15 The ylide of the previously prepared Wittig salt was coupled with paraformaldehyde to afford $2-d_1$. MS [m/z 149 (M⁺, 100)]. Treatment of the p-methoxyphenylacetic acid with NaOH in D₂O at 160 °C, and after repeating this process twice gave p-methoxyphenylacetic acid- d_2 , 2-d₂, with more than 96% deuterium atom incorporation. Esterification of the above deuterated acid followed by LiAlH4 reduction afforded 2-(pmethoxyphenyl)ethanol- d_2 , 2- d_2 , and then, following the procedure described for the preparation of $2-d_1$ (delete the step of D exchange), $2-d_2$ was prepared. MS [m/z 150 (M⁺, 100)].

Table 1. Isotope Effects for the Ene Reaction of 3-Phenyl-1-propene and 3-(p-Methoxyphenyl)-1-propene with C_{60}

00						
reaction	cubatnoto	colvent	<i>t</i> ,	convn, ^a	<i>Т</i> ,	Ir /Ir b
type	substrate	solvent	п	70	U	KH/KD
intramol.	1-d ₁	toluene	40	24	200	1.8 ± 0.1
intermol.	$1 - d_0 / 1 - d_2$	toluene	40	21	200	2.0 ± 0.1
intramol.	2-d ₁	DCB^{c}	3	24	180	1.5 ± 0.1^d
intermol.	$2 - d_0/2 - d_2$	DCB^{c}	3	25	180	1.8 ± 0.1^d

^{*a*} Based on recovered C₆₀. ^{*b*} Determined by ¹H NMR integration of the methylenic and C₆₀-H protons. ^{*c*} 1,2-Dichlorobenzene. ^{*d*} Determined as previously^{*b*} and by ¹H NMR integration of the vinylic and aromatic protons.

1). It was difficult to measure the isotope effect from the integration of the vinylic hydrogen absorption (between 7.26 and 7.05 ppm) of the ene adducts, because the phenyl hydrogens of traces of toluene, which was the solvent used in the reaction, absorb in the same chemical shift region. Several attempts to remove completely toluene from the ene adduct were unsuccessful. A crystalline form of C_{60} in which a number of toluene molecules are occluded is probably formed.¹⁶

To confirm both the intra- and the intermolecular isotope effect values found with the 3-phenyl-1-propene system, the 3-(*p*-methoxyphenyl)-1-propene-3- d_1 , **2-d**₁, and its dideuterated analogue 3-(p-methoxyphenyl)-1-propene-3,3-d₂, 2-d₂, were prepared¹³ and studied. In this case, the ene reactions of $2-d_1$ and $2-d_0/2-d_0$ with C₆₀, in separate experiments (intraand intermolecular competition correspondingly), were run in 1,2-dichlorobenzene, DCB, using molar ratios similar to those described previously. DCB dissolves C₆₀ better than toluene.¹⁷ The presence of the heteroatom oxygen in these substrates $(2-d_1, 2-d_0, and 2-d_2)$ makes the separation and purification of the C_{60} derivatives much easier. Moreover, unlike toluene, DCB is completely removed from the reaction mixture at the end of the reaction. The ene adducts were characterized both by ¹H NMR chemical shifts and by FAB-MS. The isolated ene monoadduct was stable under the reaction conditions for several hours. The results are summarized in Table 1. The measured intramolecular (product) isotope effect $k_{\rm H}/k_{\rm D}$ was 1.5 and the intermolecular $k_{\rm H}/k_{\rm D}$ was 1.8, values which are similar to those obtained from the ene reaction of C_{60} with **1-d**₁ and **1-d**₀/**1-d**₂ ($k_{\rm H}/k_{\rm D} = 1.8$ and 2.0 respectively). In this case, the determination of the isotope effects was possible by integration of the appropriate signals in two independent hydrogen absorption regions, namely, by the integration of the vinylic vs the phenylic hydrogens, as well as of the methylenic vs C₆₀-H of the ene adducts. Both integrations gave, within experimental error, the same isotope effects. Thus, these results confirm the measured values of primary isotope effects of the $1-d_1$ and $1-d_0/1-d_2$ system with C₆₀. Furthermore, these results indicate that under the reaction conditions and purification processes no H/D exchange of C₆₀-H occurs, although the hydrogens on the C₆₀ carbon (C₆₀-H) are weakly acidic.¹⁸

The similar and substantial product and kinetic isotope effects exclude a mechanism where the formation of an intermediate is a rate-determining step, case A in Scheme 1.



Had this been the case, no C-H(D) bond breaking would have occurred at the transition state and only a very small intermolecular kinetic isotope effect, if any, would have been expected. The presense of substantial and equal kinetic and product isotope effects in the inter- and intramolecular competitions (Table 1) supports a concerted one-step reaction, i.e., case B in Scheme 1, with both C–H bond breaking and C–C bond making having occurred in the transition state. This is consonant with previous findings in a variety of other concerted ene reactions.¹² However, the reversible formation of a charge transfer or a dipolar intermediate preceding the rate-determining step, case C in Scheme 1, cannot be excluded.

The relatively small primary isotope effects, $k_{\rm H}/k_{\rm D}$ about 2, measured in these studies may be attributed either to a bent transition state, i.e., nonlinear transfer of the hydrogen between the two carbons, or to a linear but early transition state in which the C–H bond breaking is not extensively developed. It is interesting also to note that the reaction rate increases by a factor of 20 in going from the nonpolar solvent toluene to the more polar DCB.¹⁹ Furthermore, the ene

Scheme 2. Unsymmetrical and Partially Charged Transition State of the Ene Reaction of C_{60} and Arylallyl Alkenes.



⁽¹⁴⁾ Vriesema, B. K.; Lemaire, M.; Buter, J.; Kellogg, R. M. J. Org. Chem. **1986**, *51*, 5169.

⁽¹⁵⁾ Orfanopoulos, M. J. Chem. Res., Synop. 1982, 188.

⁽¹⁶⁾ Balh, A. L.; Lee, J. W.; Noll,; C. B.; Olmstead, M. M. J. Chem. Soc., Chem. Commun. **1993**, 56.

⁽¹⁷⁾ Ruoff, R. S.; Tse, D. S.; Malhotra, R.; Lorents, D. C. J. Phys. Chem. **1993**, *97*, 3379.

reaction of 3-(*p*-methoxyphenyl)-1-propene, $2-d_0$, is three times faster than that of the corresponding nonsubstituted allylbenzene, $1-d_0$. These results indicate an ene transition state with a partial dipolar character. A suprafacial orbital presentation of an unsymmetrical and partially charged transition state is shown below, Scheme 2.

In conclusion, the ene reaction of ary lallyl alkenes with C_{60} occurs either by a concerted mechanism or by the reversible formation of a charged or a dipolar intermediate that is followed by the rate-determining breaking of the C-H(D) bond.

Acknowledgment. We thank professor G. J. Karabatsos for valuable comments and discussions and Dr. A. Politou for NMR assistance. This work was supported by the Secretariat of Research and Technology (Grant Π ENE Δ -1995).

Supporting Information Available: GC MS and ¹H NMR spectra for 1-d₁, 1-d₂, 2-d₁, and 2-d₂; high-resolution FAB-MS and ¹H NMR spectra for adducts C_{60} with 1-d₁, 1-d₀/1-d₂, 2-d₁, and 2-d₀/2-d₂; kinetic data. This material is available free of charge via the Internet at http://pubs.acs.org.

OL990204O

⁽¹⁸⁾ Fagan, P. J.; Krusic, P. J.; Evans, D. H.; Lerke, S. A.; Johnston, E. *J. Am. Chem. Soc.* **1992**, *114*, 9697. (b) Nigazymbetov, M. E.; Evans, D. H.; Levke, S. A.; Cahill, P. A.; Henderson, C. C. *J Phys. Chem.* **1994**, *98*, 13093.

⁽¹⁹⁾ The dipole moments of toluene and 1,2-DCB are 0.36 and 2.5 D, respectively. *CRC Handbook of Chemistry and Physics*, 64th ed.; CRC Press: Boca Ratan, FL, 1983–1984.