

Palladium-Catalyzed Stereospecific Decarboxylative Benzylation of Alkynes

Shehani N. Mendis and Jon A. Tunge*

Department of Chemistry, The University of Kansas, 2010 Malott Hall, 1251 Wescoe Hall Drive, Lawrence, Kansas 66045, United States

Supporting Information



ABSTRACT: Enantioenriched benzyl esters of propiolic acids undergo highly stereospecific decarboxylative coupling to provide 1,1-diarylethynyl methanes. This $sp-sp^3$ coupling does not require strongly basic conditions or preformed organometallics and produces CO_2 as the sole byproduct. Ultimately, this method results in the successful transfer of stereochemical information from secondary benzyl alcohols to generate enantioenriched tertiary diarylmethanes.

ransition-metal-catalyzed cross-coupling reactions that generate tertiary stereogenic centers via the alkylation of secondary sp³-hybridized reactants are potentially powerful tools for asymmetric synthesis.¹ Despite the significant advancements made in this area, the asymmetric alkynylation of secondary benzyl electrophiles is scarcely reported.² Benzyl halides are known to undergo Sonogashira coupling as well as related couplings with other organometallic acetylides.^{2,3} However, these reactions rarely utilize secondary benzyl electrophiles and they suffer from the use of relatively toxic benzyl halides. Moreover, the couplings often require cocatalysts or preformed organometallics.^{2,3a-c} For example, the nickel-catalyzed enantioselective benzyl-acetylide coupling requires preformed trialkyl indium complexes.² In addition, benzyl-acetylide cross-couplings can result in the formation of unwanted side products via isomerization of the alkyne under basic media,^{3d} or further coupling of the product alkyne.^{3c}

Catalytic decarboxylative coupling reactions have emerged as potentially powerful alternatives to standard cross-coupling reactions.⁴ Thus, we envisioned that a wide array of tertiary diaryl methane motifs that are prevalent in biologically active compounds⁵ could be accessed via decarboxylative benzylacetylide couplings of enantioenriched diarylmethanol derivatives (Scheme 1). Decarboxylative coupling reactions are particularly well suited for the alkynylation of benzyl electrophiles since the reactions occur under formally neutral conditions.⁶ The absence of added base greatly decreases the rate of deprotonation which can lead to racemization of enantioenriched benzyl alkynes or formation of allene side products.^{3d} Decarboxylative cross-coupling reactions have the further advantage that they use benzyl alcohol derivatives in lieu of more toxic benzyl halides.² Lastly, secondary benzyl alcohols





are readily available in enantioenriched form,⁷ making them ideal substrates for asymmetric cross-coupling reactions.

Received: August 20, 2015

Legros and others have shown that secondary benzylic electrophiles derivatives can undergo palladium-catalyzed stereospecific cross-coupling reactions.⁸ These reactions proceed via the intermediacy of Pd- π -benzyl complexes that act as chiral benzyl cation equivalents (Scheme 1).

More recently, Jarvo and Watson have made impressive progress on the development of nickel-catalyzed stereospecific cross-coupling reactions of secondary benzylic electrophiles to generate molecules with tertiary stereocenters.⁹ Herein we report the synthesis of enantioenriched 1,1-diarylethynyl methanes via a highly stereospecific palladium-catalyzed decarboxylative benzylation strategy. To the best of our knowledge, this is the first report of a stereospecific decarboxylative benzylation. Moreover, we are unaware of any other cross-coupling method for the direct synthesis of enantioenriched 1,1-diarylethynyl methanes.

To begin, an array of benzylic ester derivatives were prepared via esterification of propiolic acids with enantioenriched diaryl methanols. These alcohols were obtained by a slightly modified (-)-MIB-catalyzed addition of boronic acids to aldehydes, originally reported by Braga (Scheme 2).^{7b} This method





produced a wide range of alcohols in high enantiomeric excess; however, several of the precursor alcohols were obtained in lower ee (e.g., **1d**, **1i**, **1o**, **1p**, **Table 1**). Nonetheless the Braga method provided each alcohol in large enough enantiomeric excess to accurately measure the stereospecificities of decarboxylative coupling. An additional advantage of this method for alcohol preparation is that either enantiomer of the alcohol can be obtained via the appropriate selection of the aryl boronic acid and the aromatic aldehyde reaction partners.

We then investigated the scope of the stereospecific decarboxylative benzylation reaction (Table 1). As we and others have noted,^{6b,10} benzylic cross-couplings that proceed via π -benzyl intermediates are much more facile for benzylic electrophiles that have extended π -conjugation. For example phenyl propiolic ester of diphenylmethanol did not show any reactivity under standard conditions. Therefore, a variety of substituted benzyl propiolates that contain 1-naphthyl, 2naphthyl, or indolyl groups were chosen for initial study. Since π -benzyl formation is accompanied by partial dearomatization of the coordinating arene, these aromatic groups that have relatively low resonance energies more readily form such complexes. For example, selective coordination of naphthalene instead of a phenyl group would be expected on the basis of the lower resonance energy of naphthalene (1b, Figure 1). Ultimately, the decarboxylative coupling of benzyl ester 1b is facile and forms the product 2b with a high degree of stereochemical fidelity, with a conservation of enantiomeric excess (cee) of 94% [cee = (product ee/reactant ee) \times 100] (Table 1). However, oxidative addition is sterically disfavored by an ortho substituent on the naphthyl ring (1c, Figure 1 andTable 1), preventing the reaction. Substitution at the ortho position of the noncoordinating arene (1d, Figure 1) does slow the coupling; however, the reaction still proceeds with high

 Table 1. Scope of Stereospecific Decarboxylative Benzylation

 of Alkynes

	Ar ¹ O Ar ² O	10 mol % Pd(PPh ₃) ₄ A^{r^1}										
	1 Pr		10 0,		2 F	Ph						
entry	reactant ee(%) ^a	product		yield (%) ^b	ee (%) ^a	cee (%)						
		(1)										
		R R										
1b	94	R=H) (2b)	85	88	94						
1c	-	R = Me	(2c)	NR	-	-						
		ОМе	<i>(</i> - -)									
10	69	1-Np	(2d)	31	64	93						
		1-Np										
1e	97	Pr "	(2e)	81	96	99						
		ÓMe 2-Np										
1f	94		(2 f)	71	91	97						
		Me	•									
		2-Np										
R Ph												
1g	94	R = F	(2g)	83	92	98						
1h 1i	>85 62	R = Me R = SMe	(2h) (2i)	77 68	95 43	>99 69						
1j	94	R = ^t Bu	(2 j)	92	89	95						
1k	91	2-Nn	(2k)	73	93	>99						
		QMe Pł	ו									
11	81	\bigcirc	(2I)	73	79	98						
		Ŷ										
		Pł	ו									
		H										
		1-Np										
1m	-	R = NO ₂	`R (2m)	16 ^c	ND	-						
1n	-	R = CN	(2 n)	18 ^c	ND	-						
		2-Np										
10	72	X → `Pł X = S	(2o)	69	64	89						
1р	75	X = 0	(2 p)	77	64	85						
10	87		(2 a)	62	60 ^d	60						
ιq	01		(2q)	92	00-	09						

^{*a*}Determined by chiral HPLC analysis. ^{*b*}Yield of isolated product All products were stored cold upon isolation to avoid decomposition or isomerization. ^{*c*}Yield reported using the racemic benzylic ester. ^{*d*}Boc group removed prior to HPLC analysis.

enantiospecificity (1d, Table 1). Further substitution of the benzylic esters with a variety of alkyl (2f, 2h, 2j), alkoxy (2d, 2e, 2l), or halogen (2g, 2k) substituents allowed for highly



Figure 1. π -Benzyl formation: sterics and favorable coordination of arenes with extended π -conjugation.

enantiospecific (average cee = 97%) cross-coupling, and the products were isolated in good to excellent yield. However, a sulfur donor in the para-position led to partial racemization during the coupling (2i, Table 1). Furthermore, substitution of one of the arenes with a strongly electron-withdrawing p-NO₂ or p-CN substituent, which substantially acidifies the benzylic proton, drastically lowered the yield of the reaction and these products were isolated as the allene isomers (2m, 2n). Interestingly, heteroaromatic phenyl propiolates (10-1q) also underwent stereospecific decarboxylative benzylation with moderate-to-good stereochemical fidelity.

We also briefly examined the effect of the acetylide coupling partner on the yields and stereospecificity of the decarboxylative coupling (Table 2). In addition to phenyl propiolates,



	Ar ¹ 0 Ar ² 0	to	0 mol % P Iuene, 110	d(PPh) °C, 1	1 <u>3)4</u> 19 h Ar	Ar ¹ 2 2	Ph
entry	reactant ee(%	%) ^a	product		yield (%) ^b	ee (%) ^a	cee (%)
1r	92	2-Np Ph	Me	(2 r)	43 ^c	87	95
1s	93	1-Np Ph	\sim	(2s)	53	81	87
1t	91	1-Np Ph		(2t)	59	87	96
1u	91	1-Np Ph	F	(2u)	62	85	93
1v	86	1-Np Ph		(2v)	87	77	90
1w	ND	1-Np Ph	° °OM€	• (2w)	NR		

^aDetermined by chiral HPLC analysis. ^bYield of isolated product. All products were stored cold upon isolation. ^cIsolated yield after 16 h.

alkyl (1r, 1s) and aryl propiolates (1t-1v) also undergo decarboxylative benzylation with high stereospecificity. However, a cyclohexenyl acetylide (1w) did not undergo decarboxylative benzylation under optimized reaction conditions.

While it was clear that the decarboxylative couplings of most propiolic esters occur with high stereospecificity, it was desirable to determine whether the coupling proceeded with retention or inversion of configuration. With this in mind, an alcohol [(R)-3k, Scheme 3] of known configuration was

Scheme 3. Determination of the Stereochemical Course of the Reaction



prepared.^{7c,11} The alcohol was converted to the highly enantiomerically enriched phenyl propiolate ester (R)-1k via PyBop coupling. After the decarboxylative coupling, crystals of the product were subjected to single crystal X-ray diffraction analysis which revealed the absolute configuration of 2k to be (S) (Scheme 3).

The overall inversion of stereochemistry that is observed in decarboxylative coupling can be mechanistically explained as follows (Scheme 4). Substrate 1 undergoes oxidative addition

Scheme 4. Proposed Catalytic Cycle for the Decarboxylative Coupling



to Pd to generate an η^3 -benzyl-Pd carboxylate intermediate. The oxidative addition is expected to proceed with inversion of configuration via S_N2-like displacement by palladium.¹²

We propose that coordination of the alkyne to Pd, as in intermediate A, facilitates decarboxylation resulting in the formation of a benzyl-Pd-acetylide intermediate (B) which undergoes reductive elimination with retention of stereochemistry to give the cross-coupled product 2 with overall inversion of stereochemistry. This is similar to Pd-catalyzed allylic alkylations, where allylation of nonstabilized or "hard" nucleophiles proceeds with inversion of configuration.^{6a}

Organic Letters

In summary, we have shown that decarboxylative benzylation of acetylides is highly stereospecific and that the stereochemical information on a secondary alcohol is successfully transferred to generate a tertiary stereogenic center. This method provides a route for the asymmetric coupling of secondary benzylic electrophiles with alkynes that does not require preformed organometallics.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.or-glett.5b02410.

Experimental procedures, ¹H, ¹³C NMR data, characterization data and HPLC data of all novel products (PDF) Crystallographic data for **2k** (CIF)

AUTHOR INFORMATION

Corresponding Author

*E-mail: tunge@ku.edu.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank the National Science Foundation (CHE-1465172) and the Kansas Bioscience Authority Rising Star program for financial support. We also thank Dr. Victor Day for X-ray crystallographic analysis and the NSF-MRI Grant CHE-0923449 used to purchase the X-ray diffractometer and software used in this study. Support for the NMR instrumentation was provided by NSF Academic Research Infrastructure Grant No. 9512331, NIH Shared Instrumentation Grant No. 0320648.

REFERENCES

(1) (a) Taylor, B. L. H.; Jarvo, E. R. Synlett 2011, 2011, 2761–2765.
(b) Swift, E. C.; Jarvo, E. R. Tetrahedron 2013, 69, 5799–5817.
(c) Liang, Y.; Fu, G. C. J. Am. Chem. Soc. 2014, 136, 5520–5524.
(d) Xu, S.; Lee, C.-T.; Wang, G.; Negishi, E.-i. Chem. - Asian J. 2013, 8, 1829–1835.

(2) Caeiro, J.; Pérez Sestelo, J.; Sarandeses, L. A. Chem. - Eur. J. 2008, 14, 741-746.

(3) (a) Pérez, I.; Sestelo, J. P.; Sarandeses, L. A. J. Am. Chem. Soc. 2001, 123, 4155–4160. (b) Qian, M.; Negishi, E. Tetrahedron Lett. 2005, 46, 2927–2930. (c) Pottier, L. R.; Peyrat, J.-F.; Alami, M.; Brion, J.-D. Synlett 2004, 1503–1508. (d) Larsen, C. H.; Anderson, K. W.; Tundel, R. E.; Buchwald, S. L. Synlett 2006, 2006, 2941–2946.

(4) (a) Weaver, J. D.; Recio, A.; Grenning, A. J.; Tunge, J. A. Chem. Rev. 2011, 111, 1846–1913. (b) Shang, R.; Huang, Z.; Xiao, X.; Lu, X.; Fu, Y.; Liu, L. Adv. Synth. Catal. 2012, 354, 2465–2472. (c) Fields, W. H.; Chruma, J. J. Org. Lett. 2010, 12, 316–319. (d) Recio, A., III; Heinzman, J. D.; Tunge, J. A. Chem. Commun. 2012, 48, 142–144. (e) O'Hair, R. A. J. Pure Appl. Chem. 2015, 87, 391–404. (f) Rodriguez, N.; Goossen, L. J. Chem. Soc. Rev. 2011, 40, 5030–5048.

(5) (a) Xie, W.-D.; Li, X.; Weng, C.-W.; Liu, S.-S.; Row, K. H. Chem. Pharm. Bull. 2011, 59, 511–514. (b) Ali, M. S.; Banskota, A. H.; Tezuka, Y.; Saiki, I.; Kadota, S. Biol. Pharm. Bull. 2001, 24, 525–528. (6) (a) Rayabarapu, D. K.; Tunge, J. A. J. Am. Chem. Soc. 2005, 127, 13510–13511. (b) Torregrosa, R. R. P.; Ariyarathna, Y.; Chattopadhyay, K.; Tunge, J. A. J. Am. Chem. Soc. 2010, 132, 9280– 9282. (c) Park, K.; Lee, S. RSC Adv. 2013, 3, 14165–14182. (7) (a) Bolm, C.; Rudolph, J. J. Am. Chem. Soc. 2002, 124, 14850–14851. (b) Braga, A. L.; Paixão, M. W.; Westermann, B.; Schneider, P. H.; Wessjohann, L. A. J. Org. Chem. 2008, 73, 2879–2882. (c) Salvi, L.; Kim, J. G.; Walsh, P. J. J. Am. Chem. Soc. 2009, 131, 12483–12493. (8) (a) Legros, J.-Y.; Boutros, A.; Fiaud, J.-C.; Toffano, M. J. Mol. Catal. A: Chem. 2003, 196, 21–25. (b) Legros, J.-Y.; Toffano, M.; Fiaud, J.-C. Tetrahedron 1995, 51, 3235–3246. (c) Assié, M.; Legros, J.-Y.; Fiaud, J.-C. Tetrahedron: Asymmetry 2005, 16, 1183–1187. (d) Legros, J.-Y.; Toffano, M.; Fiaud, J.-C. Tetrahedron: Asymmetry 2005, 16, 1183–1187. (d) Legros, J.-Y.; Toffano, M.; Fiaud, J.-C. Tetrahedron: Asymmetry 1995, 6, 1899–1902. (e) He, A.; Falck, J. R. J. Am. Chem. Soc. 2010, 132, 2524–2525. (f) López-Pérez, A.; Adrio, J.; Carretero, J. C. Org. Lett. 2009, 11, 5514–55.

(9) (a) Yonova, I. M.; Johnson, A. G.; Osborne, C. A.; Moore, C. E.; Morrissette, N. S.; Jarvo, E. R. Angew. Chem., Int. Ed. 2014, 53, 2422– 2427. (b) Wisniewska, H. M.; Swift, E. C.; Jarvo, E. R. J. Am. Chem. Soc. 2013, 135, 9083–9090. (c) Taylor, B. L. H.; Harris, M. R.; Jarvo, E. R. Angew. Chem., Int. Ed. 2012, 51, 7790–7793. (d) Zhou, Q.; Srinivas, H. D.; Dasgupta, S.; Watson, M. P. J. Am. Chem. Soc. 2013, 135, 3307–3310. (e) Maity, P.; Shacklady-McAtee, D. M.; Yap, G. P. A.; Sirianni, E. R.; Watson, M. P. J. Am. Chem. Soc. 2013, 135, 280– 285.

(10) (a) Greene, M. A.; Yonova, I. M.; Williams, F. J.; Jarvo, E. R. Org. Lett. **2012**, 14, 4293–4296. (b) Kuwano, R.; Kondo, Y.; Matsuyama, Y. J. Am. Chem. Soc. **2003**, 125, 12104–12105.

(11) Li, K.; Hu, N.; Luo, R.; Yuan, W.; Tang, W. J. Org. Chem. 2013, 78, 6350-6355.

(12) (a) Trost, B. M.; Czabaniuk, L. C. J. Am. Chem. Soc. 2012, 134, 5778–5781. (b) Lau, K. S. Y.; Wong, P. K.; Stille, J. K. J. Am. Chem. Soc. 1976, 98, 5832–5840.