

## Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Incn20>

### Chemical Incorporation of 1-Methyladenosine, Minor tRNA Component, into Oligonucleotides

E. V. Efimtseva<sup>a c</sup>, S. N. Mikhailov<sup>a</sup>, J. Rozenski<sup>b</sup>, R. Busson<sup>b</sup>, A. Van Aerschot<sup>b</sup> & P. Herdewijn<sup>b</sup>

<sup>a</sup> Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, Moscow, Russia

<sup>b</sup> Rega Institute Katholieke Universiteit Leuven, Leuven, Belgium

<sup>c</sup> Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, Vavilov str. 32, 119991, Moscow, Russia

Published online: 31 Aug 2006.

To cite this article: E. V. Efimtseva, S. N. Mikhailov, J. Rozenski, R. Busson, A. Van Aerschot & P. Herdewijn (2003) Chemical Incorporation of 1-Methyladenosine, Minor tRNA Component, into Oligonucleotides, *Nucleosides, Nucleotides and Nucleic Acids*, 22:5-8, 1113-1115, DOI: [10.1081/NCN-120022749](https://doi.org/10.1081/NCN-120022749)

To link to this article: <http://dx.doi.org/10.1081/NCN-120022749>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

## Chemical Incorporation of 1-Methyladenosine, Minor tRNA Component, into Oligonucleotides

E. V. Efimtseva,<sup>1,\*</sup> S. N. Mikhailov,<sup>1</sup> J. Rozenski,<sup>2</sup> R. Busson,<sup>2</sup>  
A. Van Aerschot,<sup>2</sup> and P. Herdewijn<sup>2</sup>

<sup>1</sup>Engelhardt Institute of Molecular Biology, Russian Academy of Sciences,  
Moscow, Russia

<sup>2</sup>Rega Institute Katholieke Universiteit Leuven,  
Leuven, Belgium

### ABSTRACT

The synthesis of suitably protected 1-methyladenosine derivatives has been developed and its successful chemical incorporation into oligonucleotides was achieved.

*Key Words:* 1-Methyladenosine; Minor tRNA component; Chemical synthesis; Oligonucleotides.


Modified nucleosides offer an overwhelming variety of chemistry and potential biological functions. More than hundred modified nucleosides have been discovered in nature, most of them occur in tRNAs. The contribution of these modifications to the functional chemistry, structure and biological activity of the RNAs largely

\*Correspondence: Ekaterina V. Efimtseva, Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, Vavilov str. 32, 119991 Moscow, Russia; Fax: +7 095 135 1405; E-mail: smikh@imb.ac.ru.



remains to be discovered. Approximately 25% of all tRNAs have 1-methyladenosine ( $m^1A$ ) at position 58 in the T loop.

Here we present our results on the preparation of  $m^1A$  building block and its successful incorporation into oligonucleotides (ONs). Previously oligoribonucleotides containing  $m^1A$  (**1**) were prepared using T4 RNA ligase<sup>[1]</sup>. To the best of our knowledge before our recent publication<sup>[2]</sup> no attempts have been made for chemical incorporation of  $m^1A$  into ONs.



a. MMTrCl/Py; b. (ClCH<sub>2</sub>CO)<sub>2</sub>O/Py; c. tBuMe<sub>2</sub>SiCl; d. iPr<sub>2</sub>NP(O)Cl(OCH<sub>2</sub>CH<sub>2</sub>CN).

Under alkaline conditions,  $m^1A$  (**1**) rearranges to N<sup>6</sup>-methyladenosine ( $m^6A$ ).<sup>[3]</sup>  $m^1A$  is difficult to handle due to its polar character. We have examined the stability of **1** under different conditions of deblocking which are used in ON synthesis. In 25% aq. ammonia the half-time of Dimroth rearrangement of **1** to  $m^6A$  was 36 h at 20°C. However, **1** was stable at least for 3 days in 2 M NH<sub>3</sub> in MeOH and for 2 days at 20°C in 1M Bu<sub>4</sub>NF in THF.

Tritylation of the hydroiodide salt of **1**<sup>[3]</sup> gave **2** in 68% yield. The benzoyl and acetyl groups for base protection of **2** were found to be too stable towards NH<sub>3</sub> in MeOH.<sup>[2]</sup> Therefore, more labile chloroacetyl group was chosen. Compound **3** was prepared in 80% yield using chloroacetic anhydride in pyridine with subsequent cleavage of *O*-chloroacetyl groups with NH<sub>3</sub> in MeOH (0°C, 30 min). After silylation 2'-isomer **4** was converted to the phosphoramidite **5** suitable for DNA and RNA synthesis. Deblocking of base-labile groups should be done in anhydrous conditions (2M NH<sub>3</sub> in MeOH 60 h, 20°C). Mass spectrometry together with enzymatic hydrolysis were used to prove ONs structure. The modification destabilizes a DNA duplex but slightly improves the stability of a hairpin when incorporated into the loop.

Authors thank KUL Research Council and RFBR for financial support.

## REFERENCES

- Helm, M.; Giege, R.; Florentz, C. A Watson-Crick base-pair-disrupting methyl group ( $m^1A$ ) is sufficient for cloverleaf folding of human mitochondrial tRNA<sup>Lys</sup>. *Biochemistry* **1999**, 38, 13,338–13,346.

2. Mikhailov, S.N.; Rozenski, J.; Efimtseva, E.V.; Busson, R.; Van Aerschot, A.; Herdewijn, P. Chemical incorporation 1-methyladenosine into oligonucleotides. *Nucleic Acids Res.* **2002**, *30*, 1124–1131.
3. Jones, J.W.; Robins, R.K. Purine nucleosides. III. Methylation studies of certain naturally occurring purine nucleosides. *J. Am. Chem. Soc.* **1963**, *85*, 193–201.



