This article was downloaded by: [University of Haifa Library] On: 27 September 2013, At: 11:09 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/lncn20</u>

SYNTHESIS OF 1-(2-NAPHTHYLSULFONYL)PYRAZOLE-C-GLYCOSIDES

Abd EI-Hamid Ismail^a, Abdel Aleem Hassan Abdel Aleem^a, Hamed Abdel Bary^a & Samy El-Assaly^a

^a Chemistry Department, Menoufia University, Shebin El-Koom, Egypt Published online: 31 Aug 2006.

To cite this article: Abd EI-Hamid Ismail, Abdel Aleem Hassan Abdel Aleem, Hamed Abdel Bary & Samy EI-Assaly (2002) SYNTHESIS OF 1-(2-NAPHTHYLSULFONYL)PYRAZOLE-C-GLYCOSIDES, Nucleosides, Nucleotides and Nucleic Acids, 21:6-7, 469-475

To link to this article: http://dx.doi.org/10.1081/NCN-120014819

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

©2002 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.

NUCLEOSIDES, NUCLEOTIDES & NUCLEIC ACIDS Vol. 21, Nos. 6 & 7, pp. 469–475, 2002

SYNTHESIS OF 1-(2-NAPHTHYLSULFONYL)PYRAZOLE-*C*-GLYCOSIDES

Abd El-Hamid Ismail,* Abdel Aleem Hassan Abdel Aleem, Hamed Abdel Bary, and Samy El-Assaly

Chemistry Department, Faculty of Science, Menoufia University, Shebin El-Koom, Egypt

ABSTRACT

2-Naphthylsulfonylhydrazine was reacted with aromatic aldehydes or aldehydo sugars to give the corresponding hydrazones which undergo Michael addition reactions with malononitrile or ethyl cyanoacetate to form pyrazole derivatives.

INTRODUCTION

Some pyrazoles are biologically active and have been used as analgesic anti-inflammatory,^[1] chemical control substance for pharmacological characterization of histamine receptors,^[2] antibacterial activity against Escherichia coli,^[3] sedative and hypontic,^[4] saluretic effect,^[5] a good activity as platelet aggregation inhibitor,^[6] a prostaglandin synthetase inhibitor^[7] and an antirheumatic drug.^[8]

469

DOI: 10.1081/NCN-120014819 Copyright © 2002 by Marcel Dekker, Inc. 1525-7770 (Print); 1532-2335 (Online) www.dekker.com

^{*}Fax: 20 48 235689

RESULTS AND DISCUSSION

The aim of the work in this paper is the synthesis of naphthylsulfonylpyrazoles and naphthylsulfonylpyrazole-C-glycosides. We expect that the naphthylsulfonyl group will increase the biological activity of the pyrazole ring. Thus, treatment of 2-naphthalene sulphonyl chloride with hydrazine hydrate produced 2-naphthyl-sulfonylhydrazine 1. Condensation of 2-naphthylsulfonylhydrazine 1 with equimolar amounts of aromatic aldehydes namely, benzaldehyde and anisaldehyde 2 in ethanol and in presence of glacial acetic acid as a catalyst yielded the corresponding hydrazones 3. 2-Naphthyl-sulfonylhydrazones 3 were allowed to react with malononitrile and/or ethyl cyanoacetate to afford 1-(2-naphthylsulfonyl)pyrazoles 5a-dafter purification by column chromatography. The reaction may be occur as follows: Initial addition of active methylene moiety to the double bond of hydrazones 3 affords an acyclic Michael adduct 4. The intermediate 4 cyclizes by internal addition to the cyano group or internal substitution at the ester carbonyl followed by aromatization form pyrazole derivatives 5a-d(Sch. 1).



Scheme 1.



Pyrazole-*C*-glycosides were prepared via 1,3-dipolar cycloaddition reaction of nitrilimines with acetylene derivatives^[9,10] and methylacrylate.^[11] In our work 2-naphthylsulfonylhydrazine **1** was reacted with D-glucose, D-mannose or D-xylose **6** to form the corresponding hydrazones **7**. The sugar-hydrazones **7** were reacted with malononitrile in ethanol and with a catalytic amount of triethylamine via Michael addition to afford 1-(2-naph-thylsulfonyl) pyrazole-*C*-glycosides **8** (Sch. 2).

EXPERIMENTAL

NMR spectra were recorded on a Bruker 250 FT NMR spectrometer, TMS as internal standard. The silica gel (0.040–0.63 mm) used for CC was purchased from Merck. Analytical TLC was performed on percolated TLC sheets (Merk silica gel 60 F_{254} 0.2 mm). Results of elemental analysis were in acceptable range.

General Procedure for Compound 5a-d

A mixture of 2-naphthylsulfonylhydrazones **3** (1 mmole) and malononitrile or ethyl cyanoacetate (1 mmole) in 10 mL of ethanol with a catalytic amount of triethylamine was boiled under reflux 6h. The solvent was evaporated under vacuum. The residue was purified by column chromatography with chloroform/methanol (90:10, v/v) to give **5a-d**. 5-Amino-4-cyano-3-phenyl-1-(2-naphthylsulfonyl)pyrazole (5a)

Yield (54%); m.p. 182°C. ¹H-NMR (DMSO- d_{δ}): δ 11.64 (s, 1H, NH), 8.53-7.39 (m, 9H, ArH); Anal. Calcd for C₂₀H₁₄N₄O₂S (374.42): C, 64.15; H, 3.27; N, 14.96. Found: C, 64.43; H, 3.76; N, 14.69.

5-Amino-4-cyano-3-(4-methyoxyphenyl)-1-(2-naphthylsulfonyl)pyrazole (5b)

Yield (56%); m.p. 188°C. ¹H-NMR (DMSO- d_{δ}): δ 11.50 (s, 1H, NH), 9.23-8.65 (m, 10H, ArH), 3.59 (s, 3H, CH₃O); Anal. Calcd for C₂₁H₁₆N₄O₃S (404.45): C, 62.34; H, 3.99; N, 13.85. Found: C, 62.66; H, 4.20; N, 13.69.

4-Cyano-3-phenyl-1-(2-naphthylsulfonyl)pyrazole-5-one (5c)

Yield (45%); m.p. 191°C. ¹H NMR (DMSO- d_{δ}): δ 11.53 (s, 1H, NH), 8.77-7.51 (m, 8H, ArH); Anal. Calcd for C₂₀H₁₃N₃O₃S (375.4): C, 63.99; H, 3.49; N, 11.19. Found: C, 63.72; H, 3.40; N, 11.45.

4-Cyano-3-(4-methoxyphenyl)-1-(2-naphthylsulfonyl)pyrazole-5-one (5d)

Yield (54%); m.p. 182°C. ¹H NMR (DMSO- d_{δ}): δ 11.4 (s, 1H, NH), 8.50-7.86 (m, 11H, ArH), 3.11 (s, 3H, OCH₃); Anal. Calcd for C₂₁H₁₅N₃O₄S (404.43): C, 62.21; H, 3.72; N, 10.35. Found: C, 62.53; H, 3.60; N, 10.72.

General Procedure for Hydrazones 3a, b and 7a-c

A mixture of 2-naphthylsulfonylhydrazine (1 mmole) and D-Glucose, D-Mannose, D-Xylose, benzaldehyde or anisaldehyde in 10 mL of ethanol with a few drops of acetic acid was boiled under reflux for 1 h. After concentration and cooling, the separated precipitate was filtered off and crystallized from the appropriate solvents (**3a**, **b** from ethanol, **7a**-**c** from 1,4-dioxane).

Benzaldehyde (2-naphthylsulfonyl)hydrazone (3a)

Yield (81%); m.p. 188°C; Anal. Calcd for $C_{17}H_{14}N_2O_2S$ (310.40): C, 65.79; H, 4.55; N, 9.03. Found: C, 75.61; H, 4.80; N, 9.11.

1-(2-NAPHTHYLSULFONYL)PYRAZOLE-C-GLYCOSIDES

Anisaldehyde (2-naphthylsulfonyl)hydrazone (3b)

Yield (70%); m.p. 192°C; Anal. Calcd for $C_{18}H_{16}N_2O_3S$ (340.36): C, 63.52; H, 4.74; N, 8.23. Found: C, 63.29; H, 4.81; N, 8.00.

D-Glucose 2-naphthylsulfonylhydrazone (7a)

Yield (76%); m.p. 173°C; Anal. Calcd for $C_{16}H_{20}N_2O_7S$ (384.4): C, 49.99; H, 5.24; N, 7.28. Found: C, 50.5; H, 5.40; N, 7.52.

D-Mannose 2-naphthylsulfonylhydrazone (7b)

Yield (72%); m.p. 179°C; Anal. Calcd for $C_{16}N_{20}N_2O_7S$ (384.4): C, 49.99; H, 5.25; N, 7.28. Found: C, 49.52; H, 5.00; N, 7.43.

D-Xylose 2-naphthylsulfonylhydrazone (7c)

Yield (73%); m.p. 168°C; Anal. Calcd for $C_{15}H_{18}N_2O_6S$ (354.37): C, 50.84; H, 5.12; N, 7.90. Found: C, 50.73; H, 5.30; N, 7.63.

General Procedure for Compound 8a-c

A mixture of 2-naphthylsulfonylhydrazones 7 (1 mmole) and malononitrile (1 mmole) in 10 mL absolute ethanol and 1 mL triethylamine was boiled under reflux for 6 h. The solvent was evaporated under vacuum. The residue was purified by column chromatography with chloroform/methanol (90:10, v/v) to produce compounds 8a-c.

5-Amino-4-cyano-3-(D-gluco-1,2,3,4,5,pentahydroxypentyl)-1-(2-naphthyl sulfonyl)pyrazole (**8a**)

Yield (53%); m.p. 180°C; ¹H-NMR (DMSO-d_{δ}): δ 9.77 (s, 1H, NH), 8.55-7.60 (m, 7H, ArH), 5.67 (s, 1H, OH-1'), 4.61 (s, 1H, H-1'), 4.35 (m, 3H, OH-2', OH-3' and OH-4'), 4.01 (m, 1H, OH-5'), 3.91 (m, 3H, H-2', H-3' and H-4'), 2.58 (m, 2 H, <u>CH₂OH</u>). ¹³C NMR (DMSO-d_{δ}): δ 136.58, 134.26, 131.65, 129.20, 128.81, 128.62, 128.44, 127.70, 127.36, 122.90 (C arom.), 112.11(C=N), 89.96 (C-1'), 77.96 (C-2'), 76.80 (C-3'), 70.61 (C-4'), 61.77 C₅'); Anal. Calcd for C₁₉H₂₀N₄O₇S (448.45): C, 50.88; H, 4.5; N, 12.49. Found: C, 50.71; H, 4.43; N, 12.62.

5-Amino-4-cyano-3-(D-manno-1-1,2,3,4,5,pentahydroxypentyl)-(2-naphthyl sulfonyl)pyrazole (**8b**)

Yield (55%); m.p. 178°C; ¹H-NMR (DMSO-d_δ): δ 9.02 (s, 1H, NH); 8.52-7.58 (m, 7H, ArH) 4.70 (m, 2H, OH-1' and H-1') 4.58 (m, 3H, OH-2', OH-3', OH-4'), 3.98 (m, 1H, OH-5'), 3.61 (m, 3H, H-2', H-3', H-4'); 2.89 (m, 1H, C<u>H</u>₂OH). ¹³C NMR (DMSO-d_δ): δ 136.51, 134.33, 131.70, 129.22, 128.85, 128.41, 127.75, 127.32, 124.94, 122.80 (C arom.), 112.33 (C≡N), 91.71 (C-1'), 76.80 (C-2'), 70.28 (C-3'), 69.40 (C-4'), 66.75 (C-5'); Anal. Calcd for C₁₉H₂₀N₄O₇S (448.48): C, 50.88; H, 4.50; N, 12.49. Found: C, 50.90; H, 5.00; N, 12.55.

5-Amino-4-cyano-3-(D-xylo-1,2,3,4,tetrahydroxybutyl)-1-(2-naphthyl sulfonyl)pyrazole (**8c**)

Yield (53%); m.p.; 188°C; ¹H-NMR (DMSO-d_δ): δ 11.39 (s, 1H, NH), 8.51-7.86 (m, 7H, ArH), 5.40 (s, 1H, OH-1'), 4.91 (s, 1H, H-1'), 4.45 (m, 1H, OH-2'), 4.12 (m, 1H, OH-3'), 3.94 (m, 1H, OH-4'), 3.60 (m, 2H, H-2' and H-3'), 2.94 (m, 2H, CH₂OH); ¹³C NMR (DMSO-d_δ): δ 136.22, 134.39, 131.76, 129.29, 128.90, 127.84, 127.35, 123.07 (C arom.), 112.00 (C \equiv N), 91.20 (C-1') 76.63 (C-2'), 70.18 (C-3'), 69.70 (C-4'); Anal. Calcd for C₁₈H₁₈N₄O₆S (418.43): C, 51.66; H, 4.34; N, 13.38. Found: C, 15.50; H, 4.46; N, 13.22.

ACKNOWLEDGMENT

The authors would like to express their thanks to DANIDA establishment, Denmark, which supports the project "Development of new drugs against Hepatitis" at the Chemistry Department, Faculty of Science, Menoufia University for the laboratory facilities and spectroscopic measurements.

REFERENCES

- 1. Credneer, K.; Tauscher, M.; Jozic, L.; Brenner, G. Salts of 5-methylpyrazole-3-Carboxylic Acid with Basically Substituted Adenine Derivatives, Preparation and Lipolysis Inhibitory Activity. Arzneim-Forsch. **1981**, *31*, 2096.
- Bauer, V.L.; Dalalian, H.P.; Fanshawe, W.J.; Safir, R.S.; Tocus, E.C.; Boshart, C.R. 4-[3(5)-Pyrazolyl]pyridinium salts. A New Class of Hypoglycemic Agents. J. Med. Chem. 1968, 11, 981.
- 3. Kornet, M.J.; Garret, R.J. Synthesis of 1-phenyl-2(phenylcarbamoyl)pyrazolidines as Potential Anticonvulsant Agents. J. Pharm. Sci. **1979**, *68*, 377.
- 4. Berger, J.C.; Iorio, L.C. Antianxiety Agents, Anticonvulsants, and Sedativehypnotics. Annu. Rep. Med. Chem. **1980**, *15*, 26.

1-(2-NAPHTHYLSULFONYL)PYRAZOLE-C-GLYCOSIDES

- 5. Kornet, M.J. New Local Anesthetics. 1 Esters of 1,2-diethyl-(3-hydroxymethyl) Pyrazolidine. J. Med. Chem. **1966**, *9*, 493.
- 6. Zeeb, B.; Goetz, N.; Ammermann, E.; Pommer, E.H. Ger. Pat. 1981, 2926, 280.
- 7. Cassidy, P.E. Thermally Stable Polymer; Dekker: New York, 1980.
- 8. Buchanan, J.G.; Stobie, A.; Wightmann. C-Nucleosides Studies. Part 14¹. A New Synthesis of Pyrazofurin. J. Chem. Soc. Perkin Trans. **1981**, *1*, 2374.
- Tronchet, M.J.; Mlle, F.P. Helvetica Chemica Acta 1971, 54, 683. C-Glycosides IV¹). Mise en évidence d' un mécanisme concurrent de la cyclo-addition dipolar-1,3cocertée los de la réaction d'une azométhine-imine avce un dipolaraphile acéylenique.
- 10. Tronchet, J.M.; Mlle, F.P. Helvetica Chemica Acta **1970**, *53*, 70. C-Glycosides 1); Synthese d'un analogue de la pyrazomycine.
- 11. Tronchet, J.M.; Tronchet, J.F.; Barbalat Rey, F. Synthesis of Pyrazole-Cglycosides by 1,3-Dipolor Cycloaddition of Nitrilimines Formed by Lead Tetraacetate oxidation of p-Nitrophenylhydrazones of Aldehydo Sugars. Heterocycles **1993**, *36* (4), 833.

Received February 7, 2002 Accepted June 12, 2002