STRUCTURE OF 2:1 COBALT(III) COMPLEXES DERIVED FROM ARYLAZOCITRAZINIC ACID

Jaroslava Kořínková^{*a*1,*}, Antonín Lyčka^{*b*}, Aleš CEE^{*c*}, Milan Nádvorník^{*a*2}, Robert JIRÁSKO^{*a*3} and Radim HRDINA^{*a*4}

 ^a Faculty of Chemical Technology, University of Pardubice, Studentská 95, CZ-532 10 Pardubice, Czech Republic; e-mail: ¹ jaroslava.korinkova@upce.cz, ² milan.nadvornik@upce.cz,
³ robert.jirasko@upce.cz, ⁴ radim.hrdina@upce.cz

^b Research Institute for Organic Syntheses, Rybitví 296, CZ-533 54 Pardubice 20, Czech Republic, and Faculty of Education, University of Hradec Králové, Rokitanského 62, CZ-500 03 Hradec Králové, Czech Republic; e-mail: antonin.lycka@vuos.com

^c Foerstrova 1688/5, CZ-500 02 Hradec Králové, Czech Republic; e-mail: ales.cee@home-net.cz

Received October 1, 2008 Accepted January 22, 2009 Published online March 18, 2009

¹H, ¹³C NMR and MS spectra of the 2:1 cobalt(III) complexes, the azo coupling products of 2-aminophenol or 2-aminobenzoic acid with the citrazinic acid, were measured and analysed. It was found that the cobalt atom is six-coordinated being bound to two oxygens (one from primary and one from secondary component) and the nitrogen of 2-aminophenol or 2-aminobenzoic acid. Molar magnetic susceptibility data proved the presence of Co(III). Diazotisation of 2-aminophenol or 2-aminobenzoic acid and subsequent coupling reaction with citrazinic acid and complexation of the formed azo product proceeded on standard way.

Keywords: Citrazinic acid; Complexation; Co(III) complexes; Chelates; Cobalt; NMR spectroscopy; Mass spectrometry.

Metal-complex dyes play a very important role in dyestuff technology^{1–5}. The complexation converts relatively cheap dyes to attractive commercial products having appropriate stability and shade. The dyes used in preparation of metal-complex dyes are usually tridentate ligands and the resulting products can have different configurations (*fac, mer*). It would be ideal to determine the real structure of these dyes using X-ray data. The number of existing X-ray structures is very limited (e.g. refs.^{6,7}) because of extreme difficulties in growing single crystals.

X-ray crystallography and NMR spectroscopy are in same sense complementary methods for the structure determination. NMR spectra provide information about the uniformity of metal-complex dyes, dynamic behaviour, position of coordination, etc., allowing, at least partially, to solve the structures of metal-complex dyes in cases where X-ray data are not available. We have successfully studied several types of metal-complex dyes derived from ligands having either hydroxy or carboxylic groups by multi-nuclear magnetic resonance⁸⁻¹⁴.

An important group of metal-complex dyes is derived from citrazinic acid as secondary component¹⁵. Masoud et al.¹⁶ proposed the existence of structure **1** having four-membered rings as a consequence of coordination of citrazinic acid nitrogen and both oxygen atoms.



This structure differs considerably from those shown previously and also from other structures mentioned in ref.¹⁶. Since citrazinic acid and its derivatives are promising secondary components, we decided to study this type of compounds using detailed ¹H, ¹³C NMR and MS spectra.

RESULTS AND DISCUSSION

Synthesis of the Dyes

The preparation of azo dyes 2a-2c is based on the diazotisation of primary amine (2-aminophenol, 2-aminobenzoic acid or 2-methylaniline) in dilute aqueous HCl at temperature 0–5 °C, where the reaction time is about 20–30 min (Scheme 1).



Scheme 1

The obtained diazonium salt is coupled with aqueous solution of a mixture of citrazinic acid and its salt at pH ~ 8, at temperature 0–5 °C; reaction time ca. 3 h (Scheme 2).



 $X = OH (2a), COOH (2b), CH_3 (2c)$

SCHEME 2

Cobalt(III) complex dyes **3a**, **3b** are prepared in water by complexation reaction of azo dyes **2a**, **2b** with cobalt(II) sulfate at pH ~ 8–9 and reaction temperature 100 °C (reflux); reaction time ca. 3 h (Schemes 3 and 4).



SCHEME 3

The reaction mixture is vigorously stirred to facilitate the oxidation of cobalt ($Co^{2+} \rightarrow Co^{3+}$) with air oxygen.

Thus, the compound Na₃**3a** is a Co(III) complex with 3-[(2-hydroxyphenyl)azo]citrazinic acid, Na₃**3b** is a Co(III) complex with 3-[(2-carboxyphenyl)azo]citrazinic acid. The reference anion **4** is a Co(III) complex with 2-[(2-hydroxyphenyl)hydrazono]-3-oxo-*N*-phenylbutanamide⁸ and the reference anion **5** is a Co(III) complex with 2-[(2-hydroxy-1-naphtalenyl)azo]benzoic acid¹⁷.







SCHEME 4





Collect. Czech. Chem. Commun. 2009, Vol. 74, No. 4, pp. 535-544

Constitution of Prepared Dyes

The ¹H and ¹³C NMR chemical shifts of starting azo dyes **2a** and **2b** and the anion **3a** were measured in DMSO- d_6 and the obtained data are collected in Table I.

Atom	2a		2b		3a		4 (ref. ⁸)	
	δ(¹ H)	δ(¹³ C)	δ(¹ H)	δ(¹³ C)	δ(¹ H)	δ(¹³ C)	δ(¹ H)	δ(¹³ C)
2	_	а	_	b	_	с		
3	-	121.8	_	123.3	_	114.7		
4	-	146.7	_	152.1	-	d		
5	6.25	114.1	5.97	112.1	5.83	104.2		
6	-	а	-	b	-	с		
7	-	166.4	-	167.5	-	167.4		
NH-N	14.38	-	15.66	-				
1′	-	129.4	-	143.5	-	145.8	_	147.1
2′	-	146.3	-	120.1	-	168.8	_	167.7
3′	7.00	116.2	8.01	131.5	6.70	117.3	6.63	117.5
4'	7.04	126.6	7.20	123.9	7.00	128.9	6.92	127.9
5′	6.94	120.3	7.56	132.8	6.70	114.4	6.67	114.2
6'	7.47	114.7	7.86	115.3	8.03	114.6	8.23	114.9
7′	10.82	-	d	168.7	-	-	-	-

TABLE I ¹H and ¹³C NMR chemical shifts of anions 2-4 in DMSO- d_6

^a 163.9 or 161.1; ^b 164.9 or 160.7; ^c 148.7 or 148.3; ^d not observed. Additional signals: **2a**: $\delta(^{1}\text{H}) = 11.79$ (br s, 1 H, acidic proton of NH or OH group); **3a**: $\delta(^{1}\text{H}) = 11.74$ (br s, 1 H, acidic proton of NH or OH group).

The ¹H and ¹³C NMR chemical shifts were assigned using H,H-COSY, gradient-selected gs-HMQC and gs-HMBC ^{11,12} (optimised for ¹J(¹³C,H) ca. 150 Hz and ³J(¹³C,H) ca. 8 Hz). Correlation of H(3') with the carbon of C-OH or COOH group in HMBC spectra was a key information for the assignment of hydrogen and carbon resonance of C1'-C7'.

Only one set of appropriate NMR signals was obtained for anion **3a**. This means that both ligands in the complex **3a** are equivalent and there is only one arrangement or a very fast exchange on the NMR time scale. The second

possibility can be theoretically examined from the measurement at low temperature. Since melting point of DMSO is +18 °C and since our complexes are insoluble in other solvents, this low temperature technique cannot be used. In addition, both ligands are spatially so far that the use of NOE or NOESY is impossible.

We compared the ¹H and ¹³C chemical shifts measured for anion **3a** with those for anion **4** for which we undoubtedly proved⁸ that cobalt atom is six-coordinated via two oxygen and nitrogen atoms from starting 2-aminophenol. A comparison of ¹H and ¹³C chemical shifts in complexes **3a** and **4** convincingly shows that the NMR data for comparable structures in both molecules are practically the same. Consequently, the bonding arrangement around cobalt atom in both complexes is also quite the same.

To provide chemical evidence for the importance of the hydroxy group presence in the ligand molecule, we prepared compound 2c in which hydroxy group was replaced with methyl group having similar steric requirement as OH group. In accord with our expectation, no complexation of 2c with cobalt(III) cation was observed.

Moreover, in structure **1**, proposed by Masoud et al.¹⁶, it could be expected that both the hydroxy groups and even the phenylazo substituent are not of decisive importance for a formation of cobalt(III) complex. Thus, we tested the complexation of unsubstituted citrazinic acid but, again, no complexation was observed.

Schetty and Kuster¹⁷ observed the three components having considerably different shades in cobalt(III) complex **5**. We proved¹³ that all the anions **3a**, **3b**, **4** and **5** contain six-coordinated cobalt atom and the difference consists in different mutual orientation of two ligands. All three components, however, are separable, after standing of each isomer in DMSO solution, which results in the formation of the starting equilibrium.

We measured ¹H and ¹³C NMR spectra of anion **3b** and observed a mixture of four components. The ¹H NMR spectrum consists of a strongly overlapped signals (δ (¹H) 7.25–8.28) and four separated signals resonating at 6.34, 6.20, 6.14 and 6.07 ppm, having approximately the same intensities. The latter ¹H NMR resonances resemble those for H-3' protons¹³ in the compound. An attempt to separate these isomers chromatographically was not completely successful since the tendency to establish an equilibrium during evaporation of solvent is stronger in **3b** than that in **5**.

The fact that the ligand **2b** was bound to cobalt(III) ion (forming **3b**) was also convincingly proved using mass spectrometry. The value of m/z of **3b** was found to be 661 (negative mode of electrospray ionisation) and 663

(positive mode of electrospray ionisation) for each chromatographic fraction.

We studied the cobalt oxidation state by measuring specific susceptibility of compound Na₃**3a**. The experimental value χ_g was (0.85 ± 0.02) × 10⁻⁸ m³ kg⁻¹. Calculating molar magnetic susceptibility $\chi_M = 570 \times 10^{-11}$ m³ mol⁻¹ and correcting this value for diamagnetic contribution 358 × 10⁻¹¹ m³ mol⁻¹, we obtained corrected molar magnetic susceptibility $\chi_M = 928 \times 10^{-11}$ m³ mol⁻¹.

Anion **3a** exists predominantly as a diamagnetic complex with Co(III) in the low-spin state (t_{2g}^{6}) (high-spin state has been observed so far exclusively in hexafluorocobaltate(III) complexes in which their magnetic properties are determined by four unpaired electrons with nonquenched orbital angular momentum and observed magnetic moments 6.5–9.5 BM ¹⁹). A small contribution of a Co(II) impurity in the high-spin state with the configuration (t_{2g}^{5})(e_{g}^{2}) having three unpaired electrons cannot be excluded taking experimental data into account. Complexes of this type give magnetic moments μ_{eff} in the range from 4.3 to 5.2 BM ¹⁹ which correspond to the values of corrected molar magnetic susceptibility in the range of 9760 × 10⁻¹¹–14268 × 10⁻¹¹ m³ mol⁻¹ using the expression $\mu_{eff} = 797.5 \times (\chi_M T)^{1/2}$. Provided this assumption is correct, the Co(II) complex content could be estimated to be 6–9% using the experimental value of $\chi_M = 876 \times 10^{-11}$ m³ mol⁻¹.

The results of electrospray ionisation (ESI) mass spectrometry measurements both in the positive- and negative-ion modes for compounds 2a-2cand Na⁺ salts **3a**, **3b** are summarised in Experimental and they are in agreement with their molecular weights and the structures. Ions presenting in the full-scan positive- and negative-ion mass spectra strongly depend on the Na⁺ concentration in the solution which is related to the character of measured compounds. The presence of carboxylic group of ligands is confirmed by the fragment ions observed in the full-scan negative-ion mass spectra for compounds **2a**, **2b**, **2c** and Na₃**3b** and tandem mass spectra for compound Na₃**3a**.

CONCLUSIONS

¹H, ¹³C NMR and MS data, suitable model experiments and taking into account some data from literature, we proved that 2:1 cobalt(III) complexes **3a** and **3b** derived from azo coupling products of 2-aminophenol or 2-aminobenzoic acid with citrazinic acid have the same structure as the complexes prepared by coupling of 2-aminophenol or 2-aminobenzoic acid with other secondary components and subsequent complexation. Anion **3a**

is a less flexible complex with new five- and six-membered rings, which are more flexible and, thus, several isomers existing in an equilibrium can be formed. Molar magnetic susceptibility data proved the existence the Co(III) oxidation state in the complexes.

EXPERIMENTAL

General

Acetonitrile and methanol for HPLC were purchased from Sigma–Aldrich (Prague, Czech Republic). Water was purified using the Milli-Q® plus water purification system (Millipore, USA). Ammonium acetate was purchased from Lach-Ner (Czech Republic). ¹H and ¹³C NMR spectra were recorded on a Bruker Avance 500 spectrometer (500.13 MHz for ¹H and 125.76 MHz for ¹³C) in DMSO- d_6 . The ¹H and ¹³C NMR chemical shifts were referenced to internal TMS. All 2D experiments (gradient-selected gs-COSY, gs-HMQC, gs-HMBC) were performed using manufacturer's software. Magnetic susceptibility was measured at room temperature (25 °C) using a Variable Temperature Gouy Balance apparatus (Newport Instruments, UK) and Hg[Co(NCS)₄] as calibrant. The values of magnetic molar susceptibility were corrected by the Pascal constants¹⁹. Positive- and negative-ion electrospray ionisation (ESI) mass spectra were measured on an ion trap analyser (Esquire 3000, Bruker Daltonics, Germany). The full-scan mass spectra were measured in the range m/z 50–1000. The samples were dissolved in a water/acetonitrile (1:1) mixture and analysed by direct infusion at the flow rate 5 μ l min⁻¹ and 10 Pa, respectively.

Diazotisation of 2-Aminophenol

2-Aminophenol (0.02 mol, 2.18 g; Aldrich) was mixed with water (20 ml) and concentrated HCl (5 ml, 35%). The obtained fine suspension was externally cooled under intensive stirring to 0–5 °C and 5 M aqueous NaNO₂ (5 ml) was slowly added. The cooling was stopped and the suspension was stirred at ambient temperature (ca. 22 °C) for 20–30 min until a yellow-brown solution of the diazonium compound was produced. The solution was used for subsequent coupling reaction.

Preparation of 3-[(2-Hydroxyphenyl)azo]citrazinic Acid (2a)

Citrazinic acid (0.02 mol, 3.20 g; Aldrich) was mixed with water (15 ml) and 5 M aqueous NaOH (13 ml) and the temperature was maintained at 0–5 °C (external cooling). The solution of the diazonium compound was gradually added under intensive stirring. pH was adjusted to ca. 8 by the addition of 5 M aqueous NaOH. When the coupling reaction was completed (after ca. 3 h), pH was adjusted to 4 by the addition of concentrated HCl (35%) and the precipitated monoazo dye was filtered off and dried at 90–110 °C. The yield of crude dye **2a** was 5.84 g (98%). The purity of the dye was determined by TLC (Alugram SIL G/UV₂₅₄, Macherey-Nagel, Germany; mobile phase: propan-2-ol/propan-1-ol/ethyl acetate/ water 2:4:1:3 (v/v/v/v)). According to TLC, **2a** was pure (R_F 0.8). C₁₂H₉N₃O₅ (275): ESI⁺, *m/z*: 320.0 [M + 2 Na – H]⁺, 100%; 301.9 [M + 2 Na – H – H₂O]⁺; 276.0 [M + H]⁺. ESI⁻, *m/z*: 295.8 [M – 2 H + Na]⁻; 273.8 [M – H]⁻; 255.8 [M – H – H₂O]⁻; 229.8 [M – H – CO₂]⁻, 100%.

542

Preparation of 3-[(2-Carboxyphenyl)azo]citrazinic Acid (2b)

Preparation was analogous to the preparation of compound **2a**. According to TLC, **2b** was pure ($R_{\rm F}$ 0.75). C₁₃H₉N₃O₆ (303): ESI⁻, *m/z*: 301.8 [M – H]⁻, 100%; 257.8 [M – H – CO₂]⁻; 213.8 [M – H – 2 CO₂]⁻.

Preparation of Complex Na₃3a

The monoazo dye **2a** (0.02 mol, 5.94 g) was dissolved in water (50 ml) and pH was adjusted to 8–9 by the addition of 5 M aqueous NaOH. Then $CoSO_4 \cdot 7H_2O$ (2.80 g, 0.01 mol) was added and the reaction mixture was vigorously stirred at 100 °C (reflux) for ca. 3 h. When the reaction was completed, compound Na₃**3a** was precipitated by addition of NaCl, filtered off and dried. The yield of compound Na₃**3a** was ca. 78% (based on **2a**). The purity of the dye Na₃**3a** was determined by TLC (Alugram SIL G/UV₂₅₄, Macherey–Nagel, Germany; mobile phase: propan-2-ol/propan-1-ol/ethyl acetate/water 2:4:1:3 (v/v/v/v)). According to TLC, **3a** was pure (R_F 0.69). Na₃[Co(C₁₂H₆N₃O₅)₂]; C₂₄H₁₂CON₆Na₃O₁₀ (672): ESI⁺, *m/z*: 651.0 [M – Na + 2 H]⁺, 100%. ESI⁻, *m/z*: 605.1 [M – 3 Na + 2 H]⁻, 100%. MS/MS, *m/z*: 605; 560.9 [M – 3 Na + 2 H – CO₂]⁻; 517.0 [M – 3 Na + 2 H – 2 CO₂]⁻.

Preparation of Complex Na₃3b

Preparation was analogous to the preparation of complex Na₃**3a**. According to TLC, **3b** was pure $(R_{\rm F} 0.71)$. Na₃ $[Co(C_{13}H_6N_3O_6)_2]$; $C_{26}H_{12}CoN_6Na_3O_{12}$ (728): ESI⁺, *m/z*: 684.9 [M - 2 Na + 3 H]⁺; 663.0 [M - 3 Na + 4 H]⁺, 100%. ESI⁻, *m/z*: 661.0 [M - 3 Na + 2 H]⁻, 100%; 616.9 [M - 3 Na + 2 H - CO₂]⁻; 572.9 [M - 3 Na + 2 H - 2 CO₂]⁻.

Preparation of 3-[(2-Methylphenyl)azo]citrazinic Acid (2c)

Preparation was analogous to the preparation of compound **2a**. According to TLC, **2c** was pure ($R_{\rm F}$ 0.84). C₁₃H₁₁N₃O₄ (273): ESI⁻, *m/z*: 271.8 [M – H]⁻; 227.8 [M – H – CO₂]⁻, 100%.

The authors thank the Ministry of Education, Youth and Sports of the Czech Republic (Project MSM 0021627501) for financial support. R. Jirásko and J. Kořínková acknowledge the support of the the Ministry of Education, Youth and Sports of the Czech Republic (Project MSM 0021627502).

REFERENCES

- 1. Price R. in: *The Chemistry of Synthetic Dyes* (K. Venkataraman, Ed.), Vol. III. Academic Press, New York and London 1970.
- 2. Baumann H.: Ullmanns Encyclopädie der Technischen Chemie, Vol. 16. Verlag Chemie, Weinheim 1978.
- 3. Beffa F., Back G.: Rev. Prog. Color Relat. Top. 1984, 14, 33.
- 4. Zollinger H.: Color Chemistry. Syntheses, Properties and Application of Organic Dyes and Pigments. Verlag Chemie, Weinheim 1987.
- 5. Schündehüte K.-H. in: *Methoden der Organischen Chemie* (Houben–Weil) (R. Stroh, Ed), Vol. 10/3. Georg Thieme, Stuttgart 1965.
- 6. Grieb R., Nigli A.: Helv. Chim. Acta 1965, 48, 317.

- 7. Lehmann U., Rihs U. in: *Chemistry of Functional Dyes* (Z. Yoshida and T. Kitao, Eds), p. 215. Mita Press, Tokyo 1989.
- 8. Lyčka A., Jirman J., Cee A.: Magn. Reson. Chem. 1990, 28, 408.
- 9. Abilgaard J., Hansen P. E., Josephsen J., Lyčka A.: Inorg. Chem. 1994, 33, 5271.
- 10. Tušek-Božič L., Čurič M., Vikič-Topič D., Lyčka A.: Collect. Czech. Chem. Commun. 1997, 62, 1888.
- 11. Lyčka A., Rys P., Skrabal P.: Magn. Reson. Chem. 1998, 36, 279.
- 12. Lyčka A., Luštinec D., Holeček J., Nádvorník M., Holčapek M.: Dyes Pigm. 2001, 50, 203.
- 13. Lyčka A., Holeček J.: Dyes Pigm. 2003, 57, 115.
- 14. Lyčka A.: Ann. Rep. NMR Spectrosc. 2000, 42, 1.
- 15. Straley J. M., Giles R. R.: U.S. 2857372 (1958); Chem. Abstr. 1958, 53, 37169.
- 16. Masoud M. S., Khalil E. A., Hindawy A. M., Ali A. E., Mohamed E. F.: Spectrochim. Acta, Part A 2004, 60, 2807.
- 17. Schetty G., Kuster W.: Helv. Chim. Acta 1961, 44, 2193.
- 18. Figgis B. N., Lewis J.: Prog. Inorg. Chem. 1964, 6, 183.
- 19. Mabbs F. E., Machin D. J.: *Magnetism and Transition Metal Complexes*, pp. 20–21. Chapman&Hall, London 1973.