


1,2-Azaborine Cations**

Adam J. V. Marwitz, Jesse T. Jenkins, Lev N. Zakharov, and Shih-Yuan Liu*

1,2-Dihydro-1,2-azaborine is a six-membered aromatic heterocycle that is isoelectronic with benzene through the replacement of a C=C unit in benzene with an isoelectronic B–N unit.^[1,2] Since the pioneering work by Dewar et al.,^[3,4] significant advances have been made in the synthesis and reactivity studies of this family of heterocycles.^[5–7] Our continued exploration of the 1,2-azaborine motif^[8–15] has led us to consider the synthesis of cationic 1,2-azaborines, for which no examples have been reported. In particular, we envisioned that substitution of 1,2-azaborine on the boron atom with pyridine derivatives would furnish cationic biaryl-type structures^[16] having the potential for use in materials applications (Scheme 1). Herein we report the synthesis,




Scheme 1. 1,2-Azaborine cations.


structural characterization, and optoelectronic properties of pyridine-substituted 1,2-azaborine cations, including a cationic heterocyclic analogue of *para*-terphenyl.

We have previously established nucleophilic substitution of the B–Cl bond in 1,2-azaborines by anionic nucleophiles (with Cl[−] serving as the leaving group).^[8] Less reactive neutral nucleophiles did not displace the chloride from the boron atom. We hypothesized that a better leaving group on the

boron atom (e.g., OTf) could render it susceptible to nucleophilic attack by weaker neutral nucleophiles. In the course of our studies, we discovered that silver reagents facilitate the ligand exchange at the boron position in 1,2-azaborines.^[13] We were thus pleased to discover that treatment of 1,2-azaborine **1** with AgOTf produced the substituted 1,2-azaborine **2** in 59% yield as an extremely moisture-sensitive liquid (Scheme 2). The 1,2-azaborine **2** was characterized by ¹H, ¹¹B, and ¹³C NMR spectroscopy as well as IR spectroscopy.

Scheme 2. Synthesis of **2**. Tf = trifluoromethanesulfonyl.

Heterocycle **2** readily reacts with *para*-substituted pyridines to form the desired cationic 1,2-azaborines **3**. As can be seen from Scheme 3, the substitution reaction is independent of the electronic nature of the nucleophile. Excellent yields have been obtained with both electron-rich and electron-poor pyridines.

Scheme 3. Synthesis of 1,2-azaborine cations **3**.

The 1,2-azaborine cations **3** are highly crystalline solids that fluoresce under UV light. We thus explored the solid-state fluorescence of the pyridine-substituted 1,2-azaborine cations **3**. The solid-state fluorescence and quantum yields of aromatic hydrocarbons, including *para*-terphenyl, have been reported using an integrating sphere.^[17] We have recorded the fluorescence spectra of crystalline samples of **3a–e** (Figure 1), all of which were freshly recrystallized prior to making the fluorescence measurements. The solid-state fluorescence of **3a** (R = Me) shows a peak at λ_{em} = 436 nm (Φ_{PL} = 0.03) and is visibly less fluorescent than samples of **3b** and **3c** under a UV lamp (λ = 365 nm; see Figure 1, right). The solid-state emission spectrum of **3b** (R = Ph) showed a relatively narrow band at λ_{em} = 448 nm (Φ_{PL} = 0.86). The high quantum yield observed for **3b** is quite similar to the values obtained for

[*] A. J. V. Marwitz, J. T. Jenkins, Dr. L. N. Zakharov, Prof. Dr. S.-Y. Liu
Department of Chemistry
University of Oregon
Eugene, OR 97403-1253 (USA)
E-mail: lsy@uoregon.edu

[**] Support for this research has been provided by the University of Oregon and the National Institutes of Health (National Institute of General Medical Sciences, Grant R01-GM094541). Funding for the University of Oregon Chemistry Research and Instrumentation Services has been furnished in part by the National Science Foundation (CHE-0234965). Correspondence concerning X-ray crystallography should be directed to Dr. L. N. Zakharov (lev@uoregon.edu).

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201004084>.




Figure 1. Normalized solid-state fluorescence spectra and images (under UV irradiation) of 1,2-azaborine cations **3**.

para-terphenyl, though the emission maximum of the cationic **3b** is bathochromically shifted relative to the all-carbon *para*-terphenyl by approximately 75 nm.^[17] We found that solid samples of **3c** (R=H) were blue-green fluorescent (see Figure 1) with a relatively broad emission maximum at λ_{em} = 469 nm (Φ_{PL} = 0.30). Compound **3d** (R = CF₃) exhibits yellow-green emission at λ_{em} = 527 nm (Φ_{PL} = 0.11), which contrasts the blue emission observed for geometrically similar **3a**. The data illustrated in Figure 1 suggest that the *para* substituent on pyridine has a substantial effect upon the emissive properties of 1,2-azaborine cations. DMAP-substituted **3e** (R = NMe₂; DMAP = 4-dimethylaminopyridine) does not fluoresce in the solid state (Figure 1, right), which is consistent with the reported fluorescence quenching by the presence of a nitrogen lone pair of electrons.^[18]

We also determined the absorption properties of 1,2-azaborine cations **3** in solution. The absorption maximum of **3a** (R = Me) in CH₂Cl₂ was found at λ = 287 nm with an extinction coefficient of ϵ = 12713 M⁻¹ cm⁻¹. The absorption spectrum of **3b** showed a broad, featureless peak at λ = 292 nm (ϵ = 21 869 M⁻¹ cm⁻¹), which is close to that observed for **3a**, but is slightly bathochromically shifted from the absorption maximum of *para*-terphenyl (observed at λ = 280 nm in CH₂Cl₂).^[19] The absorption spectrum of **3c** (R = H) also showed a broad peak at λ = 286 nm (ϵ = 8624 M⁻¹ cm⁻¹). The observed absorption peaks of **3d** at λ = 285 nm (ϵ = 8126 M⁻¹ cm⁻¹) and **3e** at λ = 283 nm (ϵ = 21 303 M⁻¹ cm⁻¹) are relatively unchanged from the other derivatives of **3**.

Interestingly, of the prepared cationic derivatives, only terphenyl analogue **3b** was found to be fluorescent in solution. The fluorescence spectrum of **3b** in CH₂Cl₂ showed an emission peak at λ_{em} = 360 nm (Φ_{PL} = 0.06), and the absorption maximum was found at λ = 292 nm in CH₂Cl₂ (Figure 2). The large Stokes shift of 68 nm for **3b** is indicative of considerable reorganization between the ground and the excited state. We also observed a bathochromic shift in the emission peak when MeCN was used as the solvent (λ_{em} = 382 nm, Φ_{PL} = 0.17) instead of CH₂Cl₂. Furthermore, the fluorescence of **3b** in MeCN was quenched upon the addition of NaI,^[20] highlighting the potential of these materials in sensing applications.





Figure 2. Normalized absorption (dashed line) and emission (solid line) spectra of **3b** in CH₂Cl₂.

Table 1 summarizes the photophysical properties of pyridine-substituted 1,2-azaborine cations **3a–e**. To assess whether the extended conjugation provided by the 1,2-azaborine ring is critical for the observed optoelectronic properties of the 1,2-azaborine cations **3**, we prepared the corresponding protonated pyridinium species **4a–e** (Scheme 4). We determined that under the same conditions

Table 1: Photophysical data for **3a–3e**.

Compound	Absorbance [nm] ^[a]	ϵ [M ⁻¹ cm ⁻¹] ^[a]	Emission [nm]	Φ_{pl}
3a	287	8126	436 ^[b]	0.03 ^[b]
3b	292	21 869	448 ^[b] 360 ^[a] 382 ^[c]	0.86 ^[b] 0.06 ^[a] 0.17 ^[c]
3c	286	8624	469	0.30 ^[b]
3d	287	12 713	527	0.11 ^[b]
3e	283	21 303	N/A	N/A

[a] In CH₂Cl₂ (10⁻⁵ M). [b] Solid-state emission. [c] In MeCN (10⁻⁵ M). N/A = not applicable.



Scheme 4.

for **3**, pyridinium triflate salts **4a–e** do not exhibit solid-state fluorescence emission in the visible region. This is consistent with a critical role of the 1,2-azaborine moiety in the observed emission properties of 1,2-azaborine cations **3**.

We have obtained the X-ray crystal structures of 1,2-azaborine cations **3** (except for **3c**), thus unambiguously establishing their structural identity. As a representative example, the solid-state structure of **3b** is illustrated in




Figure 3. ORTEP illustration of **3b**, with thermal ellipsoids drawn at the 35 % probability level (hydrogen atoms have been omitted for clarity). Bond distances (in Å): B-N(1) 1.413(2), B-N(2) 1.531(2), B-C(3) 1.496(2), C(3)-C(4) 1.369(2), C(4)-C(5) 1.408(3), C(5)-C(6) 1.358(2), C(6)-N(1) 1.3736(19); Torsion angles: 1,2-azaborine-pyridine = 50.5°; pyridine-phenyl = 18.0°.

Figure 3 and reveals that the pyridine nitrogen atom is bound to the boron atom with the triflate group serving as a noncoordinating anion.^[21] As expected, the dative exocyclic B–N bond (B–N(2) = 1.531(2) Å) in **3b** is significantly longer than the covalent exocyclic B–NPh₂ bond in a 1,2-azaborine recently reported in our group (B–N(2) = 1.486(2) Å).^[9] The exocyclic B–N bond in cationic **3b** is slightly shorter than the B–N bond in the charge-neutral borabenzene-4-phenylpyridine adduct (B–N = 1.551(3) Å) reported by Fu and co-workers.^[22] The 1,2-azaborine ring in **3b** is completely planar and is twisted by approximately 50° relative to the pyridine ring. In contrast, the phenyl ring of **3b** is only slightly twisted relative to the pyridine ring (18°).

We were also interested in examining the structural features of the 1,2-azaborine ring in **3b**. The intra-ring B–N bond is short (B–N(1) = 1.413(2) Å), as is the intra-ring B–C bond (B–C(3) = 1.496(2) Å), which is consistent with bond distances observed for electron-deficient 1,2-azaborines.^[13]

Selected bond parameters for 1,2-azaborine cations **3** are given in Table 2. The *para* substituent in the pyridine ring has little influence on the observed bond lengths, which are virtually identical for all derivatives. The torsion angles between the pyridine and 1,2-azaborine ring are similar for derivatives **3a**, **3b**, and **3d**, although in **3e** (R = NMe₂) the 1,2-azaborine ring is nearly perpendicular to the pyridine ring. It noteworthy that the 1,2-azaborine cations **3** represent a new family of borenium cations for which only a few members have been structurally characterized by single-crystal diffraction.^[23]

Table 2: Selected bond distances [in Å] and angles [°] for 1,2-azaborine cations **3**.

Compound	B–N(1) [Å]	B–C(3) [Å]	B–N(2) [Å]	Ring torsion ^[a]
3a	1.418(5)	1.481(5)	1.528(5)	57.1°
3b	1.413(2)	1.496(2)	1.531(2)	50.5°
3d	1.416(4)	1.489(4)	1.526(4)	58.0°
3e	1.423(5)	1.489(6)	1.527(5)	77.8°

[a] Torsion angle between the 1,2-azaborine and pyridine ring.

In summary, we prepared the first examples of 1,2-azaborine cations through a nucleophilic substitution reaction between pyridine nucleophiles and the highly Lewis acidic 1,2-Azaborine **2**. 1,2-Azaborine cations **3a–3d** exhibit solid-state fluorescence that is distinct from the neutral all-carbon analogues. Furthermore, 4-phenylpyridine-substituted 1,2-azaborine cation **3b**, an analogue of terphenyl, displays solution-phase fluorescence in addition to solid-state emission. Control experiments establish the 1,2-azaborine ring as an essential component for the observed optoelectronic properties. This study highlights the unique properties of 1,2-azaborine cations and underscores the potential utility of these complexes in materials applications.

Experimental Section

3b: In a glove box, a solution of 4-phenylpyridine (0.073 g, 0.47 mmol in 1.0 mL CH₂Cl₂) was added to a stirred solution of **2** (0.100 g, 0.392 mmol in 1.0 mL CH₂Cl₂). The mixture was stirred for 1 h at room temperature. At the conclusion of the reaction, the solution was cooled to –20°C and left at that temperature for 24 h. The desired product precipitated out of the solution as a crystalline solid. The supernatant was decanted and the crystallized product was washed with *n*-pentane (3 × 5 mL). Residual solvents were removed under reduced pressure to provide **3b** as clear, colorless crystals (0.155 g, 97 %). ¹H NMR (600 MHz, CD₂Cl₂): δ = 8.82 (d, ³J_{HH} = 6.9 Hz, 2H), 8.44 (d, ³J_{HH} = 6.9 Hz, 2H), 8.02 (dd, ³J_{HH} = 9.8, 6.6 Hz, 1H), 7.97 (dd, ³J_{HH} = 8.1 Hz, ⁴J_{HH} = 1.7 Hz, 2H), 7.68 (m, 3H), 7.55 (d, ³J_{HH} = 6.6, 1H), 6.85 (app t, ³J_{HH} = 7.5 Hz, 2H), 3.83 (q, ³J_{HH} = 7.3 Hz, 2H), 1.38 ppm (t, ³J_{HH} = 7.3 Hz, 3H). ¹³C NMR (75 MHz, CD₂Cl₂): δ = 158.1, 149.5, 145.8, 139.6, 134.3, 133.2, 130.5, 128.7, 125.7, 124 (br), 115.7, 47.7, 18.2 ppm. ¹¹B NMR (192.5 MHz, CD₂Cl₂): δ = 31.0 ppm. FTIR (thin film) 3220, 3138, 3078, 2915, 1638, 1612, 1513, 1488, 1474, 1442, 1412, 1377, 1349, 1292, 1233, 1218, 1174, 1151, 1029, 833, 765, 736, 693 cm^{–1}. HRMS (EI) calcd for C₇H₆BNO₃SiF₃ [M⁺] 255.03484, found 255.03528.

Received: July 5, 2010

Published online: September 6, 2010

Keywords: boron · cations · fluorescence · heterocycles

- [1] M. J. D. Bosdet, W. E. Piers, *Can. J. Chem.* **2009**, *87*, 8–29.
- [2] Z. Liu, T. B. Marder, *Angew. Chem.* **2008**, *120*, 248–250; *Angew. Chem. Int. Ed.* **2008**, *47*, 242–244.
- [3] M. J. S. Dewar, P. A. Marr, *J. Am. Chem. Soc.* **1962**, *84*, 3782.
- [4] K. M. Davies, M. J. S. Dewar, P. Rona, *J. Am. Chem. Soc.* **1967**, *89*, 6294–6297.
- [5] A. J. Ashe III, X. Fang, *Org. Lett.* **2000**, *2*, 2089–2091.
- [6] A. J. Ashe III, X. Fang, J. W. Kampf, *Organometallics* **2001**, *20*, 5413–5418.
- [7] A. J. Ashe III, *Organometallics* **2009**, *28*, 4236–4248.
- [8] A. J. V. Marwitz, E. R. Abbey, J. T. Jenkins, L. N. Zakharov, S.-Y. Liu, *Org. Lett.* **2007**, *9*, 4905–4908.
- [9] E. R. Abbey, L. N. Zakharov, S.-Y. Liu, *J. Am. Chem. Soc.* **2008**, *130*, 7250–7252.
- [10] A. J. V. Marwitz, M. H. Matus, L. N. Zakharov, D. A. Dixon, S.-Y. Liu, *Angew. Chem.* **2009**, *121*, 991–995; *Angew. Chem. Int. Ed.* **2009**, *48*, 973–977.
- [11] L. Liu, A. J. V. Marwitz, B. W. Matthews, S.-Y. Liu, *Angew. Chem.* **2009**, *121*, 6949–6951; *Angew. Chem. Int. Ed.* **2009**, *48*, 6817–6819.
- [12] A. N. Lamm, S.-Y. Liu, *Mol. Biosyst.* **2009**, *5*, 1303–1305.

- [13] A. J. V. Marwitz, S. P. McClintock, L. N. Zakharov, S.-Y. Liu, *Chem. Commun.* **2010**, 46, 779–781.
- [14] P. G. Campbell, L. N. Zakharov, D. J. Grant, D. A. Dixon, S.-Y. Liu, *J. Am. Chem. Soc.* **2010**, 132, 3289–3291.
- [15] A. M. Daly, C. Tanjaron, A. J. V. Marwitz, S.-Y. Liu, S. G. Kukolich, *J. Am. Chem. Soc.* **2010**, 132, 5501–5506.
- [16] For a neutral boron-containing biphenyl analogue, see: R. Boese, N. Finke, J. Henkelmann, G. Maier, P. Paetzold, H. P. Reisenauer, G. Schmid, *Chem. Ber.* **1985**, 118, 1644–1654.
- [17] R. Katoh, K. Suzuki, A. Furube, M. Kotani and K. Tokumaru, *J. Phys. Chem. C* **2009**, 113, 2961–2965.
- [18] S.-P. Van, G. S. Hammond, *J. Am. Chem. Soc.* **1978**, 100, 3895–3902.
- [19] See the Supporting Information for details.
- [20] Fluorescence quenching by iodide has been reported in *para*-terphenyl, see: A. R. Watkins, *J. Phys. Chem.* **1974**, 78, 2555–2558.
- [21] Crystallographic data for **3b**: $C_{18}H_{18}BF_3N_2O_3S$, $M_r = 410.21$, crystal size $0.36 \times 0.31 \times 0.24 \text{ mm}^3$, monoclinic, space group $P2_1/n$, $a = 9.3812(9)$, $b = 14.3671(13)$, $c = 14.2217(13) \text{ \AA}$, $\beta = 110.940(1)^\circ$, $V = 1882.0(3) \text{ \AA}^3$, $Z = 4$, $\rho_{\text{calc}} = 1.448 \text{ g cm}^{-3}$, $\mu = 0.223 \text{ mm}^{-1}$, $F(000) = 848$, $\text{MoK}\alpha$ -radiation $\lambda = 0.71073 \text{ \AA}$, $T = 173(2) \text{ K}$, $2\theta_{\text{max}} = 54.00^\circ$, 17178 reflections measured [$R_{\text{int}} = 0.0255$], 4101 reflections observed, 325 refined parameters, $R1 = 0.0376$, $wR2 = 0.1050$ for reflections with $I > 2\sigma(I)$, $R1 = 0.0451$, $wR2 = 0.1131$, and $\text{GOF} = 1.024$ for all data, max/min residual electron density $+0.439/-0.201 \text{ e \AA}^{-3}$. CCDC 782514 (**3b**) 782515 (**3a**), 782517 (**3d**), and 782516 (**3e**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [22] S. Qiao, D. A. Hoic, G. C. Fu, *Organometallics* **1997**, 16, 1501–1502.
- [23] W. E. Piers, S. C. Bourke, K. D. Conroy, *Angew. Chem.* **2005**, 117, 5142–5163; *Angew. Chem. Int. Ed.* **2005**, 44, 5016–5036.