Singlet Dioxygen Fixation

Reaction of Singlet Dioxygen with Phosphine–Borane Derivatives: From Transient Phosphine Peroxides to Crystalline Peroxoboronates**

Susana Porcel, Ghenwa Bouhadir, Nathalie Saffon, Laurent Maron, and Didier Bourissou*

Dedicated to Dr. Bernard Meunier

The last two decades has witnessed a spectacular renaissance in main group chemistry.^[1] Of particular interest are ambiphilic compounds that result from the combination of donor and acceptor moieties, typically involving Group 13 and 15 elements.^[2] Most strikingly, the groups of D. W. Stephan and G. Erker have shown that phosphine-boranes R₂P-spacer- $B(C_6F_5)_2$ (R = tBu or Mes (2,4,6-Me_3C_6H_2); spacer = p-C_6H_4 or CH₂CH₂) reversibly activate dihydrogen under mild conditions, opening new avenues in metal-free catalytic hydrogenation.^[3] Recently, reversible binding of carbon dioxide has also been reported.^[4] Moreover, our group has evidenced versatile coordination properties for mono- and polyphosphine-boranes featuring o-C₆H₄ linkers, which provide a better understanding of unusual interactions between transition metals and Lewis acids.^[5] The borane moiety ortho to phosphorus was also exploited to stabilize highly reactive adducts of phosphines. Accordingly, the reaction of R₂P-(o-C₆H₄)BMes₂ with phenyl azide was shown to give a phosphazide that is dramatically stabilized by intramolecular a N→B interaction that undergoes an unusual photoisomerization process.^[6] Furthermore, treatment of the o-phenylene phosphine-borane system with diethyl azodicarboxylate and phenyl isocyanate afforded stable versions of the zwitterionic intermediates involved in the Mitsunobu reaction and cyclooligomerization of isocyanates, respectively.^[7]

[*] Dr. S. Porcel, Dr. G. Bouhadir, Dr. D. Bourissou Université de Toulouse, UPS, LHFA 118 route de Narbonne, 31062 Toulouse (France) and
CNRS, LHFA UMR 5069
31062 Toulouse (France)
Fax: (+33) 5-6155-8204
E-mail: dbouriss@chimie.ups-tlse.fr
Dr. N. Saffon Structure Fédérative Toulousaine en Chimie Moléculaire Toulouse (France)
Dr. L. Maron
Université de Toulouse, INSA, UPS, LPCNO
135 avenue de Rangueil, 31077 Toulouse (France) and
CNRS, LPCNO UMR 5215
31077 Toulouse (France)

- [**] Financial support from the Centre National de la Recherche Scientifique, the Université de Toulouse and the COST action CM0802 PhoSciNet is gratefully acknowledged. L.M. thanks the Institut Universitaire de France and S. P. thanks the Ministery of Ciencia e Innovación of Spain for a Postdoctoral Fellowship.
 - Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201000520.

6186



We began to investigate the behavior of phosphine– borane compounds towards dioxygen with the previously described phosphine–borane $\mathbf{1}$.^[12] No sign of oxidation was observed after several hours when a toluene solution of $\mathbf{1}$ was subjected to air bubbling at room temperature.^[13] In marked contrast, compound $\mathbf{1}$ rapidly reacted with ¹O₂ (generated by irradiation of ³O₂ in the presence of tetraphenylporphyrin as photosensitizer) to afford a new compound $\mathbf{2}$ (Scheme 1). The



Scheme 1. Fixation of 1O_2 by the phosphine–borane **1**. TPP=tetraphenylporphyrin.

mass spectrum (the ion $(M+Na)^+$ was found at m/z 565) indicated the incorporation of two oxygen atoms, and ¹H NMR spectroscopy revealed the desymmetrization of the two mesityl substituents at boron. Single crystals were obtained in Et₂O at -20 °C, and the molecular structure of **2** was unambiguously established by an X-ray diffraction study (Figure 1, left). Accordingly, the phosphine–borane compound **1** has split the ¹O₂ molecule. One oxygen atom sits at phosphorus (leading to a phosphine oxide moiety), and the other has been inserted into one of the B–Mes bonds. The boron atom is in a trigonal planar environment, which



Figure 1. Molecular views of **2** (left) and **5** (right) in the solid state (ellipsoids set at 50% probability). Phenyl, mesityl, and methyl groups are simplified and the hydrogen atoms are omitted for clarity. For **5**, only one of the two independent molecules present in the asymmetric unit is shown. Selected bond lengths [Å] and angles [°]: **2**: P1–O1 1.484(1), B1–O2 1.359(2). **5**: P1–O1 1.534(3), B1–O1 1.647(6), B1–O2 1.408(6), B1–O3 1.454(6), O3–O4 1.473(4); P1-O1-B1 112.4(3), O2-B1-O3 115.1(4).

corresponds with the ¹¹B NMR signal observed at $\delta = 34.7$ ppm.

From a mechanistic viewpoint, it is most likely that ${}^{1}O_{2}$ is first trapped by the ambiphilic phosphine–borane **1**, and the ensuing phosphine peroxide **3** then rearranges by migration of a mesityl group from boron to oxygen.^[14] No intermediate could be detected by ${}^{31}PNMR$ monitoring at -80 °C, suggesting that the rearrangement **3** \rightarrow **2** proceeds spontaneously. Furthermore, the reaction is totally regioselective, with the oxygen atom being inserted into one of the B–Mes bond but not into the B–C₆H₄PPh₂ bond.

To study the influence of the substituents at boron, the two mesityl groups were replaced by a pinacol unit. The phosphine-boronate $4^{[15]}$ was also found to react spontaneously with ${}^{1}O_{2}$ (but not with ${}^{3}O_{2}$), leading to a single compound **5** (Scheme 2). The mass spectrum indicated the



Scheme 2. Fixation of ${}^{1}O_{2}$ by the phosphine-boronate 4.

incorporation of two oxygen atoms and the low field shift of the ³¹P NMR signal (from $\delta = -4.1$ ppm for 4 to $\delta = +$ 46.0 ppm for 5) substantiated the oxidation of the phosphorus atom. Furthermore, the presence of two sets of ¹H and ¹³C NMR signals for the CMe₂ groups revealed the desymmetrization of the pinacol moiety, and the ¹¹B NMR resonance signal observed at $\delta = +15.9 \text{ ppm}$ (compared to +31.1 ppm for 4) suggested a tetracoordinate environment around boron. Compound 5 was isolated as a white moisturesensitive solid by recrystallisation from diethyl ether at -20°C, and its molecular structure was definitely confirmed by X-ray crystallography (Figure 1, right). The phosphineboronate 4 has thus split the dioxygen molecule in a similar fashion to that observed with 1. In this case, one oxygen atom bridges the phosphorus and boron atoms, whilst the other oxygen atom has been inserted into the B(pin) ring, leading to a B-O-O skeleton. When the reaction of 4 with ${}^{1}O_{2}$ was performed with a circa 1:1 mixture of $({}^{16}O)_2$ and $({}^{18}O)_2$, the three isotopomers $({}^{16}O)_2$, $({}^{16}O{}^{18}O)$, and $({}^{18}O)_2$ of **5** were observed by mass spectrometry.^[15] The scrambling of the labeled oxygen atoms suggests a bimolecular process for the rearrangement of the initially formed **4**: ${}^{10}O_2$ adduct into **5**.

Organylperoxides of Group 13 elements (>E-O-O-R, with E = B, Al, Ga, or In) have attracted a surge of interest in the last two decades as key intermediates in autoxidation processes and as potential oxygen transfer reagents. However, these compounds are usually extremely reactive and rather unstable,^[16] which often prevents their isolation and complete characterization. The first alkylperoxides of a Group 13 element to be structurally authenticated were the dimeric $[\{(tBu)_2 E(\mu - OOtBu)\}_2]$ (E = Ga, In)^[17] and dinuclear $[(tBuOO)(tBuO)Al(\mu-OtBu)_2Al(mesal)_2]$ (mesal = methyl salicylate) derivatives.^[18] Recently, Uhl et al. took advantage of the peculiar electronic properties of the CH(SiMe₃)₂ group to prepare and crystallographically characterize a monomeric aluminum peroxide and several original di- and trinuclear gallium peroxides,^[19] all featuring oxidizing peroxo moieties (E-O-O) and also reducing E-C bonds. Not much is known either about the related organylperoxides of boron. To date, only one such compound has been characterized by X-ray diffraction: in 2009, Piers et al. described an endoperoxide adduct of a NHC-stabilized 9-boraanthracene.[20]

Compound 5 thus provides a rare example of a structurally authenticated B-O-O skeleton, and the first included in a peroxoboronate moiety.^[21-24] The O1-P1 bond distance of 5 (1.534(3) Å) is significantly longer than that of 2 (1.484(1) Å), in agreement with the additional interaction of O1 toward B (O1-B1: 1.647(6) Å). The ensuing five-membered ring is almost planar, and the six-membered ring linked in a spiro fashion at boron adopts a chair conformation. The B-O bond is slightly shorter for the boron-alkoxide than for the boronperoxide linkage (B1-O2: 1.408(6) Å versus B1-O3: 1.454(6) Å). The O3-O4 distance (1.473(4) Å) falls in the typical range for alkylperoxides of Group 13 elements,^[17-19] and the torsion angle across the O3-O4 bond (60.8°) is similar to those observed by Uhl et al. in gallium endoperoxides^[19b-d] (values of more than 90° are encountered in acyclic peroxides).

At this stage, it should be noted that the peroxo skeleton of > E-O-O-R compounds usually comes from a dioxygen molecule (insertion into E–C bonds) or a hydroperoxide (condensation reaction), whereas for **5**, the B-O-O fragment is obtained by insertion of one oxygen atom into a B–O bond.

To gain more insight into the structure and relative stability of the various phosphine–borane: ${}^{1}O_{2}$ adducts, DFT calculations were performed at the B3PW91/SDD(P,B),6-31G**(other atoms) level of theory on the model compound I featuring an ethylene glycol unit at boron.^[15] The optimized structures of the dioxygen adducts **II–IV** are shown in Figure 2. In the cyclic adduct **II**, the O1–O2 distance is 1.54 Å and the boron atom does not interact with the phosphadioxirane moiety (the BO1 and BO2 distances exceed 2.7 Å). The formation of **II** from $I + {}^{1}O_{2}$ is strongly favored thermodynamically ($\Delta G_{I \rightarrow II} = -32.8 \text{ kcal mol}^{-1}$). In marked contrast with that predicted computationally for simple phosphines,^[25] the corresponding phosphine peroxide

Communications



Figure 2. Optimized geometries and relative energies of the ${}^{1}O_{2}$ adducts deriving from the model phosphine-boronate I (free energies *G* at 25 °C relative to $I + {}^{1}O_{2}$).

derivative **III** was also found to be a minimum on the potential energy surface, located 10.7 kcalmol⁻¹ lower in energy.^[26] Compound **III** is stabilized by a strong intramolecular $O \rightarrow B$ interaction (B–O2 distance: 1.554 Å). Overall, the phosphine–borane skeleton bridges the two oxygen atoms (O1–O2 1.50 Å), leading to a puckered six-membered ring. Finally, the peroxoboronate form **IV** (related to **5**) is a further 14.2 kcalmol⁻¹ lower in energy compared to **III**, and its optimized geometry reproduced well the structure determined crystallographically (O2–O3 1.46 Å).

To substantiate the generality of the reaction observed between **4** and ${}^{1}O_{2}$, we then investigated the behavior of the related phosphine-borane compound **6**^[15] featuring a sixmembered-ring boronate. In this case, reaction of **6** with ${}^{1}O_{2}$ under the same conditions afforded the phosphine oxideboronate **7** (Scheme 3), as deduced from mass spectrometry (incorporation of a single oxygen atom) and NMR spectros-



Scheme 3. Fixation of ${}^{1}O_{2}$ by the phosphine-boronate **6**.

copy ($\delta^{31}P = +32.2$ ppm and $\delta^{11}B = +27.4$ ppm). Taking into account the fact that **6** is reluctant toward oxidization by ${}^{3}O_{2}$, we speculated that the phosphine oxide **7** was formed by a side reaction between unreacted **6** and a ${}^{1}O_{2}$ adduct thereof. To prevent this comproportionation process, the reaction between **6** and ${}^{1}O_{2}$ was carried out at -60 °C. Gratifyingly, ${}^{31}P$ NMR spectroscopy revealed the exclusive formation of a new product **8** ($\delta = +52.9$ ppm) that was isolated in 76 % yield by precipitation from a dichloromethane/pentane mixture at -20 °C. All the spectroscopic data for **8** support a peroxoboronate structure analogous to **5**, which was unambiguously confirmed by crystallography.^[15] The O–O distance of **8** (1.433(3) Å) is slightly shorter than that of **5**, whilst the torsion angle around the O–O bond widens to 98.7° as the result of the inclusion of the B-O-O-C skeleton in a sevenmembered ring. Oxygen transfer between the peroxoboronate 8 and the starting material 6 readily occurs at room temperature, as substantiated by the instantaneous and quantitative formation of the phosphine oxide 7 upon mixing both reagents in a 1:1 ratio. This result confirms that 7 is formed from 6 and ${}^{1}O_{2}$ at room temperature.

A similar oxygen transfer was also observed between the phosphine-boronate **4** and the ensuing peroxoboronate **5** (Scheme 4). This comproportionation reaction required 3 h at room temperature, and cleanly afforded the phosphine oxide **9**. Compound **9** was isolated in 64% yield after recrystalliza-



Scheme 4. Oxygen transfer reaction from 5 to 4.

tion in Et₂O/pentane and fully characterized by NMR and Xray diffraction.^[15] Finally, the ability of the peroxoboronates **5** and **8** to oxidize simple phosphines was examined using PPh₃ as a model substrate. According to ³¹P NMR, the reaction proceeds cleanly in both cases, leading to Ph₃PO ($\delta = +$ 25.0 ppm) together with the phosphine oxide **9** or **7**. In line with observations for the comproportionation reactions, the reaction proceeds faster with **8** than with **5**, complete conversion being achieved within 2 h at room temperature with **8** but requiring heating overnight at 100 °C with **5**.

In summary, *ortho*-phenylene phosphine–borane systems have been shown to readily react with singlet dioxygen. Starting from phosphine–boronates **4** and **6**, two stable peroxoboronates **5** and **8** have been isolated and fully characterized. The ability of these compounds to undergo oxygen transfer reactions has been substantiated. The potential application of such peroxoboronates as oxidizing agents and the extension of the present work to other phosphine– borane systems is currently under study in our group.

Experimental Section

All reactions and manipulations were carried out using standard Schlenk techniques. The NMR spectra were recorded at 25 °C.

General procedure for the reactions of the phosphine–borane compounds with ${}^{1}O_{2}$: A toluene solution of the phosphine–borane compound (concentrations ranging from 0.02 M to 0.09 M) containing 5–7 mol% of tetraphenylporphyrin was irradiated with a halogen lamp under a dried atmosphere of dioxygen until ${}^{31}P$ NMR spectros-copy indicated complete conversion. Toluene was removed under vacuum, and the residue was purified by crystallization in diethyl ether (for **2** and **5**) or precipitation in dichloromethane/pentane (for **8**).

2: ³¹P{¹H} NMR (202.5 MHz, C₆D₆): $\delta = 37.8$ ppm; ¹¹B NMR (160.5 MHz, CDCl₃): $\delta = 34.7$. HRMS (ESI⁺) calcd for $[M+Na]^+$ (C₃₆H₃₇BO₂P⁺): 565.2444, found: 565.2443. **5**: ³¹P NMR (202.5 MHz, CDCl₃): $\delta = 46.0$ ppm; ¹¹B NMR (160.5 MHz, CDCl₃): $\delta = 15.9$ ppm. HRMS (ESI⁺) calcd for $[M+H]^+$ (C₂₄H₂₇BO₄P⁺): 421.1740, found: 421.1757. **8**: ³¹P{¹H} NMR (202.5 MHz, C₆D₆): $\delta = 52.9$ ppm; ¹¹B NMR (160.5 MHz, C₆D₆): $\delta = 52.9$ ppm; ¹¹B NMR (160.5 MHz, C₆D₆): $\delta = 52.9$ ppm; ¹¹B NMR (160.5 MHz, C₆D₆): $\delta = 12.9$ ppm. HRMS (CI⁺) calcd for $[M+H]^+$ (C₂₃H₂₅BO₄P⁺): 407.1584, found: 407.1578.

Received: January 28, 2010 Revised: April 6, 2010 Published online: June 22, 2010

Keywords: ambiphilic compounds · boronates · oxygen transfer · phosphines · singlet oxygen

- For a recent review highlighting unexpected similarities between main group elements and transition metals, see: P. P. Power, *Nature* 2010, 463, 171–177. See also the special issue (number 33) of *Dalton Transactions* published in 2008.
- [2] a) D. W. Stephan, *Dalton Trans.* 2009, 3129–3136; b) D. W.
 Stephan, G. Erker, *Angew. Chem.* 2010, 122, 50–81; *Angew. Chem. Int. Ed.* 2010, 49, 46–76.
- [3] a) G. C. Welch, R. R. San Juan, J. D. Masuda, D. W. Stephan, Science 2006, 314, 1124–1126; b) P. A. Chase, G. C. Welch, T. Jurca, D. W. Stephan, Angew. Chem. 2007, 119, 8196–8199; Angew. Chem. Int. Ed. 2007, 46, 8050–8053; c) P. Spies, G. Erker, G. Kehr, K. Bergander, R. Fröhlich, S. Grimme, D. W. Stephan, Chem. Commun. 2007, 5072–5074; d) P. Spies, S. Schwendemann, S. Lange, G. Kehr, R. Fröhlich, G. Erker, Angew. Chem. 2008, 120, 7654–7657; Angew. Chem. Int. Ed. 2008, 47, 7543–7546.
- [4] C. M. Mömming, E. Otten, G. Kehr, R. Fröhlich, S. Grimme, D. W. Stephan, G. Erker, *Angew. Chem.* 2009, 121, 6770-6773; *Angew. Chem. Int. Ed.* 2009, 48, 6643-6646.
- [5] a) S. Bontemps, H. Gornitzka, G. Bouhadir, K. Miqueu, D. Bourissou, Angew. Chem. 2006, 118, 1641-1644; Angew. Chem. Int. Ed. 2006, 45, 1611-1614; b) S. Bontemps, G. Bouhadir, K. Miqueu, D. Bourissou, J. Am. Chem. Soc. 2006, 128, 12056-12057; c) M. Sircoglou, S. Bontemps, M. Mercy, N. Saffon, M. Takahashi, G. Bouhadir, L. Maron, D. Bourissou, Angew. Chem. 2007, 119, 8737-8740; Angew. Chem. Int. Ed. 2007, 46, 8583-8586; d) S. Bontemps, M. Sircoglou, G. Bouhadir, H. Puschmann, J. A. K. Howard, P. W. Dyer, K. Miqueu, D. Bourissou, Chem. Eur. J. 2008, 14, 731-740; e) S. Bontemps, G. Bouhadir, W. Gu, M. Mercy, C.-H. Chen, B. M. Foxman, L. Maron, O. V. Ozerov, D. Bourissou, Angew. Chem. 2008, 120, 1503-1506; Angew. Chem. Int. Ed. 2008, 47, 1481-1484; f) M. Sircoglou, S. Bontemps, G. Bouhadir, N. Saffon, K. Miqueu, W. Gu, M. Mercy, C.-H. Chen, B. M. Foxman, L. Maron, O. V. Ozerov, D. Bourissou, J. Am. Chem. Soc. 2008, 130, 16729-16738; g) M. Sircoglou, M. Mercy, N. Saffon, Y. Coppel, G. Bouhadir, L. Maron, D. Bourissou, Angew. Chem. 2009, 121, 3506-3509; Angew. Chem. Int. Ed. 2009, 48, 3454-3457; h) P. Gualco, T.-P. Lin, M. Sircoglou, M. Mercy, S. Ladeira, G. Bouhadir, L. M. Pérez, A. Amgoune, L. Maron, F. P. Gabbaï, D. Bourissou, Angew. Chem. 2009, 121, 10076-10079; Angew. Chem. Int. Ed. 2009, 48, 9892-9895; i) M. Sircoglou, S. Bontemps, M. Mercy, K. Miqueu, S. Ladeira, N. Saffon, L. Maron, G. Bouhadir, D. Bourissou, Inorg. Chem. 2010, 49, 3983-3990.
- [6] M. W. P. Bebbington, S. Bontemps, G. Bouhadir, D. Bourissou, Angew. Chem. 2007, 119, 3397–3400; Angew. Chem. Int. Ed. 2007, 46, 3333–3336.
- [7] S. Moebs-Sanchez, G. Bouhadir, N. Saffon, L. Maron, D. Bourissou, *Chem. Commun.* 2008, 3435–3437.
- [8] The lowest singlet state, referred to as ${}^{1}\Delta_{g}$, features one doublyoccupied π^{*} orbital and one empty π^{*} orbital, so singlet dioxygen can be considered as an ambiphilic molecule.
- [9] The frustrated Lewis pair tBu₃P/B(C₆F₅)₃ was recently found to readily bind nitrous oxide (N₂O): E. Otten, R. C. Neu, D. W. Stephan, J. Am. Chem. Soc. 2009, 131, 9918–9919.

- [10] a) R. Gao, D. G. Ho, T. Dong, D. Khuu, N. Franco, O. Sezer, M. Selke, Org. Lett. 2001, 3, 3719–3722; b) D. G. Ho, R. Gao, J. Celaje, H.-Y. Chung, M. Selke, Science 2003, 302, 259–262; c) D. Zhang, B. Ye, D. G. Ho, R. Gao, M. Selke, Tetrahedron 2006, 62, 10729–10733; d) D. Zhang, R. Gao, S. Afzal, M. Vargas, S. Sharma, A. McCurdy, M. Yousufuddin, T. Stewart, R. Bau, M. Selke, Org. Lett. 2006, 8, 5125–5128; e) J. J. Wilke, F. Weinhold, J. Am. Chem. Soc. 2006, 128, 11850–11859.
- [11] For the reaction of boranes with molecular oxygen, see: H. C. Brown in *Boranes in Organic Chemistry*, Cornell University Press, 1972.
- [12] T. W. Hudnall, Y.-M. Kim, M. W. P. Bebbington, D. Bourissou, F. P. Gabbai, J. Am. Chem. Soc. 2008, 130, 10890-10891.
- The inertness of 1 toward oxidation by ³O₂ is reminiscent of that reported by Buchwald et al. for biarylphosphines: T. E. Barder, S. L. Buchwald, *J. Am. Chem. Soc.* 2007, *129*, 5096-5101.
- [14] 1,2-Shifts of aryl groups from boron to oxygen have already been observed; see, for example: R. W. Hoffmann, K. Ditrich, *Synthesis* 1983, 107–109.
- [15] See the Supporting Information for details.
- [16] a) Y. A. Alexandrov, N. V. Chikinova, J. Organomet. Chem. 1991, 418, 1–59; b) A. R. Barron, Chem. Soc. Rev. 1993, 22, 93– 99.
- [17] a) W. M. Cleaver, A. R. Barron, J. Am. Chem. Soc. 1989, 111, 8966–8967; b) M. B. Power, W. M. Cleaver, A. W. Apblet, A. R. Barron, J. W. Ziller, *Polyhedron* 1992, 11, 477–486.
- [18] a) J. Lewiński, J. Zachara, E. Grabska, J. Am. Chem. Soc. 1996, 118, 6794–6795; b) J. Lewiński, J. Zachara, P. Goś, E. Grabska, T. Kopeć, I. Madura, W. Marciniak, I. Prowotorow, Chem. Eur. J. 2000, 6, 3215–3227.
- [19] a) W. Uhl, B. Jana, Chem. Eur. J. 2008, 14, 3067-3071; b) W. Uhl,
 M. R. Halvagar, Angew. Chem. 2008, 120, 1981-1983; Angew.
 Chem. Int. Ed. 2008, 47, 1955-1957; c) W. Uhl, M. R. Halvagar,
 F. Roge, W. Massa, Eur. J. Inorg. Chem. 2009, 489-492; d) W.
 Uhl, M. R. Halvagar, M. Layh, Chem. Commun. 2009, 4269-4271.
- [20] T. K. Wood, W. E. Piers, B. A. Keay, M. Parvez, Angew. Chem. 2009, 121, 4069–4072; Angew. Chem. Int. Ed. 2009, 48, 4009– 4012.
- [21] Sodium perborate, which is commonly used as a bleaching agent in the detergent industry, features O–O bridges: W. P. Griffith, A. C. Skapski, A. P. West, *Chem. Ind.* **1984**, 185.
- [22] The peroxoborate tBuOOB(en)(en=ethylene glycol) was prepared from tBuOOH and ClB(en) and used to oxidize organolithium and Grignard reagents. See Ref. [13] and: R. W. Hoffmann, B. Hölzer, O. Knopff, K. Harms, Angew. Chem. 2000, 112, 3206–3207; Angew. Chem. Int. Ed. 2000, 39, 3072–3074.
- [23] For a crystalline peroxoboronic acid that is κ²-O,O coordinated to rhodium, see: M. Ahijado, T. Braun, *Angew. Chem.* 2008, *120*, 2996–3000; *Angew. Chem. Int. Ed.* 2008, *47*, 2954–2958.
- [24] A peroxoborate complex [Rh]OOB(pin) (pin = tetramethylethylene glycol) has been characterized by NMR at low temperature upon reacting a η²-peroxo rhodium complex with pinacolborane HB(pin): M. A. Ahijado Salomon, T. Braun, A. Pener, *Angew. Chem.* 2008, 120, 8999–9003; *Angew. Chem. Int. Ed.* 2008, 47, 8867–8871.
- [25] a) K. Nahm, Y. Li, J. D. Evanseck, K. N. Houk, C. S. Foote, J. Am. Chem. Soc. **1993**, 115, 4879–4884; b) S. Tsuji, M. Kondo, K. Ishiguro, Y. Sawaki, J. Org. Chem. **1993**, 58, 5055–5059.
- [26] The phosphadioxirane II was predicted to rearrange into the phosphine peroxide III with a barrier height of only 5.1 kcal mol^{-1} .