

# 1,1-Carboboration of 1-Alkynes: A Conceptual Alternative to the Hydroboration Reaction


Chao Chen, Tanja Voss, Roland Fröhlich, Gerald Kehr, and Gerhard Erker\*

Organisch-Chemisches Institut, Westfälische Wilhelms-Universität, Corrensstrasse 40,  
48149 Münster, Germany

erker@uni-muenster.de

Received October 20, 2010

## ABSTRACT



Strongly electrophilic boranes  $\text{R}-\text{B}(\text{C}_6\text{F}_5)_2$  react readily with a variety of 1-alkynes by means of a 1,1-carboboration reaction to yield alkenylborane products, which can subsequently be used as reagents in metal catalyzed cross-coupling reactions.

The 1,1-carboboration of alkynes previously had been a rather specific reaction leading to boryl-functionalized alkenes.<sup>1</sup> Formally it might be regarded as an insertion reaction of the vinylidene isomer of the acetylene into a boron–carbon bond of the borane reagent.<sup>2</sup> However, mechanistically it proceeds by means of an abstraction/rearrangement sequence.<sup>3</sup> So far it required the presence of particular substituents at the alkyne, mostly main group or transition metal derived, that served as good 1,2-migrating groups in this scheme.<sup>1,3</sup> We had quite recently found that 1,1-carboboration reactions of simple 1-alkynes can easily be performed with the strongly electrophilic  $\text{R}-\text{B}(\text{C}_6\text{F}_5)_2$  re-

agents,<sup>4</sup> which are readily available.<sup>5</sup> In this account we will report on some significant progress including utilization of the resulting 1,1-carboboration products in cross-coupling reactions.

In a typical example, we mixed the terminal alkyne 5-phenyl-1-pentyne (**1a**) with  $\text{B}(\text{C}_6\text{F}_5)_3$  (**2a**)<sup>6</sup> in  $\text{CD}_2\text{Cl}_2$ . Inspection by NMR spectroscopy ( $^1\text{H}$ ,  $^{13}\text{C}$ ,  $^{19}\text{F}$ ,  $^{11}\text{B}$ ) revealed that the 1,1-carboboration reaction was complete within minutes to give a close to quantitative conversion to a ca. 1:1.2 mixture of the isomeric products *E*-**3a** and *Z*-**3a** (for details see the Supporting Information). Subsequent photolysis (HPK 125, Pyrex filter, 15 min, rt) resulted in a >95% conversion of the *E*-**3a** to the *Z*-**3a** isomer (see Scheme 1). We then performed this reaction on a preparative scale. In an one-pot procedure the thermally induced 1,1-carboboration reaction of the 1-alkyne **1a** with the borane **2a** was carried out in pentane at ambient conditions. Subsequent photolysis

(1) Wrackmeyer, B. *Coord. Chem. Rev.* **1995**, *145*, 125–156.

(2) In a formal sense this reaction is remotely related to the reverse of the Fritsch–Buttenberg–Wiechell reaction. Review: Tykwinski, R. R. *Chem. Commun.* **2010**, *46*, 3235–3249.


(3) Review: (a) Wrackmeyer, B. *Heteroat. Chem.* **2006**, *17*, 188–208, and references cited therein. See also selected examples: (b) Wrackmeyer, B.; Horchler, K.; Boese, R. *Angew. Chem.* **1989**, *101*, 1563–1565. (c) Wrackmeyer, B.; Horchler, K.; Boese, R. *Angew. Chem., Int. Ed. Engl.* **1989**, *28*, 1500–1502. (d) Wrackmeyer, B.; Kehr, G.; Boese, R. *Angew. Chem.* **1991**, *103*, 1374–1376. (e) Wrackmeyer, B.; Kehr, G.; Boese, R. *Angew. Chem., Int. Ed. Engl.* **1991**, *30*, 1370–1372. (f) Wrackmeyer, B.; Kehr, G.; Sebald, A.; Kümmerlen, J. *Chem. Ber.* **1992**, *125*, 1597–1603. (g) Wrackmeyer, B.; Kundler, S.; Milius, W.; Boese, R. *Chem. Ber.* **1994**, *127*, 333–342. (h) Wrackmeyer, B.; Kenner-Hofmann, B. H.; Milius, W.; Thoma, P.; Tok, O. L.; Herberhold, M. *Eur. J. Inorg. Chem.* **2006**, 101–108.

(4) (a) Chen, C.; Eweiner, F.; Wibbeling, B.; Fröhlich, R.; Senda, S.; Ohki, Y.; Tatsumi, K.; Grimme, S.; Kehr, G.; Erker, G. *Chem. Asian J.* **2010**, *5*, 2199–2208. (b) Chen, C.; Kehr, G.; Fröhlich, R.; Erker, G. *J. Am. Chem. Soc.* **2010**, *132*, 13594–13595.

(5) For their preparation, see the Supporting Information. See also: Spence, R. E. v. H.; Piers, W. E.; Sun, Y.; Parvez, M.; MacGillivray, L. R.; Zaworotko, M. J. *Organometallics* **1998**, *17*, 2459–2469.

(6) (a) Massey, A. G.; Park, A. J. *J. Organomet. Chem.* **1964**, *2*, 245–250. (b) Wang, C.; Erker, G.; Kehr, G.; Wedeking, K.; Fröhlich, R. *Organometallics* **2005**, *24*, 4760–4773.

Scheme 1




| <b>2</b> | <b>R</b>                   | <b>1</b> | <b>R<sup>1</sup></b>       | <b>3</b> | <b>E/Z</b>   | $h\nu$ | % <b>Z<sup>a</sup></b> |
|----------|----------------------------|----------|----------------------------|----------|--------------|--------|------------------------|
| <b>a</b> | $\text{C}_6\text{F}_5$     | <b>a</b> | $(\text{CH}_2)_3\text{Ph}$ | <b>a</b> | 1:1.2        |        | 63                     |
| <b>a</b> | $\text{C}_6\text{F}_5$     | <b>b</b> | $(\text{CH}_2)_4\text{Cl}$ | <b>b</b> | 1:1.9        |        | 83                     |
| <b>a</b> | $\text{C}_6\text{F}_5$     | <b>c</b> | $t\text{Bu}$               | <b>c</b> | <sup>b</sup> |        | 84                     |
| <b>b</b> | $\text{CH}_3$              | <b>d</b> | $\text{Pr}$                | <b>d</b> | 1:1.6        |        | 69                     |
| <b>c</b> | $(\text{CH}_2)_2\text{Ph}$ | <b>d</b> | $\text{Pr}$                | <b>e</b> | 1:1.4        |        | 69                     |

<sup>a</sup> Isolated yield. <sup>b</sup> Not determined.



(3 h) followed by workup eventually gave the pure **Z-3a** product in 63% yield. Similarly, the products **Z-3b** and **Z-3c** were obtained in good yield starting from **1b** and **1c**, respectively. The X-ray crystal structure analysis of **Z-3c** clearly shows the substituent pattern at the double bond originating from the 1,1-carbaboration reaction after subsequent irradiation.

Scheme 2




| <b>2</b> | <b>R</b>               | <b>4</b> | <b>n</b> | <b>R<sup>1</sup></b>                 | <b>5</b> | <b>Z/E</b>         | $h\nu$ | % <b>E<sup>a</sup></b> |
|----------|------------------------|----------|----------|--------------------------------------|----------|--------------------|--------|------------------------|
| <b>a</b> | $\text{C}_6\text{F}_5$ | <b>a</b> | 1        | $\text{CH}_2\text{C}\equiv\text{CH}$ | <b>a</b> | 1:1.8              |        | 62                     |
| <b>a</b> | $\text{C}_6\text{F}_5$ | <b>b</b> | 1        | $\text{CH}_3$                        | <b>b</b> | 1:2.8              |        | 49                     |
| <b>a</b> | $\text{C}_6\text{F}_5$ | <b>c</b> | 1        | $\text{TMS}$                         | <b>c</b> | 1:1.5              |        | 73                     |
| <b>a</b> | $\text{C}_6\text{F}_5$ | <b>d</b> | 2        | $\text{TMS}$                         | <b>d</b> | 1:1.0              |        | 56                     |
| <b>b</b> | $\text{CH}_3$          | <b>d</b> | 2        | $\text{TMS}$                         | <b>e</b> | 1:1.2 <sup>b</sup> |        | 58                     |

<sup>a</sup> Isolated yield. <sup>b</sup> 75 °C for 5 h.

Figure 1. Molecular structure of **Z-3c**.Figure 2. Molecular structures of the 1,1-carbaboration products **E-5a** (top) and **E-5b** (bottom).

Scheme 3



<sup>a</sup> A: starting from the isolated alkenylboranes **Z-3** or **E-5**, respectively. B: one-pot procedure. <sup>b</sup> For the respective alkenylborane see ref 4a. <sup>c</sup> Trimethylsilyl groups were removed during workup.

The 1,1-carboboration reaction of 1-alkynes is chemo-selective with regard to the substituents employed at the borane. We treated 1-pentyne with the reagent  $H_3C-B(C_6F_5)_2$  (**2b**) under similar conditions and found that exclusively the methyl substituent was transferred from boron to carbon in the course of the 1,1-carboboration reaction to give a mixture of the **Z-3d** and **E-3d** isomers. Subsequent photolysis again gave the pure **Z-3d** product (69% isolated, see Scheme 1). Similarly, we obtained the product **Z-3e** selectively from the 1,1-carboboration/photochemical isomerization sequence, in which only the 2-phenylethyl substituent had migrated.

Since we could selectively prepare substituted alkenyl boranes (**Z-3**) with a *Z*-configuration by this facile 1,1-carboboration/photolysis sequence, we searched for an *E*-selective alternative. This was found by employing 1-alkyne substrates bearing alkoxyalkyl or trimethylsiloxyalkyl substituents. A typical example is the reaction of bis-propargyl ether (**4a**) with  $B(C_6F_5)_3$  (**2a**). The 1,1-carboboration reaction took place readily at room temperature selectively with one alkynyl moiety to generate a ca. 1:1.8

mixture of the products **Z-5a** and **E-5a**. Subsequent photolysis again resulted in a fast and efficient isomerization of the carbon-carbon double bond of the product, but in this case the photostationary equilibrium lay on the side of the **E-5a** isomer (>95%, see Scheme 2). We isolated this crystalline product in 62% yield and characterized it by X-ray diffraction (see Figure 2). The X-ray crystal structure analysis confirmed the formation of **5a** by the 1,1-carboboration sequence and revealed stabilization of the borane product by internal oxygen to boron coordination (B1–O5 1.617(2) Å). This structural feature of compound **E-5a** is retained in solution (e.g.,  $^{11}B$  NMR,  $\delta$  9.1;  $^{19}F$  NMR,  $\Delta\delta_{m,p}[B(C_6F_5)_2] = 8.5$ ).<sup>7</sup>

The 1,1-carboboration/photolysis sequence of the  $-OCH_3$  or  $-OTMS$  functionalized 1-alkynes **4b–d** proceeds analogously. In each case we eventually isolated the respective *E*-alkenylborane product in good yield (see Scheme 2). Compound **5b** was also characterized by X-ray diffraction

(7) (a) Piers, W. E. *Adv. Organomet. Chem.* **2005**, 52, 1–76. (b) Beringhelli, T.; Donghi, D.; Maggioni, D.; D'Alfonso, G. *Coord. Chem. Rev.* **2008**, 252, 2292–2313, and references cited therein.

(see Figure 2). The crystal structure analysis confirmed the *E*-configuration of the C=C double bond in **5b** and shows the internal stabilization of the borane by internal oxygen to boron coordination. The oxygen–boron bond in **5b** (B2–O1 1.595(3) Å) is shorter than in **5a**.

The alkenylboranes **3** and **5** are interesting products in themselves as functionalized strong boron Lewis acids. They have now also been employed as reactive reagents in Pd-catalyzed cross-coupling reactions. For this purpose the isolated alkenylborane products were used (method **A**, Scheme 3), but it was also possible to alternatively employ these reagents generated in situ without additional workup in an one-pot procedure (method **B**). The reaction starting from 5-phenyl-1-pentyne (**1a**) is a typical example. Selective 1,1-carboboration with CH<sub>3</sub>B(CF<sub>3</sub>)<sub>2</sub> took place at room temperature; this was followed by photolysis and then Pd(PPh<sub>3</sub>)<sub>4</sub> (10 mol %) catalyzed coupling with phenyl iodide (see Scheme 3)<sup>8</sup> to selectively give the product *E*-**6b**, which was isolated in a yield of 76%. The -OTMS functionalized alkynes reacted similarly, only in these cases the corresponding alcohols (*E*-**6d**, *Z*-**6e**, see Scheme 3) were isolated in 90 or 73% yield, respectively. The *p*-cyanophenyl substituted alkene *Z*-**6f** was obtained in good yield from the Pd-catalyzed cross-coupling reaction of *Z*-**3c** with *p*-iodobenzonitrile.<sup>9</sup>

The new 1,1-carboboration reaction of simple 1-alkynes makes *Z*-substituted reactive alkenylboranes readily available. In principle, such products could also be prepared by a hydroboration route of the corresponding alkyne backbones with Pier's borane HB(CF<sub>3</sub>)<sub>2</sub>.<sup>10–12</sup> Of course, our procedure

shows a different pattern of forming the essential new bonds and, thus, provides a useful alternative to the hydroboration reaction,<sup>13</sup> especially in cases (e.g., **3d**, **3e**) where the latter might encounter regioselectivity problems. This new sequence makes the 1,1-carboboration reaction of 1-alkynes an interesting and useful methodological addition to the synthetic repertoire of organo-boron reagent chemistry.

**Acknowledgment.** Financial support from the Deutsche Forschungsgemeinschaft is gratefully acknowledged. C.C. thanks the Alexander von Humboldt-Stiftung for a stipend.

**Supporting Information Available:** Experimental procedures and spectroscopic data for all new compounds **3**, **5** and **6**. CIF files for compounds *Z*-**3c**, *E*-**5a** and *E*-**5b**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL102544X

(8) (a) Colberg, J. C.; Rane, A.; Vaquer, J.; Soderquist, J. A. *J. Am. Chem. Soc.* **1993**, *115*, 6065–6071. (b) Miyauro, N.; Suzuki, A. *Chem. Rev.* **1995**, *95*, 2457–2483. (c) Flynn, A. B.; Ogilvie, W. W. *Chem. Rev.* **2007**, *107*, 4698–4745.

(9) See for a comparison: (a) Ishiyama, T.; Mori, M.; Suzuki, A.; Miyauro, N. *J. Organomet. Chem.* **1996**, *525*, 225–231. (b) Alacid, E.; Nájera, C. J. *Org. Chem.* **2009**, *74*, 2321–2337.

(10) (a) Parks, D. J.; Spence, R. E. v H.; Piers, W. E. *Angew. Chem.* **1995**, *107*, 895–897. (b) Parks, D. J.; Spence, R. E. v H.; Piers, W. E. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 809–810. (c) Parks, D. J.; Piers, W. E.; Yap, G. P. A. *Organometallics* **1998**, *17*, 5492–5503.

(11) Reviews: (a) Trost, B. M.; Ball, Z. T. *Synthesis* **2005**, 853–887. (b) Suzuki, A.; Dhillon, R. S. *Topics in Current Chemistry*, Vol. 130; Boschke, F. L., Ed.; Springer, Berlin, 1986; pp 23–88.

(12) (a) Ishiyama, T.; Yamamoto, M.; Miyauro, N. *Chem. Lett.* **1996**, *25*, 1117–1118. See for comparison: (b) Satoh, Y.; Serizawa, H.; Miyauro, N.; Hara, S.; Suzuki, A. *Tetrahedron Lett.* **1988**, *29*, 1811–1814. (c) Sugimoto, M. *Chem. Rec.* **2010**, *10*, 348–358.

(13) General hydroboration reference (for alkynes): (a) Brown, H. C.; Bowman, D. H.; Misumi, S.; Unni, M. K. *J. Am. Chem. Soc.* **1967**, *89*, 4531–4532. (b) Zweifel, G.; Arzoumanian, H.; Whitney, C. C. *J. Am. Chem. Soc.* **1967**, *89*, 3652–3653. (c) Zweifel, G.; Fisher, R. P.; Snow, J. T.; Whitney, C. C. *J. Am. Chem. Soc.* **1971**, *93*, 6309–6311. (d) Brown, H. C.; Gupta, S. K. *J. Am. Chem. Soc.* **1972**, *94*, 4370–4371. (e) Brown, H. C.; Hamaoka, T.; Ravindran, N. *J. Am. Chem. Soc.* **1973**, *95*, 5786–5788. (f) Cha, J. S.; Min, S. J.; Kim, J. M.; Kwon, O. O. *Tetrahedron Lett.* **1993**, *34*, 5113–5116. (g) He, X.; Hartwig, J. F. *J. Am. Chem. Soc.* **1996**, *118*, 1696–1702. (h) Huang, Z.; Negishi, E.-I. *J. Am. Chem. Soc.* **2007**, *129*, 14788–14792.