

Synthesis of cortistatins A, J, K and L

Alec N. Flyer, Chong Si and Andrew G. Myers*

The cortistatins are a recently identified class of marine natural products characterized by an unusual steroidal skeleton, which have been found to inhibit differentially the proliferation of various mammalian cells in culture by an unknown mechanism. We describe a comprehensive route for the synthesis of cortistatins from a common precursor, which in turn is assembled from two fragments of similar structural complexity. Cortistatins A and J, and for the first time K and L, have been synthesized in parallel processes from like intermediates prepared from a single compound. With the identification of facile laboratory transformations linking intermediates in the cortistatin L synthetic series with corresponding intermediates to cortistatins A and J, we have been led to speculate that somewhat related paths might occur in nature, offering potential sequencing and chemical detail for cortistatin biosynthetic pathways.

Since the structure of cortistatin A was elucidated by Kobayashi and colleagues in 2006, more than ten natural cortistatins have been described, which have a common modified steroidal skeleton with varying substitution of the A- and D-rings^{1–3}. Many of these have profound effects on cultured mammalian cells, and on human umbilical vein endothelial cells (HUVECs) in particular^{1–4}. In 2008, Baran and colleagues reported the first laboratory synthetic route to cortistatin A (ref. 5), from prednisone, and subsequent to this a number of different approaches to the synthesis of cortistatins have been pursued, focused largely on the most highly functionalized (and potent) member of the family, cortistatin A (1). No fewer than three independent routes to cortistatin A have been reported to date^{5–7}, as well as a number of synthetic studies^{8–18} and formal routes^{19,20}. In addition, the Nicolaou–Chen group has described a synthesis of cortistatin J (ref. 21).

Structure–activity relationships among natural cortistatins and synthetic analogues have suggested that the 7-substituted isoquinoline appendage is a key determinant of the phenotypic effects of cortistatins upon cells grown in culture^{4,21–23}. The Nicolaou–Chen group recently reported that cortistatin A inhibits the function of a number of different kinases *in vitro* and proposed that the natural product may occupy the ATP-binding site of at least one of the enzymes they identified²⁴. Before this, Kobayashi and colleagues had shown that the addition of cortistatin A to cultured HUVECs led to a marked decrease in levels of an unidentified 110 kDa phosphoprotein⁴.

We envisioned that the naturally occurring cortistatins A, J, K and L (1–4, respectively, in Fig. 1) and a diverse array of cortistatin analogues, including compounds suitable for use as biological probes in target identification experiments, could be prepared from the single azido alcohol intermediate 5 (Fig. 2). We considered it likely that one or more pathways linking the azido alcohol 5 to the cyclohexadienone 6 could be developed. The oxabicyclic core of the latter precursor could then be disconnected by an oxidative cyclization transform, a strategy independently conceived and previously demonstrated by Sarpong and colleagues, as well as others^{8,15,16,18}. To allow for a convergent assembly process, we imagined an earlier disconnection of the seven-membered B-ring by an olefin metathesis reaction, giving rise to the triene precursor 7, which we envisioned could be synthesized by coupling of the *o*-vinyl benzylzinc reagent 8 and the enol triflate intermediate 9, structures of similar complexity, but less than certain viability as stable chemical entities.

Results

The *o*-vinyl benzylzinc reagent 8, the A-ring precursor, was derived from the benzyl bromide 14, which was prepared in amounts up to 47 g by the four-step sequence shown in Fig. 3 (via intermediates 10–13). Direct insertion of zinc²⁵ into the carbon–bromine bond of the benzyl bromide 14 was attempted only once, without success; however, a two-stage process for the preparation of organozinc reagent 8 proved to be both highly reliable and readily scalable. Addition of benzyl bromide 14 to activated magnesium turnings²⁶ in ether at 23 °C led to rapid (<30 min) formation of a deep green solution of the corresponding Grignard reagent, contaminated with <5% of the product of Wurtz coupling. In contrast, formation of the Grignard reagent in tetrahydrofuran as solvent was accompanied by as much as 65% of the Wurtz coupling by-product. Addition of the freshly prepared ethereal Grignard product to a solution of anhydrous zinc chloride in tetrahydrofuran at 23 °C led to the formation of a cloudy white suspension of the presumed *o*-vinyl benzylzinc reagent 8, which was used directly for coupling with the enol triflate 9 (see below).




Figure 1 | Cortistatins A, J, K and L and their IC₅₀ values measured in cultured HUVECs. Members of the cortistatin natural product family have a common modified steroidal skeleton with varying substitutions, primarily within the A, B and C rings. These substitutional variations are found to modulate the potencies of natural cortistatins as growth inhibitors of HUVECs in culture.




Figure 2 | A key precursor to cortistatins and its retrosynthetic disconnection. The azido alcohol intermediate **5** serves as a common intermediate for the synthesis of cortistatins A, J, K and L, as well as cortistatin analogues. This intermediate arises in turn from the cyclohexadienone **6** and the triene **7**. The latter intermediate is formed by coupling of the *o*-vinyl benzylzinc reagent **8** and the enol triflate intermediate **9**. TBS, *tert*-butyldimethylsilyl; TIPS, triisopropylsilyl; Tf, trifluoromethanesulfonyl.

The enol triflate coupling partner **9**, the CD-ring precursor, was synthesized in a three-step sequence from the known α -methylene ketone **15** (Fig. 4). The latter substance was first prepared in its antipodal form by Danishefsky and colleagues²⁷ (from the Hajos–Parrish enedione)²⁸ in the context of their synthesis of taxol, and was also selected as a starting material by the Nicolaou–Chen⁶ and Sorensen¹⁸ groups in their research directed toward the synthesis of cortistatins. As illustrated in Fig. 4, phosphoniosilylation^{29,30} of α -methylene ketone **15** with triphenylphosphine (1.02 equiv.) and triethylsilyl trifluoromethanesulfonate (1.02 equiv.) at -40°C in chloroform, uniquely effective as a solvent in this particular transformation, afforded the phosphonium salt **16** (containing a tetrasubstituted triethylsilyl enol ether), which was observed to be stable towards aqueous workup and extractive isolation. Previously, phosphoniosilylation–Wittig sequences have been conducted without isolation of the intermediate phosphonium salts³⁰; this was not an option in the present case, because chloroform was not compatible with formation of the Wittig reagent. The oily phosphonium salt obtained upon extractive isolation and concentration was dissolved in tetrahydrofuran, and after cooling to -78°C , the resulting solution was treated sequentially with *n*-butyllithium (1.1 equiv.) and paraformaldehyde (5 equiv.), affording the α -vinyl triethylsilyl enol ether **17** in 75% yield over the two steps from the α -methylene ketone **15**. Methodology for triflation developed by Corey and colleagues was used to transform the triethylsilyl enol ether **17** into the corresponding enol triflate derivative **9** (92% yield after purification by flash-column chromatography)³¹. This product, a colourless oil, was stored frozen as a solution in benzene for no longer than a few days; decomposition was observed to occur over longer periods of time.

Coupling of the enol triflate **9** (1 equiv.) with the benzylzinc reagent **8** (~1.2 equiv., a solution in ether–tetrahydrofuran) in the presence of *tris*(dibenzylideneacetone)dipalladium (0.045 equiv.) and the ligand 2-dicyclohexylphosphino-2',6'-dimethoxy-biphenyl (S-Phos, 0.18 equiv.)³² in *N*-methylpyrrolidone at 70°C (20 h) afforded the metathesis substrate **7** in 70% yield on a 16 g scale. Warming of a 0.2 M solution of the latter product with the second-generation Grubbs catalyst (0.025 equiv.)³³ in dichloromethane at 45°C for 5 h led to highly efficient ring-closing metathesis, forming the tetracyclic diene **18** (>95% yield, 11 g scale). In our initial route design we had planned to hydrogenate selectively the disubstituted alkene within this product and then epoxidize the remaining (tetra-substituted) alkene; however, we were not able to achieve selective 1,2-reduction of the diene. Sarpong and colleagues have reported the successful execution of this exact two-step

sequence, albeit using a substrate with different A-ring substituents (1-methoxyl and 2-*p*-methoxybenzyl groups)⁸.

We found that we could achieve the desired transformation (**18** \rightarrow **20**) by reversing the order of the two operations and in so doing we were able to avoid an isolation step. Thus, direct addition of a solution of dimethyldioxirane (1.5 equiv.) in acetone³⁴ to the unpurified metathesis product solution at 0°C led to stereoselective epoxidation of the tetrasubstituted alkene, affording the acid-sensitive tetracyclic diene monoepoxide **19**. Without purification, the epoxide **19** was hydrogenated in benzene under 500 psi hydrogen in the presence of a mixture of Wilkinson's catalyst (0.15 equiv.) and suspended solid sodium bicarbonate (1.5 equiv.), the latter being added as an acid scavenger. The saturated product, epoxide **20**, then underwent selective eliminative opening with lithium diethylamide (2 equiv.) in tetrahydrofuran at -15°C (ref. 35), furnishing the conjugated allylic alcohol **21**, a sufficiently polar intermediate to allow for facile chromatographic isolation (>95% purity, ^1H NMR (nuclear magnetic resonance) analysis, 50% yield over the four-step sequence). Cleavage of the triisopropylsilyl protective group with tetra-*n*-butylammonium fluoride (1.1 equiv.) in tetrahydrofuran and oxidative cyclization of the resulting phenol with [*bis*(trifluoroacetoxy)iodo]benzene (1.8 equiv.) in a mixture of dichloromethane and hexafluoroisopropanol provided the cyclohexadienone **6** in 50% yield^{8,18,36}.

The synthesis of the key azido alcohol intermediate **5** was achieved in three steps from the cyclohexadienone **6**, a sequence initiated by hydrosilylation with triethylsilane (2 equiv.) in the presence of Wilkinson's catalyst (0.05 equiv.)³⁷. The intermediate triethylsilyl enol ether was not isolated; rather, pyridine was added (14% by volume) followed by *N*-bromosuccinimide (2 equiv.), forming the (3*R*)-bromo ketone **22** as a single diastereomer in 70% yield. In the absence of pyridine, or with lesser quantities of pyridine, the bromination reaction was much less stereoselective, suggesting the lower selectivity was related to the presence of Wilkinson's catalyst during the bromination (bromination of the triethylsilyl enol ether generated in the absence of Wilkinson's catalyst was highly diastereoselective). Bromide displacement with tetramethylguanidinium azide (2 equiv.) proceeded with clean inversion of C3-stereochemistry and this was followed by direct reduction of the resulting (3*S*)- α -azido ketone with catecholborane (2 equiv.) in the presence of the (*R*)-Corey–Bakshi–Shibata (CBS) catalyst³⁸ (0.2 equiv., -40°C) and tetramethylguanidine (1 equiv.), producing a 15:1 mixture of the (2*S*)- and (2*R*)-configured alcohols, respectively, in 85% yield after chromatographic isolation. Similar stereocontrol of the C3-centre could not be achieved with other reductants examined (lithium tri-*tert*-butoxyaluminum hydride




Figure 3 | Preparation of the *o*-vinyl benzylzinc reagent **8.** The functionalized benzylzinc reagent **8** is prepared from the corresponding Grignard reagent and anhydrous zinc chloride. The Grignard reagent is readily formed from the benzyl bromide **14**, which is prepared in multigram amounts by a four-step sequence. THF, tetrahydrofuran; Ph, phenyl; DIBAL-H, diisobutylaluminium hydride.




Figure 4 | Synthesis of azido alcohol 5 from α -methylene ketone 15. The α -methylene ketone **15** is transformed in three steps to the enol triflate intermediate **9**. Coupling with the functionalized benzylzinc reagent **8** affords the triene **7**, which undergoes efficient ring-closing metathesis and oxidation (*in situ*, with dimethyldioxirane) to form the tetracyclic diene monoepoxide **19**. A sequence of hydrogenation, eliminative epoxide opening, silyl ether cleavage and oxidative cyclization provides cyclohexadienone **6**, which is converted stereoselectively in three steps to the key azido alcohol intermediate **5**. TES, triethylsilyl; DME, 1,2-dimethoxyethane; Pd₂dba₃, *tris*(dibenzylideneacetone)dipalladium(0); S-Phos, 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl; DMDO, dimethyldioxirane; TBAF, tetra-*n*-butylammonium fluoride; Py, pyridine; NBS, *N*-bromosuccinimide; TMG azide, tetramethylguanidinium azide; CBS, Corey-Bakshi-Shibata. d.r., diastereomeric ratio.

was the next most selective reductant, affording a 5:1 mixture of products favouring the (2*S*)-isomer).

Shown in parallel in Fig. 5 are the sequential pathways by which the key intermediate **5** was transformed into the corresponding 17-keto precursors to cortistatins A, J, K and L. The synthetic route to the 17-keto precursor to cortistatin A (**29**) is depicted first, and was initiated by protection of the 2-hydroxyl substituent of intermediate **5** with excess chloroacetyl chloride in pyridine. Cleavage of the *tert*-butyldimethylsilyl ether within the resulting product occurred upon exposure to aqueous hydrofluoric acid in a 1:1 mixture of acetonitrile and tetrahydrofuran at 0 °C, affording the alcohol **23** in 80% yield. Significantly, we observed that if during the latter transformation the reaction mixture was allowed to warm from 0 to 23 °C, 1,6-elimination of the chloroacetate ester accompanied deprotection of the *tert*-butyldimethylsilyl ether, an observation later used to advantage in the synthesis of the 17-keto precursor to cortistatin J (**32**). Oxidation of alcohol **23** with the Dess–Martin periodinane (2.4 equiv.) in dichloromethane at 0 °C followed by addition of methanol and solid potassium carbonate gave in a single operation the keto alcohol **24**, in 85% yield. Addition of *N*-bromosuccinimide (1.02 equiv.) to a solution of keto alcohol **24** in a 4:1 mixture of acetonitrile and methanol, respectively, at 23 °C led to stereoselective (*trans*-diaxial) 1,4-bromoetherification of the conjugated diene, forming the bromide **25**, which, without purification, was directly subjected to nucleophilic displacement at 0 °C with potassium superoxide (2.1 equiv.) in a 3:1 mixture of toluene and dimethyl sulfoxide, respectively^{39,40}. After extractive isolation and chromatography, the *trans*-diol methyl ether **26** was obtained in diastereomerically pure form (41% yield, 100 mg scale, two steps). When combined with scandium triflate (0.03 equiv.) and trifluoroacetic acid (9.4 equiv.) in dioxane at 23 °C, this substance was found to undergo efficient 1,2-elimination of methanol, affording the azido diol **27** (94% yield). The methoxyl substituent of substrate **26** is axially oriented and therefore stereoelectronically disposed towards elimination; none of the other allylic C–O bonds of substrate **26** (or product **27**) is so oriented. Reductive (di)methylation of the azido group

and diol protection with chlorotriethylsilane then furnished the protected 17-keto cortistatin A precursor (**29**) in 65% yield for the two steps (75 mg scale).

Syntheses of the 17-keto precursors to cortistatins J, K and L were substantially more direct (Fig. 5). The 17-keto cortistatin J precursor (**32**) was synthesized in just three steps from the azido alcohol **5**. Following reductive (di)methylation to form the dimethylamino alcohol **30** (85% yield), 1,6-elimination of water and deprotection of the *tert*-butyldimethylsilyl ether occurred simultaneously upon stirring the dimethylamino alcohol **30** with a biphasic mixture of chloroform and concentrated hydrochloric acid (20 min, 23 °C). Without purification, the product of the latter transformation (**31**) was subjected to oxidation with the Dess–Martin periodinane, providing the 17-keto cortistatin J precursor (**32**) as a white solid (77% yield following chromatographic purification, two steps).

The synthetic route to the 17-keto precursor to cortistatin K (**35**) also proceeded through the dimethylamino alcohol **30** (derived from the azido alcohol **5**, as described in the paragraph above), in a four-step sequence that was initiated by acetylation at 0 °C with acetic anhydride (a large excess) and scandium triflate as catalyst (0.05 equiv.) in a 1:1 mixture of acetonitrile and dichloromethane (93% yield)⁴¹. Regioselective reductive cleavage of the allylic C–O bond of the resulting acetate occurred in the presence of lithium borohydride (excess) and tetrakis(triphenylphosphine)-palladium (0.2 equiv.)⁴². This afforded the dimethylamino diene **34** in complex with borane; decomplexation was achieved by treatment of this intermediate with Raney nickel in methanol⁴³, providing the free amine as a pale yellow solid (90% yield after chromatographic purification over Davisil silica gel). Silyl ether cleavage (tetra-*n*-butylammonium fluoride, 87% yield) and oxidation (Dess–Martin periodinane, 90%) then completed the route to the 17-keto cortistatin K precursor (**35**), also obtained as a pale yellow solid.

Finally, the 17-keto precursor to cortistatin L (**38**) was synthesized from the azido alcohol **5** by a straightforward four-step sequence involving silyl ether deprotection (conducted at 0 °C, as in the cortistatin A series, to avoid 1,6-elimination), selective





a Cortistatin A series**b** Cortistatin J series**c** Cortistatin K series**d** Cortistatin L series

Figure 5 | Synthetic pathways to 17-keto precursors to cortistatins A, J, K and L from azido alcohol 5. **a**, Synthesis of the 17-keto precursor to cortistatin A (**29**) by an eight-step sequence. **b**, Three-step sequence to the 17-keto precursor to cortistatin J (**32**). **c**, Synthesis of the 17-keto precursor to cortistatin K (**35**) by a five-step sequence. **d**, Four-step sequence to the 17-keto precursor to cortistatin L (**38**). DMP, Dess–Martin periodinane; 18-Cr-6, 18-crown-6; TFA, trifluoroacetic acid; DMAP, 4-dimethylaminopyridine; Ac, acetyl; DBU, 1,8-diazabicyclo[5.4.0]undec-7-ene.




Figure 6 | Syntheses of cortistatins A, L, J and K. Parallel paths of 17-keto addition of 7-lithioisoquinoline followed by reductive deoxygenation transform each 17-keto cortistatin precursor into the corresponding cortistatin. TMEDA, *N,N,N',N'*-tetramethylethylenediamine; TFAA, trifluoroacetic anhydride; AIBN, 2,2'-azobis(2-methylpropionitrile).




Figure 7 | Chemical interconversions among intermediates of cortistatin series A, J and L and a hypothesized parallel biosynthetic sequence.

Intermediates of the cortistatin L series are found to undergo facile laboratory transformations to form corresponding intermediates of the A and J series, suggesting a plausible sequencing for the latter stages of cortistatin biosyntheses, which are unknown. Ms, methanesulfonyl.

protection⁴⁴, reductive (di)methylation and oxidation (Fig. 5, 60% yield overall).

Depicted in Fig. 6 are the parallel paths by which each advanced 17-keto cortistatin precursor was transformed into the corresponding cortistatin. The method we used to introduce stereoselectively the 7-isoquinolyl substituent differs from earlier reported syntheses of cortistatin A, where in each case a variation of the sequence 17-keto \rightarrow vinyl iodide^{5,7} or vinyl triflate⁶ followed by cross-coupling with a 7-metalloisoquinoline derivative, then stereoselective hydrogenation, was pursued. The hydrogenation step in particular has been established to be challenging to execute with intermediates containing the conjugated diene function of cortistatin A^{5,7}, and we believed it likely that this would be even more problematic with precursors to cortistatins J, K and L. As a general alternative, we developed the three-step 17-keto addition–reductive deoxygenation sequence shown in Fig. 6. While our work was in progress, Hirma and colleagues reported a conceptually similar but differently executed approach to introduce the isoquinolyl substituent in their formal synthesis of cortistatin A (ref. 19). We observed that lithiation of 7-iodoisoquinoline (5 equiv.) with an equimolar amount of *n*-butyllithium in tetrahydrofuran at -78°C followed by sequential addition of *N,N,N',N'*-tetramethylethylenediamine (TMEDA, 15 equiv.) and a 17-keto cortistatin precursor (1 equiv.) reproducibly afforded the corresponding (17*S*)-1,2-addition product in 52–62% yield in all cases⁴⁵. In the absence of TMEDA, 1,2-addition did not occur, undoubtedly a consequence of competing enolization of the 17-keto substituent. Enolization likely occurs, but to a much lesser extent, in our optimized conditions as well (in the presence of TMEDA) for we typically recover between 34–40% of each 17-keto precursor. The Hirma group developed a very different protocol for metalation (using *n*-butyllithium and cerium (III) chloride with 1-chloro-7-iodoisoquinoline as substrate) and with this organometallic reagent observed highly efficient addition to the carbonyl group of the Nicolaou–Chen 17-keto cortistatin A precursor. They then transformed the resulting tertiary alcohol into a thionocarbamate derivative and the latter was deoxygenated using tri-*n*-butyltin hydride¹⁹.

We briefly explored a similar method for hydroxyl activation in our deoxygenation studies, but our attempts to form a thionocarbonyl ester intermediate were unsuccessful. Instead, we adapted methodology by Jang and colleagues for free-radical deoxygenation of alcohols via their trifluoroacetate esters^{46,47}. As shown in Fig. 6,

each tertiary alcohol intermediate was readily transformed into the corresponding trifluoroacetate ester with trifluoroacetic anhydride and pyridine-*N,N*-dimethylaminopyridine. When heated with tri-*n*-butyltin hydride (excess) and azobisisobutyronitrile in benzene in a sealed tube at 100°C for 1–2 h, each trifluoroacetate ester underwent efficient deoxygenation to provide the (17*S*)-configured reduction product. Attempted deoxygenation under the conditions originally reported by Jang and colleagues (Ph_2SiH_2 , (*t*-BuO)₂, 130°C)^{46,47} led primarily to elimination of the trifluoroacetate ester, a problem that was almost completely circumvented by the use of tri-*n*-butyltin hydride as reductant at 100°C .

In the cortistatin J and K series, deoxygenation led to these substances directly; spectroscopic data for the synthetic materials matched values reported for the natural products. In the cortistatin A and L series, cleavage of the silyl ether protective group(s) with tetra-*n*-butylammonium fluoride at 23°C following deoxygenation provided these targets; spectroscopic data obtained from the synthetic substances were found to be the same as those reported for the natural products. In all four series (A, J, K and L), the yield for the two- or three-step deoxygenation (deprotection) procedure was 61–72% (1–20 mg amounts of final product).

In addition to spectroscopic characterization, synthetic cortistatins A, J, K and L were evaluated for their ability to inhibit the growth of HUVECs in culture (96-h incubation). The GI₅₀ values we measured are summarized in Fig. 1 together with literature values. For synthetic cortistatin A, we observed growth inhibition of HUVECs consistent with reported values, whereas measurements for synthetic cortistatin J more closely matched reports from the Nicolaou–Chen group than initial reports, and GI₅₀ values for synthetic cortistatins K and L fell within the range of those originally reported or were slightly higher^{1,3,21}.

Discussion

In the course of our studies we observed that several compounds closely related to cortistatin L undergo facile 1,6-elimination to form the conjugated triene function of cortistatin J (see **30** \rightarrow **31**, Fig. 5, discussion of intermediate **23** above, and the additional example of the transformation of dienyl alcohol **30** to triene **44**, Fig. 7). In an interesting and somewhat related transformation, we observed that reaction of diene **45** (cortistatin L oxidation state) with dimethyldioxirane afforded the highly sensitive epoxide **46** as the primary product, which upon standing in benzene solution

underwent spontaneous 1,4-elimination to form a product (dienyl alcohol **47**) with functionality characteristic of cortistatin A. Consideration of these findings has led us to speculate that cortistatins A and J may derive from cortistatin L or a precursor of the same oxidation state, as opposed to an alternative sequencing such as J → A (although this cannot be ruled out)². Cortistatin L might in turn be derived by C2 oxidation of cortistatin K. We note that many sterols bearing 2β-hydroxy and 2β-sulfato groups have been isolated from marine sponges, but detailed biosynthetic routes to these substances have not been determined^{48,49}.

We have shown that the protected azido alcohol intermediate **5**, synthesized in a nine-step sequence beginning with the coupling of the benzylzinc reagent **8** and the enol triflate **9**, is readily transformed into advanced 17-keto precursors to cortistatins A, J, K and L. Each of these intermediates is in turn converted into the corresponding cortistatin final product by a three- or four-step sequence involving addition of a 7-isoquinolyl organometallic intermediate followed by deoxygenation (deprotection). The latter sequence appears to be a general route to cortistatins with divergent substitutions of the A, B and C rings and it is anticipated that it will allow for late-stage introduction of diversely substituted isoquinoline groups and other heterocycles at position C17.

Received 26 May 2010; accepted 21 June 2010;
published online 1 August 2010

References

1. Aoki, S. *et al.* Cortistatins A, B, C and D, anti-angiogenic steroidal alkaloids, from the marine sponge *Corticium simplex*. *J. Am. Chem. Soc.* **128**, 3148–3149 (2006).
2. Watanabe, Y., Aoki, S., Tanabe, D., Setiawan, A. & Kobayashi, M. Cortistatins E, F, G and H, four novel steroidal alkaloids from marine sponge *Corticium simplex*. *Tetrahedron* **63**, 4074–4079 (2007).
3. Aoki, S. *et al.* Cortistatins J, K, L, novel abeo-9(10-19)-androstane-type steroidal alkaloids with isoquinoline unit, from marine sponge *Corticium simplex*. *Tetrahedron Lett.* **48**, 4485–4488 (2007).
4. Aoki, S. *et al.* Structure–activity relationship and biological property of cortistatins, anti-angiogenic spongean steroidal alkaloids. *Bioorg. Med. Chem.* **15**, 6758–6762 (2007).
5. Shenvi, R. A., Guerrero, C. A., Shi, J., Li, C.-C. & Baran, P. S. Synthesis of (+)-cortistatin A. *J. Am. Chem. Soc.* **130**, 7241–7243 (2008).
6. Nicolaou, K. C., Sun, Y. P., Peng, X. S., Polet, D. & Chen, D. Y. K. Total synthesis of (+)-cortistatin A. *Angew. Chem. Int. Ed.* **47**, 7310–7313 (2008).
7. Lee, H. M., Nieto-Oberhuber, C. & Shair, M. D. Enantioselective synthesis of (+)-cortistatin A, a potent and selective inhibitor of endothelial cell proliferation. *J. Am. Chem. Soc.* **130**, 16864–16866 (2008).
8. Simmons, E. M., Hardin, A. R., Guo, X. & Sarpong, R. Rapid construction of the cortistatin pentacyclic core. *Angew. Chem. Int. Ed.* **47**, 6650–6653 (2008).
9. Yamashita, S., Iso, K. & Hirama, M. A concise synthesis of the pentacyclic framework of cortistatins. *Org. Lett.* **10**, 3413–3415 (2008).
10. Craft, D. T. & Gung, B. W. The first transannular [4 + 3] cycloaddition reaction: synthesis of the ABCD ring structure of cortistatins. *Tetrahedron Lett.* **49**, 5931–5934 (2008).
11. Dai, M. J. & Danishefsky, S. J. A concise synthesis of the cortistatin core. *Tetrahedron Lett.* **49**, 6610–6612 (2008).
12. Dai, M. J., Wang, Z. & Danishefsky, S. J. A novel α,β-unsaturated nitron-aryne [3 + 2] cycloaddition and its application in the synthesis of the cortistatin core. *Tetrahedron Lett.* **49**, 6613–6616 (2008).
13. Kotoku, N., Sumii, Y., Hayashi, T. & Kobayashi, M. Synthesis of CD-ring structure of cortistatin A, an anti-angiogenic steroidal alkaloid from marine sponge. *Tetrahedron Lett.* **49**, 7078–7081 (2008).
14. Kurti, L., Czako, B. & Corey, E. J. A short, scalable synthesis of the carbocyclic core of the anti-angiogenic cortistatins from (+)-estrone by B-ring expansion. *Org. Lett.* **10**, 5247–5250 (2008).
15. Dai, M. J. & Danishefsky, S. J. An oxidative dearomatization cyclization model for cortistatin A. *Heterocycles* **77**, 157–161 (2009).
16. Liu, L. Z. *et al.* A model study for the concise construction of the oxapentacyclic core of cortistatins through intramolecular Diels–Alder and oxidative dearomatization–cyclization reactions. *Chem. Commun.* 662–664 (2009).
17. Magnus, P. & Littich, R. Intramolecular cyclopropene-furan [2 + 4] cycloaddition followed by a cyclopropylcarbinyl rearrangement to synthesize the BCD rings of cortistatin A. *Org. Lett.* **11**, 3938–3941 (2009).
18. Frie, J. L., Jeffrey, C. S. & Sorensen, E. J. A hypervalent iodine-induced double annulation enables a concise synthesis of the pentacyclic core structure of the cortistatins. *Org. Lett.* **11**, 5394–5397 (2009).
19. Yamashita, S., Kitajima, K., Iso, K. & Hirama, M. Efficient and stereoselective installation of isoquinoline: formal total synthesis of cortistatin A. *Tetrahedron Lett.* **50**, 3277–3279 (2009).
20. Simmons, E. M., Hardin-Narayan, A. R., Guo, X. & Sarpong, R. Formal total synthesis of (±)-cortistatin A. *Tetrahedron* **66**, 4696–4700 (2010).
21. Nicolaou, K. C. *et al.* Total synthesis and biological evaluation of cortistatins A and J and analogues thereof. *J. Am. Chem. Soc.* **131**, 10587–10597 (2009).
22. Shi, J. *et al.* Stereodivergent synthesis of 17-α and 17-β-aryl steroids: application and biological evaluation of D-ring cortistatin analogues. *Angew. Chem. Int. Ed.* **48**, 4328–4331 (2009).
23. Czako, B., Kurti, L., Mammoto, A., Ingber, D. E. & Corey, E. J. Discovery of potent and practical antiangiogenic agents inspired by cortistatin A. *J. Am. Chem. Soc.* **131**, 9014–9019 (2009).
24. Cee, V. J., Chen, D. Y. K., Lee, M. R. & Nicolaou, K. C. Cortistatin A is a high-affinity ligand of protein kinases ROCK, CDK8 and CDK11. *Angew. Chem. Int. Ed.* **48**, 8952–8957 (2009).
25. Berk, S. C., Knochel, P. & Yeh, M. C. P. General approach to highly functionalized benzylic organometallics of zinc and copper. *J. Org. Chem.* **53**, 5789–5791 (1988).
26. Pearson, D. E., Cowan, D. & Beckler, J. D. A study of the entrainment method for making Grignard reagents. *J. Org. Chem.* **24**, 504–509 (1959).
27. Isaacs, R. C. A., Digrandi, M. J. & Danishefsky, S. J. Synthesis of an enantiomerically pure intermediate containing the CD substructure of taxol. *J. Org. Chem.* **58**, 3938–3941 (1993).
28. Hajos, Z. G. & Parrish, D. R. (+)-(7aS)-7a-methyl-2,3,7a-tetrahydro-1H-indene-1,5-dione. In *Organic Syntheses*, Vol. 63, 26–31 (Wiley & Sons, 1985).
29. Evans, D. A., Hurst, K. M. & Takacs, J. M. New silicon-phosphorus reagents in organic synthesis: carbonyl and conjugate addition reactions of silicon phosphite esters and related systems. *J. Am. Chem. Soc.* **100**, 3467–3477 (1978).
30. Kozikowski, A. P. & Jung, S. H. Phosphoniosylation: an efficient and practical method for the β-functionalization of enones. *J. Org. Chem.* **51**, 3400–3402 (1986).
31. Mi, Y., Schreiber, J. V. & Corey, E. J. Total synthesis of (+)-α-onocerin in four steps via four-component coupling and tetracyclization steps. *J. Am. Chem. Soc.* **124**, 11290–11291 (2002).
32. Walker, S. D., Barder, T. E., Martinelli, J. R. & Buchwald, S. L. A rationally designed universal catalyst for Suzuki–Miyaura coupling processes. *Angew. Chem. Int. Ed.* **43**, 1871–1876 (2004).
33. Scholl, M., Ding, S., Lee, C. W. & Grubbs, R. H. Synthesis and activity of a new generation of ruthenium-based olefin metathesis catalysts coordinated with 1,3-dimesityl-4,5-dihydroimidazol-2-ylidene ligands. *Org. Lett.* **1**, 953–956 (1999).
34. Murray, R. W. & Singh, M. Synthesis of epoxides using dimethyldioxirane: trans-stilbene oxide. In *Organic Syntheses*, Vol. 74, 91–100 (Wiley & Sons, 1997).
35. Thummel, R. P. & Rickborn, B. Base-induced rearrangement of epoxides. V. Phenyl-substituted epoxides. *J. Org. Chem.* **37**, 3919–3923 (1972).
36. Kita, Y., Tohma, H., Kikuchi, K., Inagaki, M. & Yakura, T. Hypervalent iodine oxidation of N-acetyltyramines: synthesis of quinol ethers, spirohexadienones and hexahydroindol-6-ones. *J. Org. Chem.* **56**, 435–438 (1991).
37. Kogure, T. & Ojima, I. Reduction of carbonyl compounds via hydrosilylation. 4. Highly regioselective reductions of α,β-unsaturated carbonyl compounds. *Organometallics* **1**, 1390–1399 (1982).
38. Corey, E. J. & Helal, C. J. Reduction of carbonyl compounds with chiral oxazaborolidine catalysts: a new paradigm for enantioselective catalysis and a powerful new synthetic method. *Angew. Chem. Int. Ed.* **37**, 1986–2012 (1998).
39. San Filippo, J., Chern, C. I. & Valentine, J. S. Reaction of superoxide with alkyl halides and tosylates. *J. Org. Chem.* **40**, 1678–1680 (1975).
40. Corey, E. J., Nicolaou, K. C., Shibasaki, M., Machida, Y. & Shiner, C. S. Superoxide ion as a synthetically useful oxygen nucleophile. *Tetrahedron Lett.* **37**, 3183–3186 (1975).
41. Ishihara, K., Kubota, M., Kurihara, H. & Yamamoto, H. Scandium trifluoromethanesulfonate as an extremely active acylation catalyst. *J. Am. Chem. Soc.* **117**, 4413–4414 (1995).
42. Hutchins, R. O., Learn, K. & Fulton, R. P. Reductive displacement of allylic acetates by hydride transfer via catalytic activation by palladium(0) complexes. *Tetrahedron Lett.* **21**, 27–30 (1980).
43. Couturier, M., Tucker, J. L., Andresen, B. M., Dube, P. & Negri, J. T. Palladium and Raney nickel catalyzed methanolic cleavage of stable borane–amine complexes. *Org. Lett.* **3**, 465–467 (2001).
44. Moon, S. S., Stuhmiller, L. M. & McMorris, T. C. Synthesis of oregonol. *J. Org. Chem.* **54**, 26–28 (1989).
45. Foy, N. *et al.* Synthesis, receptor binding, molecular modeling and proliferative assays of a series of 17α-arylestradiols. *Chembiochem.* **4**, 494–503 (2003).

46. Jang, D. O., Kim, J. G., Cho, D. H. & Chung, C. M. Radical deoxygenation of alcohols via their trifluoroacetate derivatives with diphenylsilane. *Tetrahedron Lett.* **42**, 1073–1075 (2001).
47. Kim, J. G., Cho, D. H. & Jang, D. O. Radical deoxygenation of tertiary alcohols via trifluoroacetates. *Tetrahedron Lett.* **45**, 3031–3033 (2004).
48. D' Auria, M. V., Minale, L. & Riccio, R. Polyoxygenated steroids of marine origin. *Chem. Rev.* **93**, 1839–1895 (1993).
49. Sarma, N. S., Krishna, M. S. R. & Rao, S. R. Sterol ring system oxidation pattern in marine sponges. *Mar. Drugs* **3**, 84–111 (2005).

Acknowledgements

Financial support from the National Institutes of Health (Stimulus grant no. CA047148-22S1) is gratefully acknowledged. A.N.F. acknowledges a scholarship from Eli Lilly and Company. We gratefully acknowledge G. Zou for measuring the GI_{50} values of cortistatins A, J, K and L and A.W.G. Burgett and M.D. Shair for their assistance with these

measurements as well as a gift of the HUVEC. We acknowledge helpful discussions with C. T. Walsh and E. Balskus.

Author contributions

A.N.F., C.S. and A.G.M. conceived the synthetic route. A.N.F. and C.S. conducted all experimental work and analysed the results. A.N.F., C.S. and A.G.M. wrote the manuscript.

Additional information

The authors declare no competing financial interests. Supplementary information and chemical compound information accompany this paper at www.nature.com/naturechemistry. Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>. Correspondence and requests for materials should be addressed to A.G.M.