

Cooperative catalysis by carbenes and Lewis acids in a highly stereoselective route to γ -lactams

Dustin E. A. Raup, Benoit Cardinal-David, Dane Holte and Karl A. Scheidt*

Enzymes are a continuing source of inspiration for the design of new chemical reactions that proceed with efficiency, high selectivity and minimal waste. In many biochemical processes, different catalytic species, such as Lewis acids and bases, are involved in precisely orchestrated interactions to activate reactants simultaneously or sequentially. This type of 'cooperative catalysis', in which two or more catalytic cycles operate concurrently to achieve one overall transformation, has great potential in enhancing known reactivity and driving the development of new chemical reactions with high value. In this disclosure, a cooperative *N*-heterocyclic carbene/Lewis acid catalytic system promotes the addition of homoenolate equivalents to hydrazones, generating highly substituted γ -lactams in moderate to good yields and with high levels of diastereo- and enantioselectivity.

Biological systems have evolved to activate substrates using acids and bases simultaneously within an active site. Proteases achieve astounding efficiency in the hydrolysis of robust amide bonds by making use of a serine or cysteine (Lewis base), an aspartic acid residue (Brønsted acid) and a histidine residue (Brønsted base) in concert. This catalytic triad is essential for life processes, and the precise placement of these catalytic functional groups precludes non-productive interactions. In general, nature takes advantage of enzyme architecture to facilitate a cooperative reaction manifold, because many catalytic sites contain functional groups and/or metals that would be incompatible in simple solution-based chemistry. For example, the reactive thiol of an acyl transferase would bind to the active iron centre in a cytochrome if either one was removed from their protein framework. It has been a significant challenge to capitalize on this powerful cooperative concept in abiotic systems. In systems using 'cooperative catalysis', the combination of Lewis acid and Lewis base activation can achieve greater selectivity and more efficient reactivity than either mode individually, while simultaneously avoiding any substantial self-inhibition^{1–6}.

However, homogeneous catalytic reactions promote random collisions of all species in solution, facilitating simple acid–base chemistry or the generation of stable Lewis acid–Lewis base complexes that will compete with a desired catalytic reaction. Inspired by the thiamine pyrophosphate cofactor^{7,8}, *N*-heterocyclic carbenes (NHC) have emerged as unique and efficient Lewis base catalysts^{9–11} for a host of new reactions over the last decade, including the generation of acyl anion equivalents^{12–17}, homoenolate equivalents^{18–23}, enolates^{24,25} and acylvinyl anion equivalents²⁶. However, NHCs have not been applied cooperatively with Lewis acids to generate new carbon–carbon bonds, presumably because these species act as exceptional ligands for late transition metals (Fig. 1)^{27–33}. This issue has been overcome previously in the literature in a reaction using NHCs and palladium in separate catalytic cycles³⁴.

We disclose a cooperative and catalytic NHC/Lewis acid system, and demonstrate its utility with the highly enantioselective addition of homoenolate equivalents to Lewis acid-activated *N*-benzoyl hydrazones. In this novel strategy, the NHC combines transiently with an aldehyde to generate an unusual nucleophile, raising the

energy of the highest occupied molecular orbital (HOMO). Simultaneously, an optimal Lewis acid catalyst activates the hydrazone by lowering the energy of the lowest unoccupied molecular orbital (LUMO) for an overall productive reaction.

We have been investigating new classes of electrophiles with homoenolate equivalents generated using NHC catalysis to gain direct access to highly useful γ -amino acid derivatives^{35,36} or natural products³⁷. To achieve this goal, an effective class of imine was required that could deliver highly substituted γ -lactams with high levels of enantioselectivity and diastereoselectivity^{23,38}. If successful, this formal [3 + 2] annulation process could, from two simple starting materials and a small organic molecule catalyst, directly produce important five-membered heterocycles with known biological activity and high occurrence in the structures of natural products. *N*-acyl hydrazones are well-known substrates for both Lewis acid-activated cycloadditions and nucleophilic allylation reactions³⁹. In these processes, the hydrazone substrates presumably undergo activation by chelation to a Lewis acid with the carbonyl oxygen and imine nitrogen atoms to form an activated electrophile *in situ*. Our studies exploring the formal cycloadditions using *N*-benzoyl hydrazones and various unsaturated aldehydes were initiated by treating cinnamaldehyde and hydrazone **1a** with a variety of azolium salts and subjecting the mixture to a number of reaction conditions. The catalyst generated from benzimidazolium **4** displayed promising reactivity during the preliminary screening, although incomplete conversion of **1a** was observed. After surveying both amine and inorganic bases, the bicyclic guanidine base 1,5,7-triazabicyclo[4.4.0]dec-5-ene (TBD) was found to efficiently promote the formal [3 + 2] cycloaddition. The basicity of TBD in tetrahydrofuran (THF) is higher (TBDH⁺ pK_a = 21.0) than diazabicycloundec-7-ene (DBU) (DBUH⁺ pK_a = 16.8), which has been heavily used as a base in reported carbene-catalysed reactions⁴⁰. With these initial conditions, we examined the impact of chiral triazolium salts on the efficiency and stereoselection of the process. Although the reaction progressed with moderate selectivity using chiral triazolium **5** (72% e.e., entry 3, Table 1), higher levels of absolute stereocontrol could not be obtained by lowering the temperature, changing the base or using various protic additives (not shown). Additionally, the yield of the reaction dropped from 70%

Department of Chemistry, Center for Molecular Innovation and Drug Discovery, Northwestern University, 2145 Sheridan Road, Evanston, Illinois 60208, USA. *e-mail: scheidt@northwestern.edu











Figure 1 | Opportunities with NHCs and metals. It has been comprehensively established that NHCs react with enals to afford a new nucleophilic species and that metal salts can activate chelating electrophiles such as *N*-acyl hydrazones. Can both of these processes operate in one flask to access reactivity that is otherwise impossible/inefficient?

Table 1 | Development of a cooperative carbene catalysis reaction.



Entry	NHC/Lewis acid	Base	% yield*	d.r. [†]	% e.e. [‡]
1	20 mol% 4	1.2 equiv. DBU	58	6:1	—
2	20 mol% 4	20 mol% TBD	70	12:1	—
3	20 mol% 5	25 mol% TBD	62	6:1	72
4	20 mol% 5 /Mg(Ot-Bu) ₂	25 mol% TBD	78	7:1	86
5	20 mol% 6 /Mg(Ot-Bu) ₂	25 mol% TBD	84	7:1	80
6	20 mol% 7	25 mol% TBD	63	6:1	90
7	20 mol% 7 /Mg(Ot-Bu) ₂	25 mol% TBD	79	7:1	97
8	5 mol% 7	10 mol% TBD	31	6:1	90
9	5 mol% 7 /Mg(Ot-Bu) ₂	10 mol% TBD	78	7:1	97



*Isolated yields; [†]Determined by ¹H NMR spectroscopy; [‡]Determined by high-performance liquid chromatography (HPLC; chiral stationary phase); d.r., diastereomeric ratio; e.e., enantiomeric excess.

(entry 2) to 62% (entry 3) due to incomplete consumption of the hydrazone, a situation that could not be remedied with a single-catalyst approach.

We hypothesized that the overall process might benefit from increasing the electrophilicity of the hydrazone substrate and thus undertook an investigation combining carbene catalysts with Lewis acidic additives. This strategy to use nucleophilic carbenes as Lewis base catalysts in the presence of hard, oxophilic Lewis acids is a challenging goal with significant potential. If successful,

the ability to modulate the available electron density of the nucleophile and the electrophile simultaneously in two distinct catalytic cycles offers unique opportunities for new reaction discovery and development (Fig. 1). After an extensive survey of early transition and alkali metal salts, the use of 20 mol% magnesium di-*tert*-butoxide⁴¹ improved the reaction significantly both in terms of yield and selectivity (entry 4). Lanthanide trifluoromethanesulfonates (triflates) such as La(OSO₂CF₃)₃ completely inhibited homoenolate addition, even with an equivalent molar amount of base (for example, sodium *tert*-butoxide) to preclude the possibility of poisoning by trace acid. Transition-metal triflates, such as Cu(OSO₂CF₃)₂ and Zn(OSO₂CF₃)₂, were not catalysts alone, but the addition of base allowed a very modest progression toward the desired lactam over 24 h (10–30% conversion of hydrazone **1a**). The addition of titanium(IV) alkoxides resulted in very sluggish reactions, with only 40% conversion of hydrazone **1a** after 24 h. Magnesium (II) halides or triflates did not impede the progress of the reaction, but did not improve the results compared to experiments with only NHCs. Magnesium (II) di-*tert*-butoxide emerged as the optimal Lewis acid in terms of enhanced reaction rates, hydrazone conversion and isolated yields. In addition to optimizing the Lewis acid component, a survey of different chiral triazolium salts identified the 2,6-diethyl substituted azolium **7** as the superior catalyst in terms of enantioselectivity both with and without the Lewis acid additive (entries 6–9). The significant and beneficial impact of this cooperative catalysis protocol becomes evident when decreased loadings of magnesium(II) and azolium salts are used. For example, both azolium and Mg(II) sources can be lowered to 5 mol% without affecting the yield of the reaction or the high levels of enantioselectivity (entry 9). Decreasing the NHC loading without the Lewis acid has a deleterious effect on both yield and conversion (entry 8).

The proposed cooperative catalytic cycles are outlined in Fig. 2. The azolium precatalyst is deprotonated by the base (TBD) and the resultant carbene (NHC) adds to the α,β -unsaturated aldehyde. The deprotonation of the aldehyde proton generates the homoenolate equivalent (enediamine intermediate **II**), in which the electron density of the heterocyclic ring can then be translated to the β -carbon via the diene portion of the molecule. This nucleophilic species (**II**) then undergoes addition to the hydrazone, which is activated by chelation to magnesium(II) (intermediate **I**). However, no




Figure 2 | Cooperative catalysis between a Lewis acid and an NHC. **a**, Proposed catalytic cycles 1 ($\text{Mg}(\text{Ot-Bu})_2$) and 2 (NHC) meeting at the critical bond-forming step (box) **b**, Average k_{init} values plotted versus the concentration of $\text{Mg}(\text{Ot-Bu})_2$ to the negative first power. The experiments were performed with **1a**, cinnamaldehyde, catalyst D, TBD, phenyltrimethylsilane as an internal standard and a range of concentrations of $\text{Mg}(\text{Ot-Bu})_2$ in 1 ml THF-d₈. ^1H NMR spectroscopy (400 MHz) was used to monitor the concentration of the product (**2a**). The error bars represent 50% of the minimum and maximum values for k_{init} calculated. (See Supplementary Information for further details.)

significant change in chemical shift of any of the hydrazone signals was observed by ^1H nuclear magnetic resonance (NMR) upon the addition of $\text{Mg}(\text{Ot-Bu})_2$. Once the key carbon–carbon bond is formed, the NHC-catalytic cycle (cycle 2) is completed by the intramolecular acylation of the magnesium-bonded nitrogen (Mg-N) with concomitant ring closure. Finally, the magnesium (II) catalyst is regenerated by dissociation from the γ -lactam product to restart in catalytic cycle 1.


Our initial mechanistic investigations of the reactions allow for a cooperative catalysis interpretation. Preliminary kinetic studies were explored to begin elucidating the role of the $\text{Mg}(\text{Ot-Bu})_2$. Initial-rate kinetic studies were undertaken in triplicate at five concentrations of the magnesium salt to determine the kinetic order of the magnesium. When the k_{init} values for each experiment were calculated and plotted against the concentration of the magnesium (II) salt used, the best-fit model of the kinetic data reaction indicated an inverse first-order relation for $\text{Mg}(\text{Ot-Bu})_2$ (Fig. 2b). This result indicates one of two mechanistic possibilities for the magnesium salt. The first is that the Lewis basic NHC species is participating in reversible binding to the Lewis acidic magnesium, and the higher concentration of the magnesium salt is simply shifting the equilibrium toward the magnesium-bound azolium species, for which ^1H and ^{13}C NMR spectroscopy as well as X-ray crystal structure experiments to observe and/or isolate this complex have not yet

been successful. If this reversible magnesium–NHC interaction is taking place, then it is not likely that the two catalytic species are part of a frustrated Lewis acid–base system⁴². However, TBD is also Lewis basic and may undergo reversible binding to the magnesium, thus shifting the equilibrium between the active azolium/protonated TBD and protonated azolium/free-base TBD toward the inactive precatalyst.

With the optimized catalysis conditions identified, a variety of hydrazones with modified *N*-aryl groups were surveyed as potential substrates. Mild to moderate electron-withdrawing groups on the aromatic ring were well tolerated (for example, *para*-halogen). Without 5 mol% $\text{Mg}(\text{II})$, less than 50% conversion was observed when hydrazone substrates had electron-rich aryl substituents (for example, 4-MeO- C_6H_5). However, in this cooperative catalysis manifold, the yields with these substrates now regularly fall between 70 and 80% (**3ca**, **3ha**, Table 2). When varying the imine portion of the substrate, glyoxylate-derived hydrazones, including an oxazolidinone-containing substrate (**3ba**), reacted efficiently, but hydrazones from aryl aldehydes (for example, benzaldehyde) are still not electrophilic enough to undergo this annulation even with $\text{Mg}(\text{II})$ activation (not shown).

The aldehyde scope was surveyed and a variety of functionalities proved to be well tolerated. Weakly electron-withdrawing groups (Br, Cl) also performed well under the reaction conditions

Table 2 | Reaction scope.



*Isolated yields; ¹% e.e. determined by HPLC; ²d.r. determined by ¹H NMR spectroscopy; [§]10 mol% NHC/Mg(O-*t*Bu)₂, 15 mol% TBD.

(**3ad–g**). Electron-rich substrates were very well accommodated, including the 2-furyl-substituted enal which was problematic in earlier studies without the Lewis acid additive (**3ah**). Finally, an alkyl-substituted enal gave moderate yield and excellent selectivity (**3aj**, **3al**, **3am**), accommodating γ -branching and even silyl protecting groups.

The cyclic products from this new reaction can be easily converted into high-value pyroglutamic acid derivatives, thereby leveraging the utility of the overall process. For example, exposure of γ -lactam **3aa** to Raney nickel and H₂ in methanol promotes N–N bond scission in high yield (92%, Fig. 3). This 3-phenyl-pyroglutamic ester (**8**) is a known advanced intermediate for the synthesis of clausenamide, a natural product that promotes a protective effect for the liver against toxins such as carbon tetrachloride and thioacetamide and also upregulates cytochrome P450 enzymes^{37,43}. Alternatively, the amide of this unnatural amino ester can be selectively reduced to afford 3-phenyl proline derivatives⁴⁴. These compounds are known agonists for the melanocortin-4 receptor (MC4R), a G-protein coupled receptor involved in the regulation of appetite, and are potential treatments for anxiety

and depression³⁵. Various substituted prolines have been used to impart desirable structural attributes and metabolic stability in peptides and proteins, and the enantioselective synthesis of compounds such as **9** should fuel further studies in this area⁴⁵. Finally, these pyrrolidines also have the necessary characteristics for secondary amine/iminium catalysis, and new proline-derived catalysts can be generated using this route^{46,47}.

In summary, a new formal [3 + 2] reaction combining α,β -unsaturated aldehydes and hydrazones has been discovered using a novel cooperative catalysis concept. The integration of two distinct catalytic cycles involving both Lewis basic NHCs and Lewis acidic magnesium(II) salts facilitates the rapid and controlled production of γ -lactams in high yields and with exceptional levels of diastereo- and enantioselectivity. Each mode of catalysis, both carbene and Lewis acid, enhances the reactivity of the nucleophilic and electrophilic substrates, respectively. This use of nucleophilic carbene catalysis in cooperation with Lewis acid activation highlights an under-explored area of catalysis with significant potential. These initial studies demonstrate the broad potential utility and feasibility of combining Lewis basic carbenes with Lewis acids and should lead




Figure 3 | N–N bond cleavage and further functionalization. The N–N bond can be cleaved by hydrogenolysis over Raney nickel. Compound **8** can be converted to the drug clausenamide, which is used in the treatment of viral hepatitis in 13 steps. Reduction of the amide with borane provides access to a variety of proline derivatives useful in protein studies as scaffolds for organocatalysts.

to many more powerful and highly efficient bond-forming reactions not currently available using either reagent singly. Cooperative carbene catalysis will continue to grow and facilitate new strategies to access important target molecules for use in chemical biology, catalysis and medicine.

Methods

Hydrazone **1** (0.273 mmol), catalyst **7** (5 mol%) and $\text{Mg}(\text{O}i\text{-Bu})_2$ (5 mol%) were combined in a 2 dram oven-dried vial and capped with a Teflon/silicon septum inside a nitrogen drybox. The vial was removed from the drybox and THF (0.25 M) added, then the vial was heated with moderate stirring in an aluminium heating block to 60 °C for 15–20 min. The white suspension turned clear and yellow over the course of the initial heating. Unsaturated aldehyde **2** (1.5 equiv.) and then TBD (10 mol%) were immediately added. The reaction was allowed to continue at 60 °C and was monitored by thin layer chromatography (5% MeOH/ CHCl_3). Upon completion (24 h), the reaction was diluted with CH_2Cl_2 (15 ml) and washed with a 1:1 mixture of saturated aqueous NH_4Cl and water (10 ml). The aqueous layer was back-extracted CH_2Cl_2 (2 × 15 ml) and the combined organic layers were dried over anhydrous sodium sulfate and concentrated. The brown oily residue was purified by flash column chromatography on a Biotage SP-1 chromatography system using a 12–100% ethyl acetate/hexanes gradient to yield the desired product **3**.

Received 24 November 2009; accepted 21 May 2010;
published online 18 July 2010

References

- Kanai, M., Kato, N., Ichikawa, E. & Shibasaki, M. Power of cooperativity: Lewis acid–Lewis base bifunctional asymmetric catalysis. *Synlett* 1491–1508 (2005).
- Breslow, R. Bifunctional acid–base catalysis by imidazole groups in enzyme mimics. *J. Mol. Catal.* **91**, 161–174 (1994).
- Breslow, R. Bifunctional binding and catalysis. *Supramol. Chem.* **1**, 111–118 (1993).
- Lee, J. K., Kung, M. C. & Kung, H. H. Cooperative catalysis: a new development in heterogeneous catalysis. *Top. Catal.* **49**, 136–144 (2008).
- Miyabe, H. & Takemoto, Y. Discovery and application of asymmetric reaction by multi-functional thioureas. *Bull. Chem. Soc. Jpn* **81**, 785–795 (2008).
- Paull, D. H., Abraham, C. J., Scerba, M. T., Alden-Danforth, E. & Lectka, T. Bifunctional asymmetric catalysis: cooperative Lewis acid/base systems. *Acc. Chem. Res.* **41**, 655–663 (2008).
- Frank, R. A. W., Titman, C. M., Pratap, J. V., Luisi, B. F. & Perham, R. N. A molecular switch and proton wire synchronize the active sites in thiamine enzymes. *Science* **306**, 872–876 (2004).
- Jordan, F. & Patel, M. S. *Thiamine: Catalytic Mechanisms in Normal and Disease States* (Marcel Dekker, 2004).
- Denmark, S. E. & Beutner, G. L. Lewis base catalysis in organic synthesis. *Angew. Chem. Int. Ed.* **47**, 1560–1638 (2008).
- Enders, D., Niemeier, O. & Henseler, A. Organocatalysis by *N*-heterocyclic carbenes. *Chem. Rev.* **107**, 5606–5655 (2007).

- Phillips, E. M., Chan, A. & Scheidt, K. A. Discovering new reactions with *N*-heterocyclic carbenes. *Aldrichimica Acta* **42**, 55–66 (2009).
- Sheehan, J. C. & Hara, T. Asymmetric thiazolium salt catalysis of the benzoin condensation. *J. Org. Chem.* **39**, 1196–1199 (1974).
- Stetter, H., Raemsch, R. Y. & Kuhlmann, H. The preparative use of thiazolium salt-catalyzed acyloin and benzoin formation, i. Preparation of simple acyloins and benzoin. *Synthesis* 733–735 (1976).
- Breslow, R. & Schmuck, C. The mechanism of thiazolium catalysis. *Tetrahedron Lett.* **37**, 8241–8242 (1996).
- Enders, D. & Kallfass, U. An efficient nucleophilic carbene catalyst for the asymmetric benzoin condensation. *Angew. Chem. Int. Ed.* **41**, 1743–1745 (2002).
- Kerr, M. S., de Alaniz, J. R. & Rovis, T. A highly enantioselective catalytic intramolecular Stetter reaction. *J. Am. Chem. Soc.* **124**, 10298–10299 (2002).
- Mattson, A. E., Bharadwaj, A. R. & Scheidt, K. A. The thiazolium-catalyzed Stetter reaction: conjugate addition of acylsilanes to unsaturated esters and ketones. *J. Am. Chem. Soc.* **126**, 2314–2315 (2004).
- Burstein, C. & Glorius, F. Organocatalyzed conjugate umpolung of α,β -unsaturated aldehydes for the synthesis of γ -butyrolactones. *Angew. Chem. Int. Ed.* **43**, 6205–6208 (2004).
- Chan, A. & Scheidt, K. A. Conversion of α,β -unsaturated aldehydes into saturated esters: an umpolung reaction catalyzed by nucleophilic carbenes. *Org. Lett.* **7**, 905–908 (2005).
- Sohn, S. S., Rosen, E. L. & Bode, J. W. *N*-heterocyclic carbene-catalyzed generation of homoenolates: γ -butyrolactones by direct annulations of enals and aldehydes. *J. Am. Chem. Soc.* **126**, 14370–14371 (2004).
- Nair, V., Vellalath, S., Poonoth, M. & Suresh, E. *N*-heterocyclic carbene-catalyzed reaction of chalcones and enals via homoenolate: an efficient synthesis of 1,3,4-trisubstituted cyclopentenones. *J. Am. Chem. Soc.* **128**, 8736–8737 (2006).
- Chan, A. & Scheidt, K. A. Highly stereoselective formal [3 + 3] cycloaddition of enals and azomethine imines catalyzed by *N*-heterocyclic carbenes. *J. Am. Chem. Soc.* **129**, 5334–5335 (2007).
- Phillips, E. M., Reynolds, T. E. & Scheidt, K. A. Highly diastereo- and enantioselective additions of homoenolates to nitrones catalyzed by *N*-heterocyclic carbenes. *J. Am. Chem. Soc.* **130**, 2416–2417 (2008).
- Phillips, E. M., Wadamoto, M., Chan, A. & Scheidt, K. A. A highly enantioselective intramolecular Michael reaction catalyzed by *N*-heterocyclic carbenes. *Angew. Chem. Int. Ed.* **46**, 3107–3110 (2007).
- Wadamoto, M., Phillips, E. M., Reynolds, T. E. & Scheidt, K. A. Enantioselective synthesis of α,α -disubstituted cyclopentenones by an *N*-heterocyclic carbene-catalyzed desymmetrization of 1,3-diketones. *J. Am. Chem. Soc.* **129**, 10098–10099 (2007).
- Reynolds, T. E., Stern, C. A. & Scheidt, K. A. *N*-heterocyclic carbene-initiated α -acylvinyl anion reactivity: Additions of α -hydroxypropargylsilanes to aldehydes. *Org. Lett.* **9**, 2581–2584 (2007).
- Burgess, K. & Perry, M. C. Chiral *N*-heterocyclic carbene–transition metal complexes in asymmetric catalysis. *Tetrahedron: Asymmetry* **14**, 951–961 (2003).
- Brown, M. K., May, T. L., Baxter, C. A. & Hoveyda, A. H. All-carbon quaternary stereogenic centers by enantioselective Cu-catalyzed conjugate additions promoted by a chiral *N*-heterocyclic carbene. *Angew. Chem. Int. Ed.* **46**, 1097–1100 (2007).
- Lee, Y., Li, B. & Hoveyda, A. H. Stereogenic-at-metal Zn- and Al-based *N*-heterocyclic carbene (NHC) complexes as bifunctional catalysts in Cu-free enantioselective allylic alkylations. *J. Am. Chem. Soc.* **131**, 11625–11633 (2009).
- Nolan, S. P. *N-Heterocyclic Carbenes in Synthesis* (Wiley-VCH, 2006).
- Diez-Gonzalez, S., Marion, N. & Nolan, S. P. *N*-heterocyclic carbenes in late transition metal catalysis. *Chem. Rev.* **109**, 3612–3676 (2009).
- Lin, J. C. Y. *et al.* Coinage metal-*N*-heterocyclic carbene complexes. *Chem. Rev.* **109**, 3561–3598 (2009).
- Glorius, F. E. *N-Heterocyclic Carbenes in Transition Metal Catalysis* (Springer-Verlag, 2007).
- Lebeuf, R., Hirano, K. & Glorius, F. Palladium-catalyzed C-allylation of benzoin and an NHC-catalyzed three component coupling derived thereof: compatibility of NHC- and Pd-catalysts. *Org. Lett.* **10**, 4243–4246 (2008).
- Tran, J. A. *et al.* Design and synthesis of 3-arylpyrrolidine-2-carboxamide derivatives as melanocortin-4 receptor ligands. *Bioorg. Med. Chem. Lett.* **18**, 1931–1938 (2008).
- Silverman, R. B. & Nanavati, S. M. Selective-inhibition of gamma-aminobutyric acid aminotransferase by (3R,4R),(3S,4S)-4-amino-5-fluoro-3-phenylpentanoic acid and (3R,4S),(3S,4R)-4-amino-5-fluoro-3-phenylpentanoic acids. *J. Med. Chem.* **33**, 931–936 (1990).
- Hartwig, W. & Born, L. Diastereoselective and enantioselective total synthesis of the hepatoprotective agent clausenamide. *J. Org. Chem.* **52**, 4352–4358 (1987).
- He, M. & Bode, J. W. Catalytic synthesis of gamma-lactams via direct annulations of enals and *N*-sulfonylimines. *Org. Lett.* **7**, 3131–3134 (2005).
- Sugiura, M. & Kobayashi, S. *N*-acylhydrazones as versatile electrophiles for the synthesis of nitrogen-containing compounds. *Angew. Chem. Int. Ed.* **44**, 5176–5186 (2005).

40. Kaljurand, I. *et al.* Extension of the self-consistent spectrophotometric basicity scale in acetonitrile to a full span of 28 pK_{a} units: unification of different basicity scales. *J. Org. Chem.* **70**, 1019–1028 (2005).
41. Beach, R. G. & Ashby, E. C. Synthesis and characterization of dialkyl(aryl)aminomagnesium hydrides and alkoxy(aryloxy)magnesium hydrides. *Inorg. Chem.* **10**, 906–910 (1971).
42. Stephan, D. W. & Erker, G. Frustrated Lewis pairs: metal-free hydrogen activation and more. *Angew. Chem. Int. Ed.* **49**, 46–76 (2010).
43. Tang, K. & Zhang, J. T. The effects of (–)-clausenamide on functional recovery in transient focal cerebral ischemia. *Neurol. Res.* **24**, 473–478 (2002).
44. Haldar, P. & Ray, J. K. Chemoselective reduction of a lactam carbonyl group in the presence of a gem-dicarboxylate by sodium borohydride and iodine: a facile entry to *N*-aryl trisubstituted pyrroles. *Tetrahedron Lett.* **44**, 8229–8231 (2003).
45. Arnold, U. *et al.* Protein prosthesis: a nonnatural residue accelerates folding and increases stability. *J. Am. Chem. Soc.* **125**, 7500–7501 (2003).
46. Mukherjee, S., Yang, J. W., Hoffmann, S. & List, B. Asymmetric enamine catalysis. *Chem. Rev.* **107**, 5471–5569 (2007).
47. MacMillan, D. W. C. The advent and development of organocatalysis. *Nature* **455**, 304–308 (2008).

Acknowledgements

Support for this work was generously provided by National Institute of General Medical Sciences (RO1 GM73072), GlaxoSmithKline, AstraZeneca and the Alfred P. Sloan Foundation. B.C.D. thanks the Fonds Quebecois de la Recherche sur la Nature et les Technologies for a postdoctoral fellowship. Funding for the Northwestern University Integrated Molecular Structure Education and Research Center (IMSERC) has been furnished in part by the National Science Foundation (CHE-9871268). T. Reynolds and J. Roberts provided X-ray crystallography support.

Author contributions

K.A.S. conceived the idea and wrote the manuscript. D.E.A.R., B.C.-D. and D.H. performed the experiments. All the authors analysed the data. K.A.S., D.E.A.R. and B.C.-D. contributed to discussions and edited the manuscript.

Additional information

The authors declare no competing financial interests. Supplementary information and chemical compound information accompany this paper at www.nature.com/naturechemistry. Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>. Correspondence and requests for materials should be addressed to K.A.S.