

Synthesis of *trans*-*N,N,N*-triethyl-2-[4-(2-phenylethenyl)phenoxy]-ethan ammonium iodide

Mohammed I. Al-Hassan

Department of Chemistry, College of Science, King Saud University, P.O. Box 2455, Riyadh-11451 (Saudi Arabia)


(Received November 8th, 1989)

Abstract


The antispasmodic olefin *trans*-*N,N,N*-triethyl-2-[4-(2-phenylethenyl)phenoxy]-ethan ammonium iodide has been made by hydroalumination of a trimethylsilylalkyne followed by palladium-catalyzed cross-coupling of the vinylalane with the appropriate aryl bromide.

trans-*N,N,N*-Triethyl-2-[4-(2-phenylethenyl)phenoxy]-ethan ammonium iodide (1) is a biologically active disubstituted olefin that is used as an antispasmodic ganglionic blocker [1,2].

We describe here the synthesis of compound **1** by hydroalumination of a trimethylsilylalkyne followed by (i) palladium-catalyzed cross-coupling [3,4] of the vinylalane with the appropriate aryl bromide (ii) removal of the trimethylsilyl group and (iii) treatment of the desilylated olefin with ethyl iodide.



Thus trimethylsilylalkyne **2** or **3** was treated with diisobutylaluminium hydride in refluxing ether for 5 h to give the corresponding vinylalane [5] which was cross-coupled with bromobenzene [6] in the presence of catalytic amount (0.5 equiv.) of tetrakis-(triphenylphosphine)palladium [3,4] to give the vinylsilane **4** or **5**. The compounds **4** and **5** were transformed into the desired product (**1**) without further purification by protodesilylation with hydroiodic acid [7] followed by treatment of the desilylated species with ethyl iodide in refluxing tetrahydrofuran for 1 h. The desired product **1** was obtained in good overall yield after recrystallization from ethyl acetate/benzene.



Experimental

Melting points are uncorrected. GLC analysis was performed on PYE Unicam series 304 chromatograph with an OV1 on glass column.

Diisobutylaluminum hydride (Fluka) was used as a 1 M solution in hexane. Reaction mixtures were stirred magnetically under nitrogen. Oven-dried (160 °C) glassware was used. Tetrahydrofuran, diethyl ether, and benzene were distilled from sodium/benzophenone.

Preparation of trans-N,N,N-triethyl-2-[4-(2-phenylethenyl)-phenoxy]ethanamonium iodide (1). To 6 mmol of the trimethylsilylalkyne, **2** or **3**, in 12 ml of ether at room temperature were added dropwise 6.5 ml of 1 M diisobutylaluminum hydride in hexane (6.5 mmol) during 0.5 h. The mixture was refluxed for 5 h and the resulting vinylsilane solution was then treated with a mixture of 6 mmol of the relevant aryl bromide and 0.3 mmol (0.05 equiv.) of tetrakis(triphenylphosphine) palladium in 18 ml of dry tetrahydrofuran. The mixture was refluxed for 24 h and then treated with an excess of the water. The usual work-up and evaporation of solvents left the crude vinylsilane, **4** or **5**, which was refluxed with hydroiodic acid in 15 ml of benzene. Basic work-up and evaporation of solvents left a crude product which was refluxed with 6 mmol of ethyl iodide in 15 ml of tetrahydrofuran for 1 h. Work-up and evaporation of solvents left a residue, which was recrystallized from ethyl acetate/benzene to give 3.7 mmol (62% overall yield) from compound **2** or 4 mmol (67% overall yield) from compound **3** of the desired compound (**1**) as a white crystalline solid. GLC showed only one peak.

Acknowledgement

This research (Chem/1408/12) was supported by the Research Center, College of Science, King Saud University, Riyadh, Saudi Arabia.

References

- 1 The Merck Index, 9th edit., Merck & Co., Inc., Rahway, NJ, 1976.
- 2 E. Cavallini, *Farmaco Ed. Sci.*, 8 (1953) 317.
- 3 M.I. Al-Hassan, *Synth. Commun.*, 17 (1987) 1787.
- 4 M.I. Al-Hassan, *J. Organomet. Chem.*, 321 (1987) 119.
- 5 M.I. Al-Hassan, *Synth. Commun.*, 16 (1986) 353.
- 6 R. Oliver and D.R.M. Walton, *Tetrahedron Lett.*, 51 (1972) 5209.
- 7 T.H. Chan and I. Fleming, *Synthesis*, (1979) 761.