View Article Online / Journal Homepage / Table of Contents for this issue

Palladium-catalysed 1,4-Arylation/Alkylation of Buta-1,3-diene

Mitsunari Uno, Terumasa Takahashi, and Shigetoshi Takahashi*

The Institute of Scientific and Industrial Research, Osaka University, Ibaraki, Osaka 567, Japan

Palladium-catalysed reactions of butadiene with halogeno-arenes and stabilized anions give arylation/alkylation products of the diene in which two carbon–carbon bonds are formed from three components in one catalytic cycle.

Many reports have appeared on organic syntheses using palladium-catalysed reactions in which a π -allylpalladium species acts as a key intermediate.¹ Recently we have shown that stabilized anions such as $-CH(CN)_2$ and $-CH(CN)CO_2R$ react smoothly with some halogeno-arenes in the presence of a palladium catalyst to give the coupled products (3) (equation 1).² The mechanism of this reaction may reasonably be explained by postulating the initial formation of a Pd–Ar intermediate, which may be able to undergo insertion of dienes to afford π -allylpalladium species. We now report that a palladium–phosphine complex catalyses the 1,4-arylation/ alkylation of buta-1,3-diene with halogeno-arenes and malononitrile or cyanoacetates in which two carbon–carbon bonds are formed from three components in one catalytic cycle (equation 2). Catalytic arylation/alkylation reactions of 1,3dienes have not been reported previously.³

Ar-X +
$$^{-}CH(CN)E \xrightarrow{Pd^{0} \text{ catalyst}} Ar-CH(CN)E + X^{-}(1)$$

(1) (2) (3)

Thus, in the presence of dichlorobis(triphenylphosphine)palladium catalyst (0.3 mmol) a mixture of iodobenzene (30 mmol) and malononitrile anion (2: E = CN), generated *in situ* from malononitrile (10 mmol) and sodium hydride, was allowed to react with a large excess of butadiene (*ca.* 120



mmol) in tetrahydrofuran at 70 °C for 67 h. After quenching with dilute hydrochloric acid, the product was extracted with diethyl ether, and then purified by column chromatography on silica gel using hexane and chloroform as eluents. A 1,4arylation/cyanoalkylation product of butadiene, *trans*-5,5-dicyano-1,5-diphenylpent-2-ene (**4a**) was obtained in 62% yield (based on malononitrile employed) which was identified by spectroscopic analyses.† *p*-Iodotoluene similarly reacted to give the corresponding product, but *p*-chloroiodobenzene required higher temperature and longer reaction time. The reactivity of the halogeno-arenes was in the same order as observed previously in the dicyanomethylation of halogenoarenes, I > Br \gg Cl. Cyanoacetates (**2**, E = CO₂R) similarly reacted to give (**4d**—**f**) in about 40% yield (Table 1).

† $C_6H_5CH_2CH=CHCH_2C(C_6H_5)(CN)_2$ (4a): m/z 272 (M^+); ¹H n.m.r. (CDCl₃, 360 MHz): δ 7.1—7.6 (10H, m), 5.88 (1H, dt, J 15, 7 Hz), 5.51 (1H, dt, J 15, 7 Hz), 3.41 (2H, d, J 7 Hz), and 2.92 (2H, d, J 7 Hz).

Table 1. 1,4-Arylation/alkylation of butadiene.

		Conditions				
		CH ₂ (CN)E			Yield	
Run	Ar–X	E	Temp./°C	Time/h	Product	(%)
1	Ph-I	CN	70	67	(4 a)	62
2	Ph–Br	CN	70	48	(4a)	10 ^ь
3	p-MeC ₆ H ₄ -I	CN	70	48	(4b)	46
4	p-ClC ₆ H ₄ -I	CN	80	120	(4 c)	36
5	Ph-I	CO ₂ Et	70	72	(4d)	40
6	PhI	CO_2Me	70	45	(4e)	42
7	p-MeC ₆ H ₄ -I	CO ₂ Me	70	45	(4f)	47

^a Isolated yield based on $CH_2(CN)E$. ^b Catalyst, $(Ph_2PCH_2CH_2PPh_2)$ -PdCl₂; solvent, dimethoxyethane; the yield was determined by g.l.c.

Under the same conditions the use of phenylmalononitrile instead of malononitrile afforded (4a) in 70% yield, indicating that the reaction step incorporating the diene proceeds after the dicyanoalkylation of halogeno-arenes, *i.e.*, initial formation of arylmalononitrile (3) from the reaction between malononitrile and halogeno-arenes (equation 1), and then attack of the anion on the π -allylpalladium intermediate yields the products (4).

Received, 18th December 1986; Com. 1794

References

- J. Tsuji, 'Organic Synthesis with Palladium Compounds', Springer, Berlin, 1980; B. M. Trost and T. R. Verhoeven, 'Comprehensive Organometallic Chemistry', Pergamon, Oxford, 1982, Vol. 5, p. 799; R. F. Heck, 'Palladium Reagents in Organic Syntheses,' Academic Press, New York, 1985.
- 2 M. Uno, K. Seto, and S. Takahashi, J. Chem. Soc., Chem. Commun., 1984, 932; M. Uno, K. Seto, W. Ueda, M. Masuda, and S. Takahashi, Synthesis, 1985, 506.
- 3 Related work: B. A. Patel, J. E. Dickerson, and R. F. Heck, J. Org. Chem., 1978, 43, 5018; L. S. Hegedus and Y. Inoue, J. Am. Chem. Soc., 1982, 104, 4917; I. Shimizu and J. Tsuji, Chem. Lett., 1984, 233; M. Ahmar, B. Cazes, and J. Gore, Tetrahedron Lett., 1984, 25, 4505.