


This article was downloaded by: [National Sun Yat-Sen University]

On: 28 December 2014, At: 19:59

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office:

Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lcar20>

### Synthesis of 2-Deoxy-2-thiophenylglucosyl Azides through 1,2 Thio Migration of Thiophenyl Mannosides

Carmen Maieranu<sup>a</sup>, Akira Kanai<sup>a</sup>, Jean-Marc Weibel<sup>a</sup> & Patrick Pale<sup>a</sup>

<sup>a</sup> Laboratoire de synthèse et réactivité organique, associé au CNRS, Institut Le Bel, Université L. Pasteur, Strasbourg, France

Published online: 02 Feb 2007.

To cite this article: Carmen Maieranu, Akira Kanai, Jean-Marc Weibel & Patrick Pale (2005) Synthesis of 2-Deoxy-2-thiophenylglucosyl Azides through 1,2 Thio Migration of Thiophenyl Mannosides, Journal of Carbohydrate Chemistry, 24:8-9, 831-842, DOI: [10.1080/07328300500409150](https://doi.org/10.1080/07328300500409150)

To link to this article: <http://dx.doi.org/10.1080/07328300500409150>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

# Synthesis of 2-Deoxy-2-thiophenylglucosyl Azides through 1,2 Thio Migration of Thiophenyl Mannosides

Carmen Maiereanu, Akira Kanai\*, Jean-Marc Weibel, and Patrick Pale

Laboratoire de synthèse et réactivité organique, associé au CNRS, Institut Le Bel, Université L. Pasteur, Strasbourg, France

2-Deoxy-2-thio glucosyl azides bearing or not bearing an azido group at the 6-position can be conveniently obtained through a 1,2-thio migration from thiophenyl mannoside derivatives.

**Keywords** Glucosyl azides, Thiophenyl mannosides, 1,2-Thio migration


## INTRODUCTION

Carbohydrates play crucial roles in important biological recognition processes, including bacterial and viral infections, cell adhesion in inflammation and metastasis, differentiation, development, regulation, and other intercellular communication and signal transduction events.<sup>[1,2]</sup> Since natural oligosaccharides are easily degraded in vivo, development of biologically stable oligosaccharide analogs is very important for studying such recognition processes and for the elaboration of carbohydrate-based therapeutics and vaccines.<sup>[3]</sup> In this context, we are investigating 6-amino-6-deoxyglycosides, glycosyl amines, and the corresponding deoxy derivatives as building blocks for some oligosaccharide mimics (Fig. 1).<sup>[4,5]</sup>

Received August 2005; accepted September 2005.

\*On leave from the Tokyo University of Science (RIKADAI), Faculty of Pharmaceutical Sciences.

Address correspondence to Patrick Pale, Laboratoire de synthèse et réactivité organique, associé au CNRS, Institut Le Bel, Université L. Pasteur, Strasbourg 67000, France. Tel./Fax: +33 390 241 517; E-mail: ppale@chimie.u-strasbg.fr




**Figure 1:** Oligosaccharides mimics based on glycosyl amines, 6-amino-6 deoxyglycosyl amines and their corresponding 2-thio and 2-deoxy analogs.

In order to have convenient access to this family of compounds, we realized that 2-deoxy-2-thio glycosyl azides, with or without an azido group at the 6 position, would offer a key entry to several series of derivatives. The azido group can indeed be reduced to the amine, and the thio group can also be reduced leading to the 2-deoxy series. A literature survey revealed that 6-amino-2,6-dideoxy-2-thioglycosyl azides are unknown and that even 2-deoxy-2-thioglycosyl azides surprisingly are almost unknown. A single paper mentioned 2-deoxy-2-thioglycosyl azides in multicomponent reactions leading to carbohydrate combinatorial libraries;<sup>[6]</sup> they were obtained as mixtures through electrophilic addition of dibenzyl disulfide-antimony pentachloride to glucals in the presence of trimethylsilyl azide and alcohols.

We thus designed and described here an alternative and more convenient route to 2-deoxy-2-thio glycosyl azides bearing or not bearing an azido group at the 6-position through a 1,2-migration of thio mannoside derivatives (Sch. 1).

The rearrangement of thio derivatives through thiiranium intermediates is a well-known process often used in organic chemistry.<sup>[7]</sup> This process has been extended in carbohydrate chemistry and applied to the synthesis of 2-thioglycosides,<sup>[8]</sup> 2-deoxyglycosides,<sup>[9–11]</sup> sugar 2-*C*-sulfonic acids,<sup>[12]</sup> and 1,2-thio-linked disaccharides.<sup>[13]</sup> It has also been used in the total synthesis of apoptolidin.<sup>[14]</sup> This rearrangement was also detected in various works as an undesirable side reaction.<sup>[15–18]</sup>

Based on these precedents, we anticipated that  $\alpha$ -thio mannosides bearing an electron-withdrawing group at the 2-position should have the right geometrical and electronic requirements for a facile 1-2-thio migration. In the presence of an azide source, the thiiranium intermediate, which should be formed, might be opened, leading to 2-deoxy-2-thioglycosyl azides (Sch. 1).




**Scheme 1:** Anticipated synthesis of 2-deoxy-2-thioglycosyl azides by 1,2-thio migration of thio mannosides (R = OPG, N<sub>3</sub>).

## RESULTS AND DISCUSSION

To validate this route, we started from the known<sup>[19]</sup>  $\alpha$ -D-thiophenyl mannoside protected by the Ley acetal<sup>[20]</sup> at the 3, 4 position (**1** in Sch. 2). Due to this selective protecting group, the 2, 6-positions of the sugar remained free but are easily distinguishable, the 6 being a primary hydroxyl group. Indeed, selective silylation<sup>[21]</sup> with tri-*isopropylsilyl* chloride in the presence of imidazole provided exclusively the 6-silylated compound **2**, which was then tosylated at the 2-position with the Bouzide's method,<sup>[22]</sup> giving **3** in excellent overall yields. Alternatively, selective tosylation at the 6 position can be achieved using Yoshida's method<sup>[23]</sup> (**4** in Sch. 2). The resulting primary tosylate was then displaced by sodium azide, leading to the azido compound **5**, which was then mesylated and tosylated (**6a–b**), thus providing an entry to the 6-amino-6-deoxyglucosyl series.

The tosylated thiophenyl  $\alpha$ -D-mannoside **3** was used as a model to find the right conditions for the expected 1,2-thio migration. Treatment of **3** with sodium azide in polar solvents such as DMF, MeCN, THF, etc., at various temperatures led mostly to the recovery of the starting material if the temperature did not reach 100°C. DMF proved to be the best solvent, but its effect seemed mainly correlated with its boiling point. Indeed, nothing happened after 24 h at 50°C and only 15% of starting material was converted at 80°C (Table 1; entries 1–2), but at 100°C, 50% of the expected product **7** was formed (entry 3). A better yield was obtained at 130°C but decomposition started occurring (entry 4). The structure of **7** was easily deduced from changes in NMR spectra. As expected, the signal corresponding to H<sub>2</sub> was



**Scheme 2:** Synthesis of the starting materials.

**Table 1:** Formation of glucosyl azides upon treatment of  $\alpha$ -thiophenyl mannosides by NaN<sub>3</sub> in DMF.

Entry	Glycosides	Products	Temperature (°C)	Time (h)	Ratio $\alpha:\beta$ <sup>a</sup>	Yield (%) <sup>b</sup>
1			50	24	1:2.5	0
2			80	15	1:2.5	11
3			100	15	1:2.5	50
4			130	15	1:2.5	64
5			80	24	—	0 <sup>c</sup>
6			130	15	1:2	46
7			130	15	2:1	45
8			130	24	2.5:1	76
9			130	15	1.2:1	62

<sup>a</sup>The ratio was derived from integration of the H<sup>1</sup> NMR anomeric signals of the crude mixture.<sup>b</sup>Yields of pure isolated product.<sup>c</sup>Starting material recovered.

shifted from 4.88 ppm in **3** to 3.27 ppm in **7 $\alpha$**  (2.97 in **7 $\beta$** ) and its multiplicity changed from a massif in **3** to a doublet of doublet in **7**, clearly reflecting profound modifications in the atomic environment of this proton and in its stereochemistry. Similarly, the signal of the anomeric proton H<sub>1</sub> was shielded, especially for the  $\beta$ -anomer of **7** (5.55 ppm in **3**, 5.38 in **7 $\alpha$** , and 4.58 ppm in **7 $\beta$** ). The multiplicity was also different: Appearing as a broad singlet in **3**, the anomeric proton in **7** is a doublet with typical constant either for axial-equatorial coupling in **7 $\alpha$**  ( $J = 4.0$  Hz) or for axial-axial coupling in **7 $\beta$**  ( $J = 9.5$  Hz).


It is worth noting that the  $\alpha/\beta$  ratio did not change whatever the conditions in this series of experiments, and that it was in favor of the  $\beta$  isomer as expected from nucleophilic opening of a thiiranium intermediate (see Sch. 1). Further control experiments showed that this ratio did not change either

upon prolonged heating. These results are thus in agreement with the intermediate formation of a thiiranium species.

During preliminary experiments to form **4** using standard tosylation procedures, we obtained mixtures of products, among which the mannoside monotosylated at position 2 (**8**) was isolated (<15%). This analog of **3** bearing a free hydroxyl group at position 6 (**8**) was also submitted to the above-mentioned conditions (Table 1, entries 5–6). Although disaccharide formation could be a competitive process via a nucleophilic attack of the free 6-alcohol to the thiiranium intermediate, the corresponding thioglucosyl azide **9** was cleanly obtained but in moderate yields. Here again, a higher temperature was required (entry 6 vs. 5) and the  $\alpha/\beta$  ratio observed was similar to the one obtained from the 6-protected **3** (entry 6 vs. 3, 4). Interestingly enough, no disaccharide can be detected, reflecting the better nucleophilicity of the azide ion over the hydroxyl group. Compound **9** exhibited NMR data similar to **7** with  $H_1$  appearing as a doublet at 5.38 ppm ( $J = 4.0$  Hz) in **7 $\alpha$**  and 4.61 ppm ( $J = 9.6$  Hz) in **7 $\beta$**  and with  $H_2$  as a doublet of doublet at 3.30 ppm ( $J = 4.0, 11.3$  Hz) in **7 $\alpha$**  and 2.98 ppm ( $J = 9.6, 11.2$  Hz) in **7 $\beta$** .

With the conditions in hand for the expected 1,2-thio migration and azidation, we started investigating the azido mannosides **6a–b**. Both gave the expected rearranged product **10** in good yields (Table 1, entries 7–9). The azido mesylated compound **6a** proved to be less reactive than its tosylated counterpart **6b**. Indeed, heating **6a** at 130°C for 1 day was required to get a satisfactory yield (entry 8 vs. 7), while heating overnight was enough for **6b** (entry 9). The new product formed **10** exhibited NMR spectra very similar to **7** and **9** and was also a mixture of anomers ( $H_1$ : doublet at 5.41 ppm ( $J = 4.2$  Hz) in **10 $\alpha$**  and 4.59 ppm ( $J = 9.6$  Hz) in **10 $\beta$** ;  $H_2$ : doublet of doublet at 3.29 ppm ( $J = 4.2, 11.5$  Hz) in **10 $\alpha$**  and 3.00 ppm ( $J = 9.6, 10.8$  Hz) in **10 $\beta$** ).

Surprisingly, the anomeric ratio of the azidoglucosyl azide so formed **10** was in favor of the  $\alpha$  isomer when starting from the mesyl derivative **6a**, but did not show any significant preference for one anomer or the other when starting from the tosyl derivatives **6b**. Such differences obviously reflect some isomerization of the thiiranium intermediate to a thia-carbenium intermediate, but the role of the leaving group in this process is unclear (Sch. 3).



**Scheme 3:** Isomerization of the thiiranium intermediate, leading to an  $\alpha/\beta$  mixture of products.

## CONCLUSION

This work demonstrated that concomitant 1,2-thio migration and azidation of  $\alpha$ -thiophenyl mannosides can be achieved and offer a convenient synthesis of various glucosyl azides, substituted or not. The compounds so obtained could then be reduced to the corresponding glycosyl amines and to their 2-deoxy derivatives. Such compounds may find broad applications as building blocks and tools in glycoscience. Progress in this area is under way.

## EXPERIMENTAL

### General

Reactions were monitored by TLC on silica gel MERCK DC Alufolien Kieselgel 60 PF<sub>254</sub> with detection by UV absorption (254 nm). All moisture-sensitive reactions were carried out under an atmosphere of argon in oven-dried glassware. Solvents were freshly distilled; THF was distilled from Na/benzophenone; C<sub>6</sub>H<sub>6</sub>, MeCN, and CH<sub>2</sub>Cl<sub>2</sub> were distilled from CaH<sub>2</sub>. Column chromatographies were performed with Merck silica gel (0.040–0.063 mesh). Yields were based on isolated products. <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on a Bruker AC-300 spectrometer. For <sup>1</sup>H NMR, TMS was used as internal standard; for <sup>13</sup>C NMR, the solvent peak at 77.00 ppm (CDCl<sub>3</sub>) was used. IR spectra were recorded on a spectrafiler IR<sup>TM</sup> Plus MIDAC spectrophotometer. Mass Spectra were recorded on a Bruker Daltonis Data Analysis 3.1 mass spectrometer. Specific rotations were measured at 589 nm with an ADP 220 Bellingham-Stanley polarimeter or a Perkin Elmer 241 polarimeter. Melting points were determined on a Bibby Sterilin Stuart Scientific melting point apparatus and are uncorrected.

### Phenyl 6-O-triisopropylsilyl-3,4-O-(2',3'-dimethoxybutane-2',3'-diyl)-1-thio- $\alpha$ -D-mannopyranoside (2)

A solution of diol **1**<sup>[19]</sup> (0.04 g, 0.104 mmol), triisopropylsilylchloride (0.034 mL, 0.156 mmol), and imidazole (0.011 g, 0.156 mmol) in DMF (1 mL) was stirred for 7 h at rt. After quenching with water, the mixture was extracted three times with EtOAc. The combined organic layers were then washed with brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The solvent was evaporated in vacuo and the resulting crude solid was grossly purified by column chromatography (cyclohexane/EtOAc, 4/1), leading to a white solid (0.059 g; quantitative yield), which was directly engaged in the next step.

**Phenyl 6-O-trisopropylsilyl-3,4-O-(2',3'-dimethoxybutane-2',3'-diyl)-2-O-toluenesulfonyl-1-thio- $\alpha$ -D-mannopyranoside (3)**

A suspension of **2**, silver oxide (0.104 g, 0.448 mmol), potassium iodide (0.074 g, 0.448 mmol), and *p*-toluenesulfonyl chloride (TsCl) (0.085 g, 0.448 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was refluxed for 11 h, then cooled down to rt. The mixture was filtered with Celite. The residue was evaporated in vacuo and then purified by column chromatography (cyclohexane/EtOAc, 4/1), yielding a colorless oil (0.062 g, 79% yield).  $[\alpha]_D^{24} +113.1$  (c 1.23, CHCl<sub>3</sub>). <sup>1</sup>H NMR: 7.83 (d, *J* = 8.3, 2H, Ph), 7.44–7.46 (m, 2H, Ph), 7.27–7.29 (m, 3H, Ph), 5.55 (br.s, 1H, H<sub>1</sub>), 4.88 (m, 1H, H<sub>2</sub>), 4.15 (br.dt, *J* = 3.4 and 10.1 Hz, 1H, H<sub>5</sub>), 4.04 (dd, *J* = 10.0 and 10.2 Hz, 1H, H<sub>4</sub>), 3.95 (dd, *J* = 3.0 and 10.0 Hz, 1H, H<sub>3</sub>), 3.90–3.91 (m, 2H, H<sub>6</sub>), 3.20 (s, 3H, OCH<sub>3</sub>), 3.14 (s, 3H, OCH<sub>3</sub>), 2.42 (s, 3H, PhCH<sub>3</sub>), 1.25 (s, 3H, CH<sub>3</sub>), 1.07 (s, 3H, Me), 1.03 (s, 21H, TIPS). <sup>13</sup>C NMR: 144.3, 133.9, 133.7, 131.7, 129.3, 129.0, 128.2, 127.6, 100.1, 99.6, 86.8, 78.6, 72.9, 66.7, 62.7, 61.6, 21.6, 17.9, 17.8, 17.7, 17.4, 12.3, 11.9. IR (KBr): 3017, 2944, 2867, 1597, 1463, 1370, 1112, 921, 756. HR-MS: calcd for C<sub>34</sub>H<sub>52</sub>O<sub>9</sub>NaSiS<sub>2</sub> ([M + Na]<sup>+</sup>): 719.2714. Found: 719.2734.

**Phenyl 3,4-O-(2',3'-dimethoxybutane-2',3'-diyl)-6-O-toluenesulfonyl-1-thio- $\alpha$ -D-mannopyranoside (4)**

To a stirred solution of **1** (1.878 g, 4.86 mmol) in acetonitrile (50 mL) was added TsCl (1.85 g, 9.72 mmol) and TMEDA (1.5  $\mu$ L, 9.72 mmol). The mixture was then stirred at 1.5 h at rt, poured into ice water, and then extracted with methylene chloride. The organic layer was dried (MgSO<sub>4</sub>) and evaporated under vacuum. The residue was chromatographed on silica gel using EtOAc/cyclohexane 3/2 as eluent giving **4** (2.556 g, 2.63 mmol; 97%).  $[\alpha]_D^{25} +216.6$  (c 2.19, CHCl<sub>3</sub>). IR (neat): 3465, 3015, 2951, 2835, 1638, 1599, 1366, 1177, 755. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): 7.7 (d, *J* = 8.1 Hz, 2H, Har), 7.41–7.37 (m, 2H, Har), 7.30–7.24 (m, 5H, Har), 5.37 (br.s, 1H, H<sub>1</sub>), 4.40 (m, 1H, H<sub>5</sub>), 4.26–4.01 (m, 4H, H<sub>2</sub>, H<sub>3</sub>, H<sub>4</sub>, H<sub>6</sub>), 3.94 (dd, *J* = 2.8 and 11.1 Hz, 1H, H<sub>6</sub>), 3.29 (s, 3H, -OCH<sub>3</sub>), 3.22 (s, 3H, -OCH<sub>3</sub>), 2.49 (d, *J* = 1.8 Hz, 1H, -OH), 2.41 (s, 3H, -PhCH<sub>3</sub>), 1.31 (s, 3H, CH<sub>3</sub>), 1.29 (s, 3H, CH<sub>3</sub>). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): 144.5 (1C, Cq), 133.4 (1C, Cq), 132.7 (1C, Cq), 131.7 (1C, CHar), 129.5 (1C, CHar), 128.9 (1C, CHar), 127.9 (1C, CHar), 127.5 (1C, CHar), 100.4 (1C, Cq), 100.0 (1C, Cq), 87.9 (1C, C1), 70.8 (1C), 69.2 (1C), 68.6 (1C), 67.6 (1C), 62.9 (1C), 48.1 (1C, -OCH<sub>3</sub>), 48.0 (1C, -OCH<sub>3</sub>), 21.5 (PhMe), 17.6 (1C, -CH<sub>3</sub>), 17.5 (1C, -CH<sub>3</sub>). HR-MS: calcd for C<sub>25</sub>H<sub>32</sub>O<sub>9</sub>NaS<sub>2</sub> ([M + Na]<sup>+</sup>): 563.1379. Found: 563.1393.



**Phenyl 6-azido-6-deoxy-3,4-O-(2',3'-dimethoxybutane-2',3'-diyl)-1-thio- $\alpha$ -D-mannopyranoside (5)**

To a solution of **4** (125 mg, 0.23 mmol) in DMF (3 mL) was added at rt sodium azide (45 mg, 0.69 mmol). The resulting mixture was heated at 80°C for 40 h. The mixture was then poured in ice water and extracted three times with EtOAc. The organic solvent was dried (MgSO<sub>4</sub>) and evaporated under vacuum. The residue was chromatographed on silica gel with EtOAc/cyclohexane 1/2 as eluent giving **5** (93 mg, 0.22 mmol; 98%).  $[\alpha]_D^{25} +249$  (c 1.62, CHCl<sub>3</sub>). IR (neat): 3447, 2993, 2948, 2833, 2100, 1583, 1480, 1376, 1281, 1140, 1049, 753. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): 7.48–7.44 (m, 2H, Har), 7.30 = 7.21 (m, 3H, Har), 5.53 (d, *J* = 0.5 Hz, 1H, H<sub>1</sub>), 4.40 (m, 1H, H<sub>5</sub>), 4.20 (m, 1H, H<sub>2</sub>), 4.05 (dd, *J* = 9.8 and 10.1 Hz, 1H, H<sub>4</sub>), 3.96 (dd, *J* = 2.8 and 10.1 Hz, 1H, H<sub>3</sub>), 3.44 (s, 1H, OH), 3.42–3.36 (m, 2H, H<sub>6</sub>), 3.27 (s, 3H, -OCH<sub>3</sub>), 3.19 (s, 3H, -OCH<sub>3</sub>), 2.49 (d, *J* = 1.8 Hz, 1H, -OH), 2.41 (s, 3H, -PhCH<sub>3</sub>), 1.32 (s, 3H, CH<sub>3</sub>), 1.27 (s, 3H, CH<sub>3</sub>). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): 133.6 (1C, Cq), 131.6 (2C, CHar), 129 (2C, CHar), 127.5 (1C, CHar), 100.4 (1C, Cq), 100.0 (1C, Cq), 88.1 (1C, C<sub>1</sub>), 71.0 (1C), 70.9 (1C), 68.7 (1C), 64.3 (1C), 50.6 (1C, C<sub>6</sub>), 48.1 (1C, -OCH<sub>3</sub>), 47.9 (1C, -OCH<sub>3</sub>), 17.7 (1C, -CH<sub>3</sub>), 17.5 (1C, -CH<sub>3</sub>). HR-MS: calcd for C<sub>18</sub>H<sub>25</sub>O<sub>6</sub>NaS ([M + Na]<sup>+</sup>): 434.1356. Found: 434.1359.

**Phenyl 6-azido-6-deoxy-3,4-O-(2',3'-dimethoxybutane-2',3'-diyl)-2-O-methylsulfonyl-1-thio- $\alpha$ -D-mannopyranoside (6a)**

To a stirred solution of alcohol **5** (112 mg, 0.27 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 mL) was added MsCl (0.33  $\mu$ L, 3.99 mmol) and triethylamine (0.57 mL, 4.1 mmol). The mixture was stirred for 1 day and then quenched with ice water. The water layer was extracted with dichloromethane. The organic layers were dried (MgSO<sub>4</sub>) filtrated and the solvent evaporated in vacuo. The product was purified by flash chromatography, using cyclohexane/EtOAc 3/1 as eluent giving **6a** (112 mg, 0.23 mmol; 84%).  $[\alpha]_D^{25} +162.5$  (c 2.00, CHCl<sub>3</sub>). IR (neat): 2994, 2948, 2835, 2101, 1583, 1364, 1177, 1142, 1109, 923. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): 7.50–7.46 (m, 2H, Har), 7.35–7.29 (m, 3H, Har), 5.58 (d, *J* = 1.0 Hz, 1H, H<sub>1</sub>), 5.09 (dd, *J* = 1.5 and 2.7 Hz, 1H, H<sub>2</sub>), 4.38 (m, 1H, H<sub>5</sub>), 4.11 (dd, *J* = 2.7 and 10.3 Hz, 1H, H<sub>3</sub>), 3.99 (dd, *J* = 10.0 and 10.3 Hz, 1H, H<sub>4</sub>), 3.55–3.40 (m, 2H, H<sub>6</sub>), 3.31 (s, 3H), 3.24 (s, 3H), 3.16 (s, 3H), 1.29 (s, 6H, CH<sub>3</sub>). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): 132.5 (1C, Cq), 132.2 (2C, CHar), 129.3 (2C, CHar), 128.3 (1C, CHar), 100.5 (1C, Cq), 100.0 (1C, Cq), 87.5 (1C, C<sub>1</sub>), 79.4 (1C), 71.3 (1C), 67.0 (1C), 64.1 (1C), 50.4 (1C, C<sub>6</sub>), 48.4 (1C, -OCH<sub>3</sub>), 48.2 (1C, -OCH<sub>3</sub>), 39.0 (1C, -SO<sub>2</sub>CH<sub>3</sub>), 17.6 (1C, -CH<sub>3</sub>), 17.5 (1C, -CH<sub>3</sub>). HR-MS: calcd for C<sub>19</sub>H<sub>27</sub>O<sub>8</sub>NaS<sub>2</sub> ([M + Na]<sup>+</sup>): 512.1131. Found: 512.1107.

**Phenyl 6-azido-6-deoxy-3,4-O-(2',3'-dimethoxybutane-2',3'-diyl)-2-O-toluenesulfonyl-1-thio- $\alpha$ -D-mannopyranoside (6b)**

To a stirred solution of alcohol **5** (49 mg, 0.12 mmol) in  $\text{CH}_2\text{Cl}_2$  (2 mL) was added  $\text{Ag}_2\text{O}$  (110 mg, 0.48 mmol),  $\text{TsCl}$  (91 mg, 0.48 mmol), and  $\text{KI}$  (79 mg, 0.48 mmol). The mixture was refluxed for 48 h. After cooling at rt, the mixture was filtered and a large excess of TMEDA was added. After stirring for 10 min, the solvent was evaporated in vacuo. The product was purified by flash chromatography, using cyclohexane/EtOAc 4/1 as eluent giving **6b** (65 mg, 0.11 mmol; 96%). IR (neat): 3060, 2993, 2952, 2835, 2101, 1738, 1597, 1444, 1370, 918.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ): 7.84–7.80 (m,  $J = 8.3$  Hz, 2H, Har), 7.47–7.34 (m, 2H, Har), 7.34–7.25 (m, 5H, Har), 5.57 (d,  $J = 1.0$  Hz, 1H,  $\text{H}_1$ ), 4.86 (dd,  $J = 1.4$  and 2.4 Hz, 1H,  $\text{H}_2$ ), 4.29 (m, 1H,  $\text{H}_5$ ), 3.94–3.90 (m, 2H,  $\text{H}_3$  and  $\text{H}_4$ ), 4.28 (dd,  $J = 2.6$  and 13.2 Hz, 1H,  $\text{H}_6$ ), 4.19 (dd,  $J = 6.4$  and 13.2 Hz, 1H,  $\text{H}_6$ ), 3.18 (s, 3H,  $-\text{OCH}_3$ ), 3.11 (s, 3H,  $-\text{OCH}_3$ ), 2.02 (s, 3H,  $-\text{C}_6\text{H}_4\text{-CH}_3$ ), 1.22 (s, 3H,  $\text{CH}_3$ ), 1.04 (s, 3H,  $\text{CH}_3$ ).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ): 144.6 (1C, Cq), 133.6 (1C, Cq), 132.7 (1C, Cq), 132.1, 129.4, 129.3, 128.3, and 128.2 (CHar), 100.2 (1C, Cq), 99.8 (1C, Cq), 87.2 (1C,  $\text{C}_1$ ), 78.5 (1C), 71.3 (1C), 66.3 (1C), 64.1 (1C), 50.5 (1C,  $\text{C}_6$ ), 48.0 (2C,  $-\text{OCH}_3$ ), 21.6 (1C, Ar-Me), 17.6 (1C,  $-\text{CH}_3$ ), 17.3 (1C,  $-\text{CH}_3$ ). HR-MS: calcd for  $\text{C}_{25}\text{H}_{31}\text{O}_8\text{NaS}_2$  ( $[\text{M} + \text{Na}]^+$ ): 588.1444. Found 588.1401.

**Procedure for 1,2-Thio Migration/Azidation**

To a solution of thiophenyl  $\cdot \alpha$ -D-mannoside (1 eq.) in DMF (0.05 M) was added at rt sodium azide (3 eq.). The resulting mixture was heated at  $130^\circ\text{C}$  for several hours until disappearance of the starting material as judged from TLC. The mixture was then poured in ice water and extracted three times with  $\text{CH}_2\text{Cl}_2$ . The combined organic layers were washed with water and brine, dried over  $\text{MgSO}_4$ , and concentrated in vacuum. The residue was chromatographed on silica gel with EtOAc/cyclohexane 7/1 as eluent.

**2-Deoxy-6-O-triisopropylsilyl-3,4-O-(2',3'-dimethoxybutane-2',3'-diyl)-2-thiophenyl-D-mannopyranosyl azide (7)**

(Mixture  $\alpha/\beta$ )  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ): 7.60–7.53 (m, 2H, Har), 7.29–7.24 (m, 3H, Har), 5.38 (d,  $J = 4.0$  Hz, 1H,  $\text{H}_1$  minor), 4.58 (d,  $J = 9.5$  Hz, 1H,  $\text{H}_1$  major), 3.95–3.89 (m, two  $\text{H}_6$  major;  $\text{H}_3$ ,  $\text{H}_4$ ,  $\text{H}_5$ , and  $\text{H}_6$  minor), 3.82 (dd,  $J = 9.6$  and 9.9 Hz, 1H,  $\text{H}_4$  major), 3.67 (dd,  $J = 9.6$  and 11.4 Hz, 1H,  $\text{H}_3$  major), 3.48 (ddd,  $J = 2.2$ , 3.1 and 9.5 Hz, 1H,  $\text{H}_5$  major), 3.38 (s, 3H,  $-\text{OCH}_3$  major), 3.35 (s, 3H,  $-\text{OCH}_3$  minor), 3.27 (dd,  $J = 4.0$  and 11.2 Hz, 1H,  $\text{H}_2$

minor), 3.28 (s, 3H, -OCH<sub>3</sub> major), 3.26 (s, 3H, -OCH<sub>3</sub> minor), 2.97 (dd,  $J = 9.5$  and 11.4 Hz, 1H, H<sub>2</sub> major), 1.35 (s, 3H, CH<sub>3</sub>), 1.30 (s, 3H, CH<sub>3</sub>), 1.05 (s, 21H, TIPS).

### 2-Deoxy-3,4-O-(2',3'-dimethoxybutane-2',3'-diyl)-2-thiophenyl- $\beta$ -mannopyranosyl azide (9)

(Mixture  $\alpha/\beta$ ) <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): 7.60–7.52 (m, 2H, Har), 7.29–7.24 (m, 3H, Har), 5.38 (d,  $J = 4.0$  Hz, 1H, H<sub>1</sub> minor), 4.61 (d,  $J = 9.6$  Hz, 1H, H<sub>1</sub> major), 4.05–3.68 (m, H<sub>6</sub>, H<sub>4</sub>, and H<sub>3</sub> of both anomers, H<sub>5</sub> minor), 3.55 (m, 1H, H<sub>5</sub> major), 3.38 (s, 3H, -OCH<sub>3</sub> major), 3.35 (s, 3H, -OCH<sub>3</sub> minor), 3.30 (dd,  $J = 4.0$  and 11.3 Hz, 1H, H<sub>2</sub> minor), 3.26 (s, 3H, -OCH<sub>3</sub>), 2.98 (dd,  $J = 9.6$  and 11.2 Hz, 1H, H<sub>2</sub> major), 1.34 (s, 3H, CH<sub>3</sub> major), 1.31 (s, 3H, CH<sub>3</sub> minor), 1.30 (s, 6H, CH<sub>3</sub> major and minor).

### 6-Azido-2,6-dideoxy-3,4-O-(2',3'-dimethoxybutane-2',3'-diyl)-2-thiophenyl- $\alpha$ - $\beta$ -mannopyranosyl azide (10 $\alpha$ )

$[\alpha]_D^{24} +268$  (c 0.51, CHCl<sub>3</sub>). IR (neat): 2993, 2948, 2833, 2107, 1440, 1379, and 1136. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): 7.55 (m, 2H, Har), 7.29 (m, 3H, Har), 5.41 (d,  $J = 4.2$  Hz, 1H, H<sub>1</sub>), 4.09 (m, 1H, H<sub>5</sub>), 3.92 (dd,  $J = 9.6$  and 11.5 Hz, 1H, H<sub>3</sub>), 3.71 (dd,  $J = 9.6$  and 9.8 Hz, 1H, H<sub>4</sub>), 3.58 (dd,  $J = 2.5$  and 13.3 Hz, 1H, H<sub>6</sub>), 3.40 (dd,  $J = 4.8$  and 13.3 Hz, 1H, H<sub>6</sub>), 3.35 (s, 3H, -OCH<sub>3</sub>), 3.29 (dd,  $J = 4.2$  and 11.5 Hz, 1H, H<sub>2</sub>), 3.26 (s, 3H, -OCH<sub>3</sub>), 1.31 (s, 3H, CH<sub>3</sub>), 1.29 (s, 3H, CH<sub>3</sub>). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): 133.9 (1C, Cq ar), 132.7 (2C, CHar), 128.9 (2C, CHar), 127.7 (1C, CHar), 100.2 (1C, Cq), 100.1 (1C, Cq), 90.1 (1C, C<sub>1</sub>), 71.3 (1C, C<sub>5</sub>), 68.2 (1C, C<sub>3</sub>), 68.1 (1C, C<sub>4</sub>), 51.7 (1C, C<sub>2</sub>), 50.1 (1C, C<sub>6</sub>), 48.2 (1C, -OCH<sub>3</sub>), 48.1 (1C, -OCH<sub>3</sub>), 17.6 (1C, -CH<sub>3</sub>), 17.5 (1C, -CH<sub>3</sub>). HR-MS: calcd for C<sub>18</sub>H<sub>24</sub>N<sub>6</sub>O<sub>5</sub>NaS ([M + Na]<sup>+</sup>): 459.1426. Found: 459.1413.

### 6-Azido-2,6-dideoxy-3,4-O-(2',3'-dimethoxybutane-2',3'-diyl)-2-thiophenyl- $\beta$ - $\beta$ -mannopyranosyl azide (10 $\beta$ )

$[\alpha]_D^{24} +116$  (c 0.52, CHCl<sub>3</sub>). IR (neat): 2995, 2948, 2836, 2116, 1441, 1376, and 1136. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): 7.58 (m, 2H, Har), 7.28 (m, 3H, Har), 4.59 (d,  $J = 9.6$  Hz, 1H, H<sub>1</sub>), 3.64 (m, 4H, H<sub>3</sub>, H<sub>4</sub>, H<sub>5</sub>, H<sub>6</sub>), 3.37 (m, 1H, H<sub>6</sub>), 3.38 (s, 3H, -OCH<sub>3</sub>), 3.26 (s, 3H, -OCH<sub>3</sub>), 3.00 (dd,  $J = 9.6$  and 10.8 Hz, 1H, H<sub>2</sub>), 1.34 (s, 3H, CH<sub>3</sub>), 1.29 (s, 3H, CH<sub>3</sub>). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): 133.6 (1C, Cq ar), 133.2 (2C, CHar), 128.9 (2C, CHar), 127.8 (1C, CHar), 100.4 (1C, Cq), 99.9 (1C, Cq), 91.5 (1C, C<sub>1</sub>), 75.4 (1C, C<sub>3</sub>), 70.5 (1C, C<sub>4</sub>), 67.8 (1C, C<sub>5</sub>), 53.5 (1C, C<sub>2</sub>), 50.2 (1C, C<sub>6</sub>), 48.4 (1C, -OCH<sub>3</sub>), 48.1 (1C, -OCH<sub>3</sub>), 17.6

(1C, -CH<sub>3</sub>), 17.5 (1C, -CH<sub>3</sub>). HR-MS: calcd for C<sub>18</sub>H<sub>24</sub>N<sub>6</sub>O<sub>5</sub>NaS ([M + Na]<sup>+</sup>): 459.1426. Found: 459.1445.

## ACKNOWLEDGMENT

The authors thank the CNRS for financial support. C.M. thanks the ANRS for financial support; A.K. thanks the Science University of Tokyo for fellowship.

## REFERENCES

- [1] Dwek, R.A. Glycobiology: toward understanding the function of sugars. *Chem. Rev.* **1996**, *96*, 683–720.
- [2] Ernst, B.; Hart, G.; Sinay, P. *Carbohydrates in Chemistry and Biology*; Ernst, B., Hart, G., Sinay, P., Eds.; Wiley-VCH: Weinheim, 2002.
- [3] Wertz, D.B.; Seeberger, P.H. Carbohydrates as the next frontier in pharmaceutical research. *Chem. Eur. J.* **2005**, *11*, 3194–3206.
- [4] Schweizer, F. Glycosamino acids: building blocks for combinatorial synthesis—implications for drug discovery. *Angew. Chem. Int. Ed.* **2002**, *41*, 230–253.
- [5] Seitz, O. Glycopeptide synthesis and the effects of glycosylation on protein structure and activity. *Chem. Bio. Chem* **2000**, *1*, 214–246.
- [6] Goebel, M.; Ugi, I. Beyond peptide and nucleic acid combinatorial libraries—applying unions of multicomponent reactions towards the generation of carbohydrate combinatorial libraries. *Tetrahedron Lett.* **1995**, *36*, 6043–6046.
- [7] Fox, D.J.; House, D.; Warren, S. Mechanisms of sulfanyl (RS) migrations: synthesis of heterocycles. *Angew. Chem., Int. Ed.* **2002**, *41*, 2462–2482.
- [8] Acton, E.M.; Ryan, K.J.; Goodman, L. Synthesis of 2-thio-D-ribose and 2'-thioadenosine derivatives. *J. Org. Chem.* **1971**, *36*, 2646–2657.
- [9] Nicolaou, K.C.; Laduwahetty, T.; Randall, J.L.; Chcucholowski, A. Stereospecific 1,2-migrations in carbohydrates. Stereocontrolled synthesis of  $\alpha$ - and  $\beta$ -2-deoxyglycosides. *J. Am. Chem. Soc.* **1986**, *108*, 2466–2467.
- [10] Zuurmond, H.M.; van der Klein, P.A.M.; van der Marel, G.A.; van Boom, J.H.A. stereospecific approach towards the synthesis of 2-deoxy  $\alpha$ - and  $\beta$ -glycosides based on a 1,2-ethyl (phenyl) thio group migration. *Tetrahedron* **1993**, *49*, 6501–6514.
- [11] Liu, B.; Yang, Z. Stereoselective synthesis of 2-S-phenyl-2-deoxy- $\beta$ -glycosides using phenyl 2,3-O-thionocarbonyl-1-thioglycoside donors via 1,2-migration and concurrent glycosidation. *Org. Lett.* **2001**, *3*, 377–379.
- [12] Lipták, A.; Sajtos, F.; Jánosy, L.; Gehle, D.; Szilágyi, L. A general method for the synthesis of sugar 2-C-sulfonic acids by 1, 2-arylthio group migration in acid-sensitive thioglycosides. Direct transformation of thiotrityl ethers into C-sulfonic acids. *Org. Lett.* **2003**, *5*, 3671–3674.
- [13] Johnston, B.D.; Pinto, B.M. Synthesis of thio-linked disaccharides by 1,2-intramolecular thioglycosyl migration: oxacarbenium versus episulfonium ion intermediates. *J. Org. Chem.* **2000**, *65*, 4607–4617.
- [14] Nicolaou, K.C.; Fylaktakidou, K.; Monenschein, H.; Li, Y.; Weyershausen, B.; Mitchell, H.J.; Wei, H.; Guntupalli, P.; Hepworth, D.; Sugita, K. Total synthesis

- of apoptolidin: construction of enantiomerically pure fragments. *J. Am. Chem. Soc.* **2003**, *125*, 15443–15452.
- [15] Auzanneau, F.-I.; Bundle, D.R. Incidence and avoidance of stereospecific 1,2-ethylthio group migration during the synthesis of ethyl 1-thio- $\alpha$ -L-rhamnopyranoside 2,3-orthoester. *Carbohydr. Res.* **1991**, *212*, 13–24.
- [16] Pozsgay, V. A simple method for avoiding alkylthio group migration during the synthesis of thioglycoside 2,3-orthoesters. An improved synthesis of partially acylated 1-thio- $\alpha$ -L-rhamnopyranosides. *Carbohydr. Res.* **1992**, *235*, 295–302.
- [17] Gadikota, R.R.; Callam, C.S.; Wagner, T.; Del Fraino, B.; Lowary, T.L. 2,3-Anhydro sugars in glycoside bond synthesis. Highly stereoselective syntheses of oligosaccharides containing  $\alpha$ - and  $\beta$ -arabinofuranosyl linkages. *J. Am. Chem. Soc.* **2003**, *125*, 4155–4165.
- [18] Crich, D.; Hutton, T.K.; Banerjee, A.; Jayalath, P.; Picione, J. Disarming, non-participating 2-*O*-protecting groups in manno- and rhamnopyranosylation: scope and limitations of sulfonates, vinylogous esters, phosphates, cyanates, and nitrates. *Tetrahedron: Asymmetry* **2005**, *16*, 105–119.
- [19] Crich, D.; Cai, W.; Dai, Z. Highly diastereoselective  $\alpha$ -mannopyranosylation in the absence of participating protecting groups. *J. Org. Chem.* **2000**, *65*, 1291–1297.
- [20] Hense, A.; Ley, S.V.; Osborn, H.M.I.; Owen, D.R.; Poisson, J.-F.; Warriner, S.L.; Wesson, K.E. Direct preparation of diacetals from 1,2-diketones and their use as 1,2-diol protecting groups. *J. Chem. Soc. Perkin Trans. 1* **1997**, 2023–2030.
- [21] Ogilvie, K.K.; Thompson, E.A.; Quilliam, M.; Westmore, J.B. Selective protection of hydroxyl groups in deoxynucleosides using alkylsilyl reagents. *Tetrahedron Lett.* **1974**, 2865–2868.
- [22] Bouzide, A.; Berre, N.; Sauvé, G. Silver(I) oxide-mediated facile and practical sulfonylation of alcohols. *Tetrahedron Lett.* **2001**, *42*, 8781–8783.
- [23] Yoshida, Y.; Shimonishi, K.; Sakakura, Y.; Okada, S.; Aso, N.; Tanabe, Y. Facile and practical methods for the sulfonylation of alcohols using Ts(Ms)Cl and Me<sub>2</sub>N(CH<sub>2</sub>)<sub>n</sub>NMe<sub>2</sub> as a key base. *Synthesis* **1999**, 1633–1636.