

¹H NMR SPECTRA OF SOME ARYLTELLURIUM COMPOUNDS

FRANK J. BERRY, EDWARD H. KUSTAN, MINA ROSHANI and BARRY C. SMITH*

*Department of Chemistry, Birkbeck College, University of London, Malet Street,
 London WC1E 7HX (Great Britain)*

(Received May 7th, 1975)

Summary

The ¹H NMR spectra of some anisyl- and phenetyl-tellurium derivatives confirm that *para*-substitution of methoxybenzene (anisole) and ethoxybenzene (phenetole) occurs on reaction with tellurium tetrachloride. Some new aryltellurium(IV) halides are reported.

Introduction

Phenetole [1] and anisole [2] undergo quasi-Friedel–Crafts reactions with tellurium tetrachloride in boiling organic solvents to form yellow aryltellurium(IV) trichlorides. Further reactions at higher temperatures and in the absence of solvent give diaryltellurium(IV) dichlorides [1,2]. The trichlorides and dichlorides react with stoichiometric quantities of silver fluoride to form aryltellurium(IV) trifluorides and diaryltellurium(IV) difluorides.

Hydrolysis of the trichlorides gives aryltellurium(IV) oxochlorides [3,4], oxohydroxides [3] or tellurinic anhydrides [4], and of the dichlorides gives diaryltellurium(IV) oxochlorides, dihydroxides, or telluroxides [5–7]. The hydrolysis products are converted into trihalides and dihalides on reaction with aqueous HCl, HBr, and HI.

Phenyl- and *p*-tolyl-tellurium(IV) trichlorides are prepared from arylmercury(II) chlorides and tellurium tetrachloride [8]. The reduction of aryltellurium(IV) trichlorides by aqueous K₂S₂O₅ gives red-brown diaryl ditellurides(II) [1,2,8] which react with an excess of bromine or iodine to give the aryltellurium(IV) trihalides [9,10]. Dianisyl ditelluride(II) reacts with methyl iodide to give anisyl dimethyltellurium(IV) iodide and anisyl methyltellurium(IV) diiodide [1], and it is shown below that the other ditellurides behave similarly:



Reduction of dianisyl- and diphenetyl-tellurium(IV) dichlorides with zinc dust in boiling benzene [2,7] and reactions of phenyl- and *p*-tolyl-mercury(II)

TABLE 1

¹H NMR CHEMICAL SHIFTS FOR ARYL PROTONS IN DMSO

Ar compound	<i>p</i> -MeC ₆ H ₄		<i>p</i> -MeOC ₆ H ₄		<i>p</i> -EtOC ₆ H ₄	
	δA	δB	δA	δB	δA	δB
Ar ₂ Te	7.51	7.03	7.75	6.98	7.73	6.98
Ar ₂ Te ₂	7.71	7.09	7.86	6.99	7.64	6.78
Ar ₂ TeF ₂			7.93	7.26	7.92	7.23
Ar ₂ TeCl ₂	7.83	7.32	7.98	7.17	7.88	7.09
Ar ₂ TeBr ₂	7.88	7.29	8.14	7.24	8.10	7.21
Ar ₂ TeI ₂	7.89	7.24	8.87	7.07	8.05	7.16
ArTeF ₃			8.08	7.26	8.20	7.20
ArTeCl ₃	8.09	7.18	8.35	7.16	8.32	7.13
ArTeBr ₃	8.23	7.15	8.50	7.01	8.60	7.20
ArTeI ₃	8.19	7.41	8.26	6.85	8.22	6.91
ArMe ₂ TeI	7.93	7.56	7.80	7.11	7.80	7.05
ArMeTeI ₂	7.83	6.67	8.00	7.08	7.92	7.01

with elemental tellurium [11,12] give diaryl tellurides(II) which react with chlorine, bromine, and iodine to give diaryltellurium(IV) dihalides [5,7,11-16].

Results and discussion

This work confirms that substitution of anisole and phenetole by tellurium(IV) occurs exclusively at *para* positions. The ¹H NMR spectra of the substitution products show four line AABB signals characteristic of 1,4-disubstituted benzenes, *J*_{AB} 8–9 Hz, which resemble those of corresponding *p*-tolyl derivatives. The protons in *ortho* positions to tellurium appear as broad doublets at lower field.

Chemical shifts with respect to TMS, of the aryl protons of some typical compounds in DMSO-*d*₆ are recorded in Table 1. All the aryltellurium compounds are soluble in DMSO; some are soluble in CCl₄ and C₆H₆. The differences in chemical shifts, δA – δB, increase in the order DMSO < CCl₄ < C₆H₆, and for a particular solvent, in the order Ar₂Te < Ar₂TeX₂ < ArTeX₃.

Experimental

Methods of preparation of typical new organotellurium compounds are described below. Analytical data for new compounds are summarised in Table 2. Other organotellurium compounds were prepared by standard methods [1-16].

Di-*p*-phenetyltellurium(IV) difluoride. Di-*p*-phenetyltellurium(IV) dichloride (0.88 g, 1.99 mmol) was added in small quantities to a mixture of silver fluoride (0.5 g, 4.0 mmol) in dry toluene (30 ml) which was boiled under reflux (1 h). Silver chloride (0.51 g, 3.55 mmol) was removed by filtration. The addition of the cold, clear, concentrated filtrate to light petroleum (b.p. 60-80°C) gave colourless crystalline di-*p*-phenetyltellurium(IV) difluoride (0.13 g, 0.32 mmol), m.p. 169-170°C.

Dimethyl-*p*-phenetyltellurium(IV) iodide and methyl-*p*-phenetyltellurium-

TABLE 2
ANALYTICAL DATA FOR NEW COMPOUNDS

Compound	m.p. (°C)	Found (%)			Molecular formula	Calcd. (%)		
		C	H	Hal		C	H	Hal
<i>p</i> -MeOC ₆ H ₄ TeF ₃	176-178	29.22	2.59		C ₇ H ₇ F ₃ OTe	28.84	2.46	19.54
(<i>p</i> -MeOC ₆ H ₄) ₂ TeF ₂	120-121	42.73	3.58		C ₁₄ H ₁₄ F ₂ O ₂ Te	44.30	3.69	10.01
<i>p</i> -EtOC ₆ H ₄ TeF ₃	204	30.92	2.91		C ₈ H ₉ F ₃ OTe	31.44	2.95	18.65
(<i>p</i> -EtOC ₆ H ₄) ₂ TeF ₂	169-170	46.92	4.28		C ₁₆ H ₁₈ F ₂ O ₂ Te	47.15	4.41	9.32
PhMe ₂ TeI	168	26.45	3.09	35.32	C ₈ H ₁₁ I ₂ Te	26.57	3.04	35.09
PhMeTeI ₂	145	18.36	1.79	51.95	C ₇ H ₈ I ₂ Te	17.76	1.69	53.60
<i>p</i> -TolMe ₂ TeI	154	28.79	3.62	32.12	C ₉ H ₁₃ I ₂ Te	28.78	3.46	33.79
<i>p</i> -TolMeTeI ₂	126-127	20.11	2.15	52.14	C ₈ H ₁₀ I ₂ Te	19.71	2.07	52.05
<i>p</i> -EtOC ₆ H ₄ Me ₂ TeI	163	29.06	3.63	32.29	C ₁₀ H ₁₅ I ₂ OTe	29.62	3.70	31.29
<i>p</i> -EtOC ₆ H ₄ MeTeI ₂	108	21.47	2.43	50.07	C ₉ H ₁₂ I ₂ OTe	20.89	2.32	49.05

(IV) diiodide. Di-*p*-phenetyl ditelluride(II) (1 g, 1.98 mmol) was boiled under reflux with an excess of methyl iodide (50 g). The yellow microcrystalline precipitate was removed by filtration and washed with benzene (2 × 15 ml) to give yellow dimethyl-*p*-phenetyltellurium(IV) iodide (0.89 g, 2.19 mmol), m.p. 163°C. Addition of the cold red filtrate to light petroleum (b.p. 60-80°C) (30 ml) gave red needles of methyl-*p*-phenetyltellurium(IV) diiodide (0.34 g, 0.66 mmol), m.p. 108°C.

References

- 1 G.T. Morgan and H.D.K. Drew, J. Chem. Soc., (1925) 2307.
- 2 G.T. Morgan and R.E. Kellett, J. Chem. Soc., (1926) 1080.
- 3 L. Reichel and E. Kirschbaum, Ann., 523 (1936) 211.
- 4 N. Petragani and G. Vicentini, Univ. São Paulo, Fre. Filosoff. Cienc. Letras. Bol. Comm., 5 (1959) 75.
- 5 K. Lederer, Ann., 391 (1912) 326.
- 6 H. Rheinboldt and E. Giesbrecht, J. Amer. Chem. Soc., 69 (1947) 2310.
- 7 G.T. Morgan and F.H. Burstall, J. Chem. Soc., (1930) 2599.
- 8 W.V. Farrar, Research London, 4 (1951) 177.
- 9 W.R. McWhinnie and P. Thavornnyutikarn, J. Chem. Soc. Dalton Trans., (1972) 551.
- 10 N. Petragani, Tetrahedron, 11 (1960) 11.
- 11 F. Krafft and R.E. Lyons, Ber., 27 (1894) 1768.
- 12 F. Zeiser, Ber., 28 (1895) 1670.
- 13 K. Lederer, Ber., 49 (1916) 1076.
- 14 K. Lederer, Ber., 50 (1917) 238.
- 15 K. Lederer, Ber., 48 (1915) 1345.
- 16 L. Reichel and E. Kirschbaum, Ber., 76B (1943) 1105.